



National Comprehensive
Cancer Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

B-Cell Lymphomas

Version 2.2020 — July 9, 2020

NCCN.org

Continue



Download Clinical Guidelines



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2020

B-Cell Lymphomas

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

Andrew D. Zelenetz, MD, PhD/Chair † ‡
Memorial Sloan Kettering Cancer Center

Leo I. Gordon, MD/Co-Vice Chair ‡ §
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

Jeremy S. Abramson, MD † ‡
Massachusetts General Hospital
Cancer Center

Ranjana H. Advani, MD †
Stanford Cancer Institute

Nancy L. Bartlett, MD †
Siteman Cancer Center at Barnes-
Jewish Hospital and Washington
University School of Medicine

L. Elizabeth Budde, MD, PhD † §
City of Hope National Medical Center

Paolo F. Caimi, MD ‡ †
Case Comprehensive Cancer Center/
University Hospitals Seidman Cancer Center
and Cleveland Clinic Taussig Cancer Institute

Julie E. Chang, MD ‡
University of Wisconsin
Carbone Cancer Center

Julio C. Chavez, MD †
Moffitt Cancer Center

Beth Christian, MD †
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Bita Fakhri, MD †
UCSF Helen Diller Family
Comprehensive Cancer Center

Luis E. Fayad, MD ‡ † †
The University of Texas
MD Anderson Cancer Center

Martha J. Glenn, MD † ‡ † §
Huntsman Cancer Institute
at the University of Utah

Thomas M. Habermann, MD ‡ §
Mayo Clinic Cancer Center

Francisco Hernandez-Ilizaliturri, MD †
Roswell Park Cancer Institute

Mark S. Kaminski, MD †
University of Michigan
Rogel Cancer Center

Christopher R. Kelsey, MD §
Duke Cancer Institute

Nadia Khan, MD †
Fox Chase Cancer Center

Susan Krivacic, MPAff ¥
Consultant

Ann S. LaCasce, MD †
Dana-Farber/Brigham and Women's
Cancer Center

Amitkumar Mehta MD † ‡ †
O'Neal Comprehensive Cancer Center at UAB

Auayporn Nademanee, MD † ‡ §
City of Hope National Medical Center

Mayur Narkhede, MD † ‡
O'Neal Comprehensive Cancer Center at UAB

Rachel Rabinovitch, MD §
University of Colorado Cancer Center

Nishitha Reddy, MD ‡ §
Vanderbilt-Ingram Cancer Center

Erin Reid, MD ‡
UC San Diego Moores Cancer Center

Kenneth B. Roberts, MD §
Yale Cancer Center/Smilow Cancer Hospital

Stephen D. Smith, MD ‡
Fred Hutchinson Cancer Research Center/
Seattle Cancer Care Alliance

Jakub Svoboda, MD ‡
Abramson Cancer Center
at the University of Pennsylvania

Lode J. Swinnen, MB, ChB † ‡ §
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

Julie M. Vose, MD, MBA ‡ §
Fred & Pamela Buffett Cancer Center

NCCN
Mary Dwyer, MS
Hema Sundar, PhD

[NCCN Guidelines Panel Disclosures](#)

Continue

§ Bone marrow transplantation	¥ Patient advocacy
‡ Hematology/Hematology oncology	§ Radiotherapy/Radiation oncology
‡ Internal medicine	* Discussion Writing Committee Member
† Medical oncology	
≠ Pathology	



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2020

B-Cell Lymphomas

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

[NCCN B-Cell Lymphoma Panel Members](#) [Summary of the Guidelines Updates](#)

- [Diagnosis \(DIAG-1\)](#)
- [Follicular Lymphoma \(FOLL-1\)](#)
- [Marginal Zone Lymphomas \(MZL-1\)](#)
 - ▶ [Gastric MALT Lymphoma \(MALT-1\)](#)
 - ▶ [Nongastric MALT Lymphoma \(Noncutaneous\) \(NGMLT-1\)](#)
 - ▶ [Nodal Marginal Zone Lymphoma \(NODE-1\)](#)
 - ▶ [Splenic Marginal Zone Lymphoma \(SPLN-1\)](#)
- [Mantle Cell Lymphoma \(MANT-1\)](#)
- [Diffuse Large B-Cell Lymphoma \(BCEL-1\)](#)
- [High-Grade B-Cell Lymphomas](#)
 - ▶ [High-Grade B-Cell lymphomas with Translocations of MYC and BCL2 and/or BCL6 \(Double-/Triple-Hit Lymphoma\) \(HGBL-1\)](#)
 - ▶ [High-Grade B-Cell Lymphomas, NOS \(HGBLNOS-1\)](#)
- [Burkitt Lymphoma \(BURK-1\)](#)
- [AIDS-Related B-Cell Lymphomas \(AIDS-1\)](#)
- [Lymphoblastic Lymphoma \(BLAST-1\)](#)
- [Post-Transplant Lymphoproliferative Disorders \(PTLD-1\)](#)
- [Castleman's Disease \(CD-1\)](#)
- [Use of Immunophenotyping/Genetic Testing in Differential Diagnosis of Mature B-Cell and NK/T-Cell Neoplasms \(NHODG-A\)](#)
- [Supportive Care for B-Cell Lymphomas \(NHODG-B\)](#)
- [Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#)
- [Principles of Radiation Therapy \(NHODG-D\)](#)
- [Special Considerations for the Use of Small-Molecule Inhibitors \(NHODG-E\)](#)

Clinical Trials: NCCN believes that the best management for any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

To find clinical trials online at NCCN Member Institutions, [click here: nccn.org/clinical_trials/member_institutions.aspx](#).

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

NCCN Categories of Preference: All recommendations are considered appropriate.

See [NCCN Categories of Preference](#).

[Classification and Staging \(ST-1\)](#)

[Primary CNS Lymphoma \(See NCCN Guidelines for CNS\)](#)

[Waldenström Macroglobulinemia/Lymphoplasmacytic Lymphoma \(See NCCN Guidelines for WM/LPL\)](#)

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2020.



Updates in Version 2.2020 of the NCCN Guidelines for B-Cell Lymphomas from Version 1.2020 include:

Follicular Lymphoma

FOLL-8

- Histologic transformation of FL to DLBCL after multiple lines of prior therapies,
 - Selinexor (only after at least two lines of systemic therapy) was added as a category 2A recommendation.

FOLL-B 2 of 4

- Second-line and Subsequent Therapy
 - Other recommended regimens, tazemetostat was added as a category 2A recommendation for the following indications
 - ◊ EZH2 mutation positive relapsed/refractory disease after 2 prior therapies
 - ◊ EZH2 wild type relapsed/refractory disease in patients who have no satisfactory alternative treatment options

Diffuse Large B-Cell Lymphoma

BCEL-C 2 of 4

- A new heading was added, "Third-line and Subsequent Therapy (including patients with disease progression after transplant or CAR T-cell therapy)."
 - "Selinexor (only after at least two lines of systemic therapy)" was added as a category 2A recommendation with corresponding footnote p, "Selinexor is FDA approved only for DLBCL and transformed DLBCL arising from FL".

[Continued](#)



Updates in Version 1.2020 of the NCCN Guidelines for B-Cell Lymphomas from Version 7.2019 include:

Global changes

- Suggested treatment regimen and other references were updated throughout the guidelines.

Follicular Lymphoma

FOLL-1

- Additional Diagnostic Testing
 - ▶ Essential, 2nd sub-bullet was revised, "Cell surface marker analysis by flow cytometry *with peripheral blood and/or biopsy specimen: ...*" (Also throughout the algorithm where applicable.)
 - ▶ Useful, 4th bullet was added, "NGS panel including *TNFRSF14* and *STAT6* mutation."
 - ▶ Footnote a was revised, "Follicular lymphoma (FL), grade 1-2. FL, grade 3b is commonly treated as diffuse large B-cell lymphoma (BCL-1). ~~The management of FL, grade 3a is controversial an area of controversy and management should be individualized. The distinction between follicular grade 3a and 3b has not been shown to have clinical significance to date. However, controversy exists regarding management of FL grade 3. Some may treat FL grade 3a as FL and others may treat it as diffuse large B-cell lymphoma (DLBCL).~~ Any area of DLBCL in an FL of any grade should be diagnosed and treated as a DLBCL."

FOLL-2

- Workup, Essential,
 - ◊ 8th bullet was revised, "Chest/abdominal/pelvic (C/A/P) CT with contrast of diagnostic quality ~~and/or~~" and "Whole-body PET/CT..." was moved to its own bullet.
 - ◊ 9th bullet was revised, "Whole-body PET/CT scan essential if RT for stage I, II disease ~~or systemic therapy is planned.~~"
- ▶ Footnotes
 - ◊ Footnote h was revised, "... For patients with risk factors or previous history of hepatitis B, add e-antigen (*See NHODG-B*). If positive, check viral load and *consider* consult with gastroenterologist." Footnote g was revised by adding, "*See NHODG-B*." Also revised in same footnote through the guidelines.
 - ◊ Footnote i was revised by adding, "Selection of patients requires adequate marrow cellularity >15% and <25% involvement of lymphoma in bone marrow, and platelets >100,000. In patients with prior autologous stem cell rescue, referral to a tertiary care center is highly recommended for ibritumomab tiuxetan."

FOLL-3

- Stage I and contiguous stage II size qualifiers were removed, "<7 cm" and "≥7 cm."
- Stage I or Contiguous Stage II, Initial Therapy,
 - ▶ "ISRT" was listed as "preferred."
 - ▶ "Anti-CD20 monoclonal antibody ± chemotherapy (*See FOLL-B*) + ISRT (category 2B)" was added.
- Non-contiguous Stage II, Initial Therapy,
 - ▶ "± ISRT for local palliation" was added to "Anti-CD20 monoclonal antibody ± chemotherapy (*See FOLL-B*) + ISRT."
 - ▶ "Anti-CD20 monoclonal antibody ± chemotherapy (*See FOLL-B*) + ISRT (category 2B)" was removed.
 - ▶ Response to Therapy
 - ◊ PR or NR was revised, "Consider ISRT *if not previously given*"
- Footnotes
 - ▶ Footnote k was revised, "Observation may be appropriate in circumstances where potential toxicity of involved-site RT (ISRT) or systemic therapy outweighs potential clinical benefit *in consultation with a radiation oncologist.*"
 - ▶ Footnote o was revised by adding, "Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated." (Also on [FOLL-4](#) and [FOLL-5](#))

FOLL-4

- Header was revised, "Initial Management and Follow-Up" (Also for [NODE-3](#))
- Footnote t was revised by adding, "Perform FISH for BCL2..."
- Footnote v was added, "NGS may be useful for treatment selection."

FOLL-7

- Histologic transformation to DLBCL, Response to Therapy,
 - ▶ For PR, NR or progressive disease, option was revised, "Axicabtagene ciloleucel or tisagenlecleucel *Anti-CD19 CAR T-cell Therapy* (only after ≥2 prior chemoimmunotherapy regimens)" and listed as a preferred option. Axicabtagene ciloleucel or tisagenlecleucel were added to FOLL-B. (Also for FOLL-8)
 - ▶ NR or progressive disease was added, "Polatuzumab vedotin ± bendamustine ± rituximab." (Also for FOLL-8)
- Footnotes
 - ▶ Footnote o was added "Imaging should be performed whenever there are clinical indications. For surveillance imaging, see Discussion for consensus imaging recommendations. Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated." (Also for [FOLL-8](#) and [FOLL-9](#))

[Continued](#)

UPDATES



Updates in Version 1.2020 of the NCCN Guidelines for B-Cell Lymphomas from Version 7.2019 include:

FOLL-8

- Histologic transformation to DLBCL after multiple lines of prior therapies,
 - ▶ For PR, "Observation" was added.

FOLL-B 1 of 4

- First-line Therapy
 - ▶ Lenalidomide + obinutuzumab was added as a category 2B, Other recommended option.
- First-line Therapy for Elderly or Infirm
 - ▶ Preference stratification was applied to the recommendations.
- First-line Consolidation or Extended Dosing (optional)
 - ▶ Preference stratification was applied to the recommendations.
 - ▶ Rituximab maintenance dosing was changed from "one dose every 8 weeks" to "one dose every 8-12 weeks."
- Footnote g, 2nd sentence was revised from, "Karyotype ± FISH for known" to "Cytogenetics/FISH assessment for MDS markers." (Also for [FOLL-B 2 of 4](#))

FOLL-B 2 of 4

- Second-line and Subsequent Therapy
 - ▶ Within the preference stratification, the regimens are now listed in alphabetical order.
 - ▶ Preferred,
 - ◊ "Rituximab" was moved from Preferred regimens to Other recommended regimens.
 - ◊ "Lenalidomide ± rituximab" was changed to "Lenalidomide + rituximab."
 - ▶ Other recommended regimens, the following options were added as category 2A recommendations
 - ◊ Lenalidomide (if not a candidate for anti-CD20 monoclonal antibody therapy)
 - ◊ Lenalidomide + obinutuzumab
 - ◊ Obinutuzumab
- Second-line and Subsequent Therapy for Elderly or Infirm
 - ▶ Preference stratification was applied to the recommendations.
- Second-line Consolidation or Extended Dosing (optional)
 - ▶ Preference stratification was applied to the recommendations.

Gastric MALT Lymphoma

MALT-1

- ▶ Workup, Essential
 - ◊ 1st bullet was revised, "Physical exam ~~with attention to nongastric sites (eyes, skin):~~."
 - ◊ 9th bullet was revised, "C/A/P CT with contrast of diagnostic quality ~~and/or wholebody PET/CT scan (especially if ISRT anticipated)~~" and "PET/CT scan (especially if ISRT anticipated)" was added to Useful in Selected Cases.

MALT-2

- Stage I₁ or I₂ or Stage II₁, H. pylori positive, t(11;18) positive disease
 - ▶ Initial Therapy
 - ◊ "Currently accepted antibiotic therapy for H. pylori" recommendation was revised by adding, "+ ISRT."
 - ◊ "Rituximab (if ISRT is contraindicated)" was added.
 - ◊ The algorithm is then directed to MALT-5.

MALT-4

- Footnotes
 - ▶ Footnote p was revised by adding, "Reassessment to rule out H. Pylori by institutional standards." (Also for [MALT-5](#) and [MALT-6](#))
 - ▶ Footnote was removed, "If patient originally had clinical stage I2 or stage IIE, early RT should be considered if there is no response to antibiotics."

Nongastric MALT Lymphoma

- Title was revised, "Nongastric MALT Lymphoma (*Noncutaneous*)."

NGMLT-1

- Workup, Essential,
 - ▶ 1st bullet was revised, "Physical exam *with attention to nongastric sites*"
 - ▶ 8th bullet was revised, "C/A/P CT and other suspected sites with contrast of diagnostic quality ~~and/or~~" and "Whole-body PET/CT scan..." was moved to its own bullet.
 - ▶ 9th bullet was revised by adding, "Whole-body...essential if RT for stage I, II disease or systemic therapy is planned."
 - ▶ Useful in Selected Cases
 - ◊ 5th bullet was added, "MRI of head/neck, cranial, ocular adnexa."
 - ◊ 6th bullet was added, "Autoimmune disease testing, particularly Sjogren's."
- Footnotes
 - ▶ Footnote a was revised, "...head and neck, lung, *dural*, ocular adnexa,..." and "...is not required for management *in the United States*."

[Continued](#)



Updates in Version 1.2020 of the NCCN Guidelines for B-Cell Lymphomas from Version 7.2019 include:

Nodal Marginal Zone Lymphoma

NODE-1

- Workup
 - ▶ Essential
 - ◊ 7th bullet was revised, "Chest/abdominal/pelvic (C/A/P) CT with contrast of diagnostic quality **and/or**" and "Whole-body PET/CT" was moved to its own bullet.
 - ◊ 8th bullet was revised, "Whole-body PET/CT scan essential if RT for stage I, II disease *or systemic therapy is planned*"

NODE-2

- Stage I and contiguous stage II size qualifiers were removed, "<7 cm" and "≥7 cm"
- Stage I or Contiguous Stage II, Initial Therapy,
 - ▶ "ISRT" was listed as "preferred."
 - ▶ "Anti-CD20 monoclonal antibody ± chemotherapy (See MZL-A) + ISRT (category 2B)" was added.
- Non-contiguous Stage II, Initial Therapy,
 - ▶ "± ISRT for local palliation" was added to "Anti-CD20 monoclonal antibody ± chemotherapy (See MZL-A) + ISRT"
 - ▶ "Anti-CD20 monoclonal antibody ± chemotherapy (See FOLL-B) + ISRT (category 2B)" was removed.
- Response to Therapy
 - ◊ PR or NR was revised, "Consider ISRT *if not previously given*"
- Footnotes
 - ▶ Footnote g was revised, "Observation may be appropriate in circumstances where potential toxicity of ISRT *or systemic therapy* outweighs potential clinical benefit *in consultation with a radiation oncologist.*"
 - ▶ Footnote k was revised by adding, "Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated." (Also for [NODE-3](#) and [NODE-4](#))

NODE-5

- Footnote q was added, "NGS may be useful for treatment selection."

NODE-6

- Histologic Transformation to DLBCL after multiple lines of prior therapy
 - ▶ "Anti-CD19 CAR T-cell Therapy (only after ≥2 prior chemoimmunotherapy regimens, if not previously given)" was added as a category 2A, preferred recommendation.
 - ▶ "Polatuzumab vedotin ± bendamustine ± rituximab" was added as a category 2A recommendation.
- Footnotes

- ▶ Footnote w was added, "See Guidance for Treatment of Patients with Chimeric Antigen Receptor (CAR) T-cell Therapy (BCEL-D)."
- ▶ Footnote x was added, "Patients should have received at least one anthracycline or anthracenedione-based regimen, unless contraindicated."

Splenic Marginal Zone Lymphoma

SPLN-1

- Workup
 - ▶ 7th bullet was revised, "Chest/abdominal/pelvic (C/A/P) CT with contrast of diagnostic quality **and/or**" and "Whole-body PET/CT" and was added to "Useful in Selected Cases."

SPLN-2

- Splenomegaly, hepatitis C negative, management was revised, "Splenectomy (~~if not responsive to rituximab~~) (category 2B)."

Marginal Zone Lymphomas

MZL-A 1 of 3

- First-line Therapy,
 - ▶ Preferred, 4th sub-bullet was revised, "Rituximab (375 mg/m² weekly for 4 doses) (~~preferred~~ for SMZL)."
 - ▶ Other recommended regimens, 3rd sub-bullet was added, "Rituximab (375 mg/m² weekly for 4 doses) for extranodal (MALT) and nodal MZL."
- First-line Therapy for Elderly and Infirm
 - ▶ Preference stratification was applied to the recommendations.
- First-line Extended Therapy (optional)
 - ▶ Rituximab bullet was revised by adding, "for up to 2 years."
- Footnote d, sentence was revised from, "Karyotype ± FISH for known..." to "Cytogenetics/FISH assessment..." (Also for [MZL-A 2 of 3](#))

MZL-A 2 of 3

- Second-Line and Subsequent Therapy
 - ▶ Preference stratification was applied to the recommendations.
 - ▶ Preferred regimens, the following regimens were revised,
 - ◊ Bendamustine + rituximab (*not recommended if treated with prior bendamustine*) (*may be considered for both nodal MZL and extranodal [MALT] lymphoma*)
 - ◊ Bendamustine + obinutuzumab (*not recommended if treated with prior bendamustine*)
 - ◊ Lenalidomide ± + rituximab (*may be considered for both nodal MZL and extranodal [MALT] lymphoma*)



Updates in Version 1.2020 of the NCCN Guidelines for B-Cell Lymphomas from Version 7.2019 include:

- ▶ **Other recommended regimens**
 - ◊ 3rd sub-bullet revised, "Rituximab (*if longer duration of remission*)."
- **Second-line and Subsequent Therapy for Elderly and Infirm**
 - ▶ Preference stratification was applied to the recommendations.
 - ▶ Preferred therapies, the following were added as category 2A recommendations
 - ◊ Ibrutinib
 - ◊ Lenalidomide + rituximab
- **Second-line Consolidation or Extended Dosing (optional)**
 - ▶ Preference stratification was applied to the recommendations.

Mantle Cell Lymphoma

MANT-1

- **Additional Diagnostic Testing**
 - ▶ Essential, 2nd bullet moved from Useful Under Certain Circumstances, "TP53 sequencing (patients with typical MCL with an expected aggressive clinical course or particularly if upfront transplant anticipated)."
- **Workup**
 - ▶ "Bone marrow biopsy ± aspirate" moved from Essential to Useful Under Certain Circumstances.
 - ▶ 7th bullet was revised, "Chest/abdominal/pelvic (C/A/P) CT with contrast of diagnostic quality and/or" and "Whole-body PET/CT" was moved to its own bullet.
 - ▶ 8th bullet was revised, "Whole-body PET/CT scan essential if RT for stage I, II disease or systemic therapy is planned."

MANT-2

- Stage I, II phrase "localized presentation, extremely rare" has been moved to footnote h.
- Stage II has been divided into "contiguous" and "noncontiguous."
- The options for both stage I or contiguous stage II, nonbulky and noncontiguous stage II, nonbulky were clarified.
- Noncontiguous stage II option, "Observe in highly selected cases" has been directed to "see MANT-4."
- **Follow-up**
 - ▶ Clinical recommendation has been revised, "Follow-up H&P and labs every 3–6 mo for 5 y and then yearly or as clinically indicated."
 - ▶ Surveillance imaging recommendations added,
 - ◊ Up to 2 y post completion of treatment: C/A/P CT scan with contrast no more than every 6 mo
 - ◊ >2 y: No more than annually

MANT-3

• Footnotes

- ▶ Footnote n was added, "Patients who have achieved near CR, can proceed to HDT/ASCR. Patients who have achieved minimal PR with substantial disease should be treated as having stable, refractory disease. Patients who have achieved a very good PR, may be treated with additional therapy to achieve CR with the goal of proceeding to HDT/ASCR."
- ▶ Footnote o was added, "Patients who have achieved a very good PR or better, can be observed or consider rituximab maintenance. Patients who have achieved minimal PR with substantial disease should be treated as having stable, refractory disease."

MANT-A 1 of 4

• Induction Therapy

▶ Less aggressive therapy

- ◊ Option was moved from preferred to other recommended, "Modified rituximab-HyperCVAD in patients older than 65 y."
- ◊ Other recommended regimen, 2nd option was revised, "RBAC500 (rituximab, bendamustine, cytarabine)" and the category was changed from a category 2B to a category 2A recommendation."
- Maintenance after less aggressive therapy, 2nd sub-bullet was revised, "Untested after VR-CAP, RBAC500"

MANT-A 2 of 4

• Second-line therapy

- ▶ "Partial response with intention to proceed to transplant" options were added with "Preferred regimens (in alphabetical order)" including the following:
 - ◊ Bendamustine ± rituximab (if not previously given)
 - ◊ Bortezomib ± rituximab
 - ◊ Lenalidomide ± rituximab
 - ◊ RCHOP or VR-CAP (if not previously given)"
- ▶ Extended response duration to prior chemotherapy (> expected median PFS),
 - ◊ Preferred regimens,
 - BTK inhibitors were moved from Other recommended,
 - Acalabrutinib
 - Ibrutinib ± rituximab
 - Zanubrutinib
 - Lenalidomide ± rituximab was added.

[Continued](#)

UPDATES



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Updates in Version 1.2020 of the NCCN Guidelines for B-Cell Lymphomas from Version 7.2019 include:

• Footnotes

- ▶ Footnote e was added, "Patients who have achieved near CR, can proceed to HDT/ASCR. Patients who have achieved minimal PR with substantial disease should be treated as having stable, refractory disease. Patients who have achieved a very good PR, may be treated with additional therapy to achieve CR with the goal of proceeding to HDT/ASCR."
- ▶ Footnote f was added, "In patients intended to receive HDT/ASCR, bendamustine should be used with caution as there are conflicting data regarding ability to collect peripheral progenitor cell collection."

Diffuse Large B-Cell Lymphoma

BCEL-1

• Additional Diagnostic Testing

- ▶ Essential, 2nd bullet was added, "Karyotype or FISH for *MYC*."
- ▶ Useful, 2nd bullet was revised, "Karyotype or FISH for *BCL2*, *BCL6* rearrangements if *MYC* positive."

• Footnotes

- ▶ Footnote was removed, "Cases with double expression of *MYC* and either *BCL2* or *BCL6* by IHC having a GCB-like immunophenotype should undergo FISH testing for *MYC*, *BCL2*, and *BCL6* rearrangement."
- ▶ Footnote g was revised, "~~Controversy exists regarding management of FL grade 3. Some may treat FL grade 3a as follicular lymphoma and others may treat it as DLBCL. FL, grade 3b is commonly treated as diffuse large B-cell lymphoma (BCEL-1). The management of FL, grade 3a is controversial and treatment should be individualized.~~"

BCEL-2

• Workup,

- ▶ Essential, 9th bullet was revised, "Adequate bone marrow biopsy (>1.6 cm) ± aspirate; ~~bone marrow may not be needed if PET/CT scan negative unless finding of another lymphoma subtype is important for treatment decision bone marrow biopsy is not necessary if PET/CT scan demonstrates bone disease. Bone marrow biopsy with a negative PET/CT scan may reveal discordant lymphoma~~"
- ▶ Useful in Selected Cases, 6th bullet was revised, "Lumbar puncture for patients at risk for CNS involvement, see BCEL-A 2 of 2 consider for patients with 4–6 factors according to prognostic model HIV lymphoma, testicular lymphoma, high-grade B-cell lymphomas (HGBL) with translocations of *MYC* and *BCL2* and/or *BCL6*, or HGBL, NOS"

BCEL-3

- Stage I and II, nonbulky tumors <7.5 cm was separated into "All others" and "IPI=0"

- "RCHOP x 4 cycles + rituximab x 2 cycles" was added as a first-line therapy option for IPI=0.

BCEL-4

- Pre RT Evaluation, Stage I, II partial response category was separated into two categories and the options were clarified,
 - ▶ Partial response (PET positive [5-PS 4])
 - ▶ Partial response (PET positive [5-PS 5])
- Footnote w was added, "Chemosensitive partial response."

BCEL-6

- Footnote aa was added, "A 5-PS = 4 response can reflect post-treatment inflammation as well as active disease. If there is uncertainty regarding interpretation of the response, consider brief interval restaging to clarify post-treatment inflammation response."

BCEL-7

• Consolidation/Additional Therapy

- ▶ Option for partial response for patients intended to proceed to transplant was revised, "*Anti-CD19 CAR T-cell therapy Axicabtagene ciloleucel or tisagenlecleucel*." Axicabtagene ciloleucel or tisagenlecleucel were added to BCEL-B. Also for BCEL-8.

BCEL-A 2 of 2

- Additional indications for CNS prophylaxis independent of CNS risk score were added as follows:
 - ▶ HIV lymphoma
 - ▶ Testicular lymphoma
 - ▶ High-grade B-cell lymphomas (HGBLs) with translocations of *MYC* and *BCL2* and/or *BCL6* HGBL, NOS
 - ▶ Primary cutaneous DLBCL, Leg type
 - ▶ Stage IE DLBCL of the breast
- Suggested CNS prophylactic therapies (Optimal management is uncertain) were added as follows:
 - ▶ Intrathecal methotrexate and/or cytarabine (4–8 doses) during the course of treatment
 - ▶ Systemic methotrexate (3–3.5 g/m² for 2–4 cycles) during the course of treatment
- Footnote was removed, "There may be an increased risk of CNS events in selected patients with 4–6 risk factors according to the above prognostic model, HIV lymphoma, testicular lymphoma, high-grade B-cell lymphomas (HGBL) with translocations of *MYC* and *BCL2* and/or *BCL6*, HGBL, NOS or kidney or adrenal involvement. The optimal management of CNS events is uncertain, but CNS prophylaxis can be considered with 4–8 doses of intrathecal methotrexate and/or cytarabine, or systemic methotrexate (3–3.5 g/m²) during the course of treatment. Recent data suggest that stage IE

[Continued](#)
UPDATES



Updates in Version 1.2020 of the NCCN Guidelines for B-Cell Lymphomas from Version 7.2019 include:

DLBCL of the breast is a potential risk factor for CNS disease."

[BCEL-B 1 of 3](#)

• Primary Mediastinal Large B-Cell Lymphoma

- ▶ Bullet was removed, "Role of RT in first-line therapy is controversial. If PET/CT scan was negative at the end of treatment and initial disease was non-bulky, observation may be considered."
- ▶ 6th bullet, 3rd sub-bullet, "Nivolumab ± brentuximab vedotin" was added as a category 2B recommendation with corresponding footnote f.

[BCEL-B 2 of 3](#)

• Grey Zone Lymphoma

- ▶ Prognosis and Treatment, 5th bullet was revised, "If localized disease, then *consolidative* RT is preferred."

[BCEL-B 3 of 3](#)

- Footnote j was added, "Selection of patients requires adequate marrow cellularity >15% and <25% involvement of lymphoma in bone marrow, and platelets >100,000...Cytogenetics/FISH assessment for MDS markers. As of 2010, updates suggest a trend towards an increased risk of MDS with RIT."

[BCEL-C 1 of 4](#)

• Preference stratification was applied to the recommendations for

- ▶ First-line therapy
- ▶ First-line Therapies for Patients with Poor Left Ventricular Function
- ▶ First-line Therapies for Very Frail Patients and Patients >80 Years of Age with Comorbidities

[BCEL-C 2 of 4](#)

• Preference stratification was applied to the recommendations for

- ▶ Second-line and Subsequent Therapy (intention to proceed to transplant) and (non-candidates for transplant)
- "Bendamustine, rituximab and polatuzumab vedotin-piiq" was changed to "Polatuzumab vedotin ± bendamustine ± rituximab (after ≥2 prior therapies)."

High-Grade B-Cell Lymphomas with Translocations of MYC and BCL2 and/or BCL6 (Double-/Triple-Hit Lymphoma)

[HGBL-1](#)

• Treatment

- ▶ 5th bullet, was revised by removing, "consider CNS prophylaxis according to institutional standards." (Also for [HGBLNOS-1](#))
- ▶ 6th bullet was added, "If localized disease, then consolidative RT is preferred."

[Burkitt Lymphoma](#)

[BURK-1](#)

- Workup, Useful in Selected Cases, 1st bullet added, "Hepatitis C testing"
- Footnotes
 - ▶ Footnote c was revised by removing, "simple karyotype with MYC rearrangement as sole abnormality"
 - ▶ Footnote f was revised by adding, "Most common karyotype is MYC rearrangement as a sole abnormality."

[BURK-2](#)

- Footnote m was added, "If there is no clinical trial available, some NCCN member institutions use non-cross resistant second-line therapy regimens listed on BURK-A in selected cases." (Also for BURK-3)

[BURK-A 1 of 2](#)

- Induction therapy, preference stratification was applied to the recommendations.
- Induction Therapy <60 y
 - ▶ Low Risk, Dose-adjusted EPOCH was revised, "... (Data ~~are for~~ included patients with leptomeningeal CNS disease; patients with parenchymal CNS disease were excluded in the clinical trials of this regimen without CNS disease." Also for High Risk <60 y and Induction Therapy ≥60 y.
 - ▶ High Risk, 1st bullet was added, "High-risk patients presenting with symptomatic CNS disease should be started with the portion of the systemic therapy that contains CNS penetrating drugs."
- Induction Therapy ≥60 y,
 - ▶ Dose-adjusted EPOCH, sub-bullet added, "In high-risk patients presenting with symptomatic CNS disease, the management of the CNS disease should be addressed with the initial regimen."
- Footnote was removed, "Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan." (Also for BURK-A 2 of 2, AIDS-3, AIDS-4)

[BURK-A 2 of 2](#)

• Second-line Therapy

- ▶ Preference stratification was applied to the recommendations.
- ▶ 1st bullet was revised, "Patients with disease relapse >6-18 mo after appropriate first-line therapy *should be treated with alternate regimens.*"
- ▶ Other recommended regimens, 1st sub-bullet was revised, "... (Data ~~are for~~ included patients with leptomeningeal CNS disease; patients with parenchymal CNS disease were excluded in the clinical trials of this regimen without CNS disease."

[Continued](#)
UPDATES



Updates in Version 1.2020 of the NCCN Guidelines for B-Cell Lymphomas from Version 7.2019 include:

AIDS-Related B-cell Lymphomas

AIDS-2

• Workup

- ▶ Bullet "bone marrow biopsy ± aspirate" was moved from Essential to Useful in Selected Cases and revised, "*Adequate bone marrow biopsy (>1.6 cm) ± aspirate; bone marrow biopsy is not necessary if PET/CT scan demonstrates bone disease. Bone marrow biopsy with a negative PET/CT scan may reveal discordant lymphoma.*"

AIDS-3

• Burkitt lymphoma

- ▶ "CODOX-M/IVAC ± rituximab" was changed to "CODOX-M/IVAC + rituximab."
- ▶ "EPOCH-R" was clarified as dose-adjusted.

• Primary CNS Lymphoma

- ▶ 4th bullet was revised, "*Consider RT alone as initial management for palliation of patients who are not candidates for systemic therapy at presentation, then reassess eligibility for systemic therapy after RT.*"

• Footnotes

- ▶ Footnote h was added, "High-risk patients presenting with symptomatic CNS disease should be started with the portion of the systemic therapy that contains CNS penetrating drugs."
- ▶ Footnote i was revised by adding, "Roschewski M, Dunleavy K, Abramson JS, et al. Risk-adapted therapy in adults with Burkitt lymphoma: Results of NCI 9177, a multicenter prospective phase II study of DA-EPOCH-R [abstract]. Blood 2017;130 (Suppl 1):Abstract 188."
- ▶ Footnote j was added, "Gupta N, et al. Long-term survival in AIDS-related primary central nervous system lymphoma. Neuro Oncol 2017;19:99-108."

AIDS-4

• DBLCL; HHV8-positive DLBCL, NOS; and PEL

- ▶ For relapse, therapy option was added, "Bortezomib-ICE ± rituximab (for DLBCL)" as a category 2B recommendation.

• Plasmablastic lymphoma

- ▶ "For relapse, see BCEL-C second-line therapy (without rituximab)" was added.

• Footnotes

- ▶ Footnote was removed, "Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan."
- ▶ Footnote l was revised by removing, "CNS prophylaxis indicated if double-

hit lymphoma or lymphomatous involvement of bone marrow, testes, sinuses, or epidural regions. The specific regimen is at the discretion of the primary oncologist but should include 4–6 doses of therapy (IT cytarabine or IT methotrexate). If a patient presents with positive CSF or leptomeningeal involvement, intrathecal therapy should be administered twice weekly until CSF is negative, then weekly x 1 month, then monthly x 6 months. Although the specific regimen may vary, a suggested regimen is: cytarabine 50 mg + methotrexate 12 mg + hydrocortisone 50 mg."

AIDS-B

• EPOCH ± rituximab dosing recommendations for cycle 1

- ▶ 5th bullet, 3rd sub-bullet was revised by removing, "For cycle 1, consider a starting dose of 187.5 mg/m² and consider initial therapy with EPOCH alone with the addition of rituximab in future cycles if the EPOCH is tolerated and CD4 count has improved with concurrent antiretroviral therapy."

[Continued](#)



Updates in Version 1.2020 of the NCCN Guidelines for B-Cell Lymphomas from Version 7.2019 include:

Post-Transplant Lymphoproliferative Disorders

PTLD-1

- **Workup**
 - ▶ **Essential**, 7th bullet was added, "EBV PCR for cell-free plasma EBV DNA marker."
 - ▶ **Useful**, bullet was removed, "EBV PCR for cell-free plasma EBV DNA marker"
- **PTLD subtype** was revised, "*Non-destructive early lesions*" (Also for [PTLD-2](#))
- **Footnotes**
 - ▶ Footnote was removed, "BCL6 positivity has been associated with a poor response to reduction in immunosuppressive therapy."
 - ▶ Footnote b was added, "If EBV-negative, should not be used as response marker."
 - ▶ Footnote d was revised, "*Early Non-destructive lesions...*" (Also for [PTLD-2](#))

PTLD-A

- **Monomorphic PTLD (T-cell type)**, 1st bullet was added, "Brentuximab vedotin + CHP (cyclophosphamide, doxorubicin, prednisone) for CD30+ cases" was added as a category 2A recommendation.
- **Primary CNS PTLD (B-cell type)**, "High-dose methotrexate + rituximab" was added as a category 2A recommendation.
- Footnote e was added, "Patients with PTLD often have renal insufficiency. High-dose methotrexate should be used with caution. Alternate regimens (cytarabine-based) should be considered."

Castleman's Disease

CD-1

- **Workup**
 - ▶ **Useful**, 3rd bullet was added, "Reticulin fibrosis of bone marrow (particularly in patients with TAFRO syndrome)."
- Footnote e was revised by adding, "Rule out other diseases that can mimic idiopathic MCD (see Exclusion criteria on CD-B 1 of 2)."

CD-3

- **MCD (Criteria for active disease present but no organ failure)**
 - ▶ **HIV-1(-) HHV8 (-)**
 - ◊ Preference stratification was applied to the primary treatment recommendations.
 - ◊ "Siltuximab (for plasmacytic/mixed histology)" was clarified as preferred.
 - ◊ Other recommended regimen, "Thalidomide, cyclophosphamide,

prednisone (for hyaline vascular histology)" was added as a category 2A other recommended option.

- **Footnotes**
 - ▶ Footnote n was added, "See Diagnostic Criteria for Idiopathic MCD (CD-B)."
 - ▶ Footnote p was added, "Patients with hyaline vascular histology do not benefit from siltuximab."

CD-A

- A new section, "Subtypes of Idiopathic MCD" was added.

CD-B

- Table for "Consensus Diagnostic Criteria for Idiopathic MCD" was added.

Supportive Care for B-Cell Lymphomas

NHODG-B 3 of 4

- "Prophylaxis for PJP and VZV" section was added with a bullet, "See NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections"
- "Hypogammaglobulinemia" section was added with a bullet, "Patients receiving anti-CD20 MAB and anti-CD19 directed CAR T-cell therapy may experience hypogammaglobulinemia. Patients with recurrent infections may benefit from IVIG replacement."

Principles of Radiation Therapy

NHODG-D

- The principles pages were extensively revised.

**DIAGNOSIS**

- **Excisional or incisional biopsy.** An FNA biopsy alone is not generally suitable for the initial diagnosis of lymphoma. A core needle biopsy is not optimal but can be used under certain circumstances. In certain circumstances, when a lymph node is not easily accessible for excisional or incisional biopsy, a combination of core biopsy (multiple biopsies preferred) and FNA biopsies in conjunction with appropriate ancillary techniques for the differential diagnosis (immunohistochemistry [IHC], flow cytometry, PCR for *IGHV* and *TCR* gene rearrangements, karyotype, and FISH for major translocations^a) may be sufficient for diagnosis.
- **Histologic grading** cannot be performed on an FNA.
- **Hematopathology review** of all slides with at least one paraffin block representative of the tumor. Rebiopsy if consult material is nondiagnostic.



- | | ADDITIONAL
DIAGNOSTIC TESTING |
|--------------------------------------------------------------------------------------------------------------------------|------------------------------------------|
| • Follicular lymphoma | → See FOLL-1 |
| • Gastric MALT lymphoma | → See MALT-1 |
| • Nongastric MALT lymphoma | → See NGMLT-1 |
| • Nodal marginal zone lymphoma | → See NODE-1 |
| • Splenic marginal zone lymphoma | → See SPLN-1 |
| • Mantle cell lymphoma | → See MANT-1 |
| • Diffuse large B-cell lymphoma | → See BCEL-1 |
| • High-grade B-cell lymphomas with translocations of <i>MYC</i> and <i>BCL2</i> and/or <i>BCL6</i> (double-hit lymphoma) | → See HGBL-1 |
| • Burkitt lymphoma | → See BURK-1 |
| • AIDS-related B-cell lymphomas | → See AIDS-1 |
| • Lymphoblastic lymphoma | → See BLAST-1 |
| • Post-transplant lymphoproliferative disorders | → See PTLD-1 |
| • Castleman's disease | → See CD-1 |

^a If a high suspicion of a clonal process remains and other techniques have not resulted in a clear identification of a clonal process, then next-generation sequencing (NGS) can be used.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Follicular Lymphoma (grade 1–2)


ADDITIONAL DIAGNOSTIC TESTING^a

ESSENTIAL:

- Adequate immunophenotyping to establish diagnosis^{b,c}
 - ▶ IHC panel: CD20, CD3, CD5, CD10, BCL2,^d BCL6, CD21, or CD23, with or without
 - ▶ Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: kappa/lambda, CD19, CD20, CD5, CD23, CD10

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- Molecular analysis to detect: antigen receptor gene rearrangements; *BCL2* rearrangements^d
- Karyotype or FISH^{e,f}: t(14;18); *BCL6*, 1p36, *IRF4/MUM1* rearrangements^d
- IHC panel: Ki-67^g; *IRF4/MUM1* for FL grade 3, cyclin D1
- NGS panel including *TNFRSF14* and *STAT6* mutation



Germinal center or follicular center cell phenotype type is not equivalent to follicular lymphoma and occurs in Burkitt lymphoma and some DLBCL.

^a Follicular lymphoma (FL), grade 1–2. FL, grade 3b is commonly treated as diffuse large B-cell lymphoma ([BCEL-1](#)). The management of FL, grade 3a is controversial and treatment should be individualized. Any area of DLBCL in an FL of any grade should be diagnosed and treated as a DLBCL.

^b Typical immunophenotype: CD10+, BCL2+, CD23+/-, CD43-, CD5-, CD20+, BCL6+. Rare cases of follicular lymphoma may be CD10- or BCL2-.

^c [See Use of Immunophenotyping/Genetic Testing in Differential Diagnosis of Mature B-Cell and NK/T-Cell Neoplasms \(NHODG-A\)](#).

^d In young patients with localized disease that lacks BCL2 expression or t(14;18), consider entity of PTFL. Analysis of *BCL6* rearrangement may be useful for evaluating the diagnosis of PTFL.

^e FL with 1p36 deletions have a predominant diffuse pattern in inguinal nodes, large localized mass, CD23+, typically grade 1–2 and have a good prognosis.

^f Lymphomas with *IRF4* translocations are usually DLBCL but occasionally are purely FL grade 3b and often DLBCL with FL grade 3b. Patients typically present with Waldeyer's ring involvement and are often children/young adults. The tumor is locally aggressive but responds well to chemotherapy +/- RT. These lymphomas do not have a *BCL2* rearrangement and should not be treated as low-grade FL.

^g There are reports showing that Ki-67 proliferation fraction of >30% may be associated with a more aggressive clinical behavior, but there is no evidence that this should guide treatment decisions.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Follicular Lymphoma (grade 1–2)


WORKUP

ESSENTIAL:

- Physical exam: attention to node-bearing areas, including Waldeyer's ring, and to size of liver and spleen
- Performance status
- B symptoms
- CBC with differential
- LDH
- Comprehensive metabolic panel
- Hepatitis B testing^h
- Chest/abdominal/pelvic (C/A/P) CT with contrast of diagnostic quality
- Whole-body PET/CT scan essential if RT for stage I, II disease or systemic therapy is planned
- Bone marrow biopsy + aspirate to document clinical stage I–II diseaseⁱ
- Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)

USEFUL IN SELECTED CASES:

- Echocardiogram or MUGA scan if anthracycline or anthracenedione-based regimen is indicated
- Neck CT with contrast
- Beta-2-microglobulin (necessary for calculation of FLIPI-2)
- Uric acid
- SPEP and/or quantitative immunoglobulin levels
- Hepatitis C testing
- Discussion of fertility issues and sperm banking



^h Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen ([See NHODG-B](#)). If positive, check viral load and consider consult with gastroenterologist.

ⁱ If ibritumomab tiuxetan is considered, bilateral cores are recommended and the pathologist should provide the percent of overall cellular elements and the percent of cellular elements involved in the marrow. If observation is initial therapy, bone marrow biopsy may be deferred. Selection of patients requires adequate marrow cellularity >15% and <25% involvement of lymphoma in bone marrow, and platelets >100,000. In patients with prior autologous stem cell rescue, referral to a tertiary care center is highly recommended for ibritumomab tiuxetan.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Follicular Lymphoma (grade 1–2)



^j See Principles of Radiation Therapy (NHODG-D).

^k Observation may be appropriate in circumstances where potential toxicity of involved-site RT (ISRT) or systemic therapy outweighs potential clinical benefit in consultation with a radiation oncologist.

^l Anti-CD20 monoclonal antibodies include rituximab or obinutuzumab. Obinutuzumab is not indicated as single-agent therapy.

^m Initiation of chemotherapy or more extended RT can improve failure-free survival (FFS), but has not been shown to improve overall survival. These are options for therapy.

ⁿ See Lugano Response Criteria for Non-Hodgkin Lymphoma (NHODG-C). PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS).

^o Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated. Imaging should be performed whenever there are clinical indications. For surveillance imaging, see Discussion for consensus imaging recommendations.

^p Consider possibility of histologic transformation in patients with progressive disease, especially if LDH levels are rising, single site is growing disproportionately, extranodal disease develops, or there are new B symptoms. If clinical suspicion of transformation, FDG-PET may help identify areas suspicious for transformation. FDG-PET scan demonstrating marked heterogeneity or sites of intense FDG avidity may indicate transformation, and biopsy should be directed biopsy at the most FDG-avid area. Functional imaging does not replace biopsy to diagnose transformation. If transformation is histologically confirmed, treat with anthracycline-based therapy. See Management of Transformation (FOLL-6).


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020


Follicular Lymphoma (grade 1–2)

^j See [Principles of Radiation Therapy \(NHODG-D\)](#).ⁿ See [Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#). PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS).^o Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated. Imaging should be performed whenever there are clinical indications. For surveillance imaging, see [Discussion](#) for consensus imaging recommendations.^p Consider possibility of histologic transformation in patients with progressive disease, especially if LDH levels are rising, single site is growing disproportionately, extranodal disease develops, or there are new B symptoms. If clinical suspicion of transformation, FDG-PET may help identify areas suspicious for transformation. FDG-PET scan demonstrating marked heterogeneity or sites of intense FDG avidity may indicate transformation, and biopsy should be directed biopsy at the most FDG-avid area. Functional imaging does not replace biopsy to diagnose transformation. If transformation is histologically confirmed, treat with anthracycline-based therapy. See [Management of Transformation \(FOLL-6\)](#).^q See [GELF criteria \(FOLL-A\)](#).^r When disease (eg, bone) is only imaged on PET/CT scan, it is appropriate to be used in follow-up.**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Follicular Lymphoma (grade 1–2)



^j See Principles of Radiation Therapy (NHODG-D).

ⁿ See Lugano Response Criteria for Non-Hodgkin Lymphoma (NHODG-C). PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS).

^o Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated. Imaging should be performed whenever there are clinical indications. For surveillance imaging, see Discussion for consensus imaging recommendations.

^p Consider possibility of histologic transformation in patients with progressive disease, especially if LDH levels are rising, single site is growing disproportionately, extranodal disease develops, or there are new B symptoms. If clinical suspicion of transformation, FDG-PET may help identify areas suspicious for transformation. FDG-PET scan demonstrating marked heterogeneity or sites of intense FDG avidity may indicate transformation, and biopsy should be directed biopsy at the most FDG-avid area. Functional imaging does not replace biopsy to diagnose transformation. If transformation is histologically confirmed, treat with anthracycline-based therapy. See Management of Transformation (FOLL-6).

^q See GELF criteria (FOLL-A).

^r When disease (eg, bone) is only imaged on PET/CT scan, it is appropriate to be used in follow-up.


^s A PET-positive PR is associated with a shortened PFS (See Discussion); however, additional treatment at this juncture has not been shown to change outcome.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



HISTOLOGIC TRANSFORMATION TO DIFFUSE LARGE B-CELL LYMPHOMA



^j [See Principles of Radiation Therapy \(NHODG-D\)](#).

ⁿ [See Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#). PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS).

^t Perform FISH for *BCL2* rearrangement [t(14;18)], and *MYC* rearrangements [t(8;14) or variants, t(8;22), t(2;8)].

^u ISRT alone or one course of single-agent therapy including rituximab.

^v NGS may be useful for treatment selection.

^w This includes ≥2 of chemoimmunotherapy regimens for indolent or transformed disease. For example, prior treatment with BR and RCHOP.

^x Consider ISRT for localized presentations, bulky disease, and/or limited osseous disease.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Follicular Lymphoma (grade 1–2)

HISTOLOGIC TRANSFORMATION TO DIFFUSE LARGE B-CELL LYMPHOMA

RESPONSE TO THERAPY^o

CR^{n,y}

Observation
or
Clinical trial
or
High-dose therapy with autologous stem cell rescue (HDT/ASCR) or allogeneic hematopoietic cell transplant (HCT) in selected cases ± ISRT if not previously given^{j,x}

Anti-CD19 CAR T-cell therapy (only after ≥2 prior chemoimmunotherapy regimens)^{bb,cc} (preferred)
or
HDT/ASCR^z or allogeneic HCT in selected cases^z ± ISRT if not previously given^{j,x}

PR^{n,z,aa}

Clinical trial
or
Ibritumomab tiuxetan
or
ISRT^j for localized residual and/or residual FDG-avid disease not previously irradiated
or
Observation

NR or progressive diseaseⁿ

Clinical trial
or
Anti-CD19 CAR T-cell therapy (only after ≥2 prior chemoimmunotherapy regimens)^{bb,cc}
or
Polatuzumab vedotin ± bendamustine ± rituximab
or
Ibritumomab tiuxetan or Second-line therapy ([See BCEL-C](#))
or
Best supportive care

Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

→ Relapsed or progressive diseaseⁿ → Biopsy^{aa}

[See Histologic Transformation to DLBCL After Multiple Prior Therapies \(FOLL-8\)](#)

^j [See Principles of Radiation Therapy \(NHODG-D\)](#).

ⁿ [See Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#). PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS).

^o Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated. Imaging should be performed whenever there are clinical indications. For surveillance imaging, see [Discussion](#) for consensus imaging recommendations.

^x Consider ISRT for localized presentations, bulky disease, and/or limited osseous disease.

^y If transformation is co-existing with extensive FL, consider maintenance ([see FOLL-5](#), Optional Extended Therapy).

^z If proceeding to transplant, consider additional systemic therapy ± ISRT to induce CR prior to transplant. Axicabtagene ciloleucel or tisagenlecleucel are not appropriate treatment options for patients with a CR.

^{aa} Repeat biopsy should be strongly considered if PET-positive prior to additional therapy because PET positivity may represent post-treatment inflammation. Patients with a durable response for transformed disease may recur with the original indolent lymphoma. In that case, the management should be as per [FOLL-5](#). If biopsy negative, follow CR pathway.

^{bb} [See Guidance for Treatment of Patients with Chimeric Antigen Receptor \(CAR\) T-Cell Therapy \(BCEL-D\)](#).

^{cc} Patients should have received at least one anthracycline or anthracenedione-based regimen, unless contraindicated.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Follicular Lymphoma (grade 1–2)

HISTOLOGIC TRANSFORMATION TO DIFFUSE LARGE B-CELL LYMPHOMA^o

^j See [Principles of Radiation Therapy \(NHODG-D\)](#).ⁿ See [Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#). PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS).^o Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated. Imaging should be performed whenever there are clinical indications. For surveillance imaging, see [Discussion](#) for consensus imaging recommendations.^w This includes ≥2 of chemoimmunotherapy regimens for indolent or transformed disease. For example, prior treatment with BR and RCHOP.^x Consider ISRT for localized presentations, bulky disease, and/or limited osseous disease.^z If proceeding to transplant, consider additional systemic therapy ± ISRT to induce CR prior to transplant. Axicabtagene ciloleucel or tisagenlecleucel are not appropriate treatment options for patients with a CR.^{aa} Repeat biopsy should be strongly considered if PET-positive prior to additional therapy because PET positivity may represent post-treatment inflammation. Patients with a durable response for transformed disease may recur with the original indolent lymphoma. In that case, the management should be as per [FOLL-5](#). If biopsy negative, follow CR pathway.^{bb} See [Guidance for Treatment of Patients with Chimeric Antigen Receptor \(CAR\) T-Cell Therapy \(BCEL-D\)](#).^{cc} Patients should have received at least one anthracycline or anthracenedione-based regimen, unless contraindicated.^{dd} Data on transplant after treatment with axicabtagene ciloleucel or tisagenlecleucel are not available. HDT/ASCR is not recommended after axicabtagene ciloleucel. Allogeneic HCT could be considered but remains investigational.**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Follicular Lymphoma (grade 1–2)

PEDIATRIC-TYPE FOLLICULAR LYMPHOMA IN ADULTS

PATHOLOGIC AND CLINICAL PRESENTATION^{d,ee}

- Pathologic
 - Morphology: expansile follicles, effacement of architecture, absence of diffuse area
 - Expresses: BCL6, CD10, ± IRF4/MUM1 (~20%)
 - Proliferation index (Ki-67/MIB-1) >30%
 - No rearrangement of *BCL2*, *BCL6*, *IRF4/MUM1*
- Clinical
 - Localized disease (stage I, II)
 - Head and neck (cervical, submandibular, submental, postauricular, or periparotid lymph nodes) or less common inguinal lymph nodes
 - Male sex predominant
 - Younger age than typical FL (though can occur in adults older than age 60)

STAGING WORKUP

- PET/CT scan
- Bone marrow biopsy


Stage
I, II

TREATMENT

Excision (preferred) or
ISRT^j
or
RCHOP for patients with
extensive local disease who
are not candidates for
excision or ISRT

Consider prophylaxis for tumor
lysis syndrome ([See NHODG-B](#))

See monoclonal antibody and
viral reactivation ([NHODG-B](#))



^d In young patients with localized disease that lacks BCL2 expression or t(14;18), consider entity of PTFL. Analysis of *BCL6* rearrangement may be useful for evaluating the diagnosis of PTFL.

^j [See Principles of Radiation Therapy \(NHODG-D\)](#).

ⁿ [See Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#). PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS).

^{ee} Localized disease (stage I, II) is more common than advanced-stage disease (stage III, IV). If the patient has disease >stage II, it is by definition not PTFL.

^{ff} If patients have an excellent prognosis, no surveillance imaging is necessary. There are no data to support maintenance therapy.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Follicular Lymphoma (grade 1–2)

GELF CRITERIA^{a,b}

- Involvement of ≥ 3 nodal sites, each with a diameter of ≥ 3 cm
- Any nodal or extranodal tumor mass with a diameter of ≥ 7 cm
- B symptoms
- Splenomegaly
- Pleural effusions or peritoneal ascites
- Cytopenias (leukocytes $<1.0 \times 10^9/L$ and/or platelets $<100 \times 10^9/L$)
- Leukemia ($>5.0 \times 10^9/L$ malignant cells)

FLIPI - 1 CRITERIA^{a,c,d}


Age	≥ 60 y
Ann Arbor Stage	III-IV
Hemoglobin level	<12 g/dL
Serum LDH level	$>ULN$ (upper limit of normal)
Number of nodal sites ^d	≥ 5

Risk group according to FLIPI chart

	Number of factors
Low	0-1
Intermediate	2
High	≥ 3

^a This provides useful prognostic information that may be used to guide therapeutic decisions.

^b Solal-Celigny P, Lepage E, Brousse N, et al. Doxorubicin-containing regimen with or without interferon alfa 2b for advanced follicular lymphomas: final analysis of survival and toxicity in the Groupe d'Etude des Lymphomes Folliculaire 86 trial. J Clin Oncol 1998;16:2332-2338.



Mannequin used for counting the number of involved areas.^e

© 2007 Dana-Farber Cancer Institute, Inc.

All rights reserved. Permission is hereby granted for copying this image by photocopy or similar process for use in the practice of medicine or for research purposes. No other use is permitted which will infringe the copyright without the express written consent of Dana-Farber Cancer Institute, Inc.

^c This research was originally published in Blood. Solal-Celigny P, Roy P, Colombat P, et al. Follicular lymphoma international prognostic index. Blood 2004;104:1258-1265. (c) the American Society of Hematology.

^d FLIPI-2 (Federico M, Bellei M, Marcheselli L, et al. J Clin Oncol 2009;27:4555-4562) predicts for outcomes after active therapy; [see Discussion](#).

^e The map is used to determine the number of nodal sites in FLIPI-1 criteria and is different than the conventional Ann Arbor site map.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Follicular Lymphoma (grade 1–2)

SUGGESTED TREATMENT REGIMENS^{a,b,c}

An FDA-approved biosimilar is an appropriate substitute for rituximab.

First-line Therapy

- Preferred regimens (in alphabetical order)
 - Bendamustine^d + obinutuzumab^e or rituximab
 - CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) + obinutuzumab^e or rituximab
 - CVP (cyclophosphamide, vincristine, prednisone) + obinutuzumab^e or rituximab
 - Lenalidomide + rituximab
- Other recommended regimens
 - Lenalidomide + obinutuzumab (category 2B)
 - Rituximab (375 mg/m² weekly for 4 doses) (consider for low tumor burden)^f

First-line Therapy for Elderly or Infirm (if none of the above are expected to be tolerable in the opinion of treating physician)

- Preferred regimen
 - Rituximab (375 mg/m² weekly for 4 doses)
- Other recommended regimens
 - Chlorambucil ± rituximab
 - Cyclophosphamide ± rituximab
 - Ibritumomab tiuxetan^g (category 2B)

First-line Consolidation or Extended Dosing (optional)

- Preferred regimens following chemoimmunotherapy
 - Rituximab maintenance 375 mg/m² one dose every 8–12 weeks for 12 doses for patients initially presenting with high tumor burden (category 1)^h
 - Obinutuzumab maintenance (1000 mg every 8 weeks for 12 doses)
- Other recommended regimens
 - If initially treated with single-agent rituximab, consolidation with rituximab 375 mg/m² one dose every 8 weeks for 4 doses
 - Ibritumomab tiuxetan^{g,i} (category 2B)

[See Second-line and Subsequent Therapy on FOLL-B 2 of 4](#)

Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

^a See references for regimens on FOLL-B 3 of 4 and [FOLL-B 4 of 4](#).

^b The choice of initial therapy requires consideration of many factors, including age, comorbidities, and future treatment possibilities (eg, HDT with ASCR). Therefore, treatment selection is highly individualized.

^c Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^d In the GALLIUM study, there was an increased risk of mortality from opportunistic infections and secondary malignancies in patients receiving bendamustine. Increased risk of mortality occurred over the entire treatment program and extending beyond maintenance. Prophylaxis for PJP and VZV should be administered; [see NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections](#).

^e The clinical trial evaluating this regimen included obinutuzumab maintenance. The use without maintenance was an extrapolation of the data.

^f Rituximab may be appropriate in patients initially observed and with progression of low tumor burden disease not meeting GELF criteria ([FOLL-A](#)). Immediate initial therapy with rituximab in patients not meeting GELF criteria has not improved OS (Ardeshna K, et al. Lancet Oncol 2014;15:424-435).

^g Selection of patients requires adequate marrow cellularity >15% and <25% involvement of lymphoma in bone marrow, and platelets >100,000. In patients with prior autologous stem cell rescue, referral to a tertiary care center is highly recommended for ibritumomab tiuxetan. If ibritumomab tiuxetan is considered, bilateral cores are recommended and the pathologist should provide the percent of overall cellular elements and the percent of cellular elements involved in the marrow. As of 2010, updates suggest a trend towards an increased risk of MDS with RIT. Cytogenetics/FISH assessment for MDS markers is recommended for patients receiving RIT.

^h This is based on the PRIMA study for patients with high tumor burden treatment with RCVP and RCHOP. There are no data following other regimens.

ⁱ The full impact of an induction regimen containing rituximab on RIT consolidation is unknown.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Follicular Lymphoma (grade 1–2)

SUGGESTED TREATMENT REGIMENS^{a,b,c}

An FDA-approved biosimilar is an appropriate substitute for rituximab.

Second-line and Subsequent Therapy

- Preferred regimens^j (alphabetical order)
 - ▶ Bendamustine^k + obinutuzumab^l or rituximab
 - ▶ CHOP + obinutuzumab^l or rituximab
 - ▶ CVP + obinutuzumab^l or rituximab
 - ▶ Lenalidomide + rituximab
- Other recommended regimens (alphabetical order)
 - ▶ Ibritumomab tiuxetan^g
 - ▶ Lenalidomide (if not a candidate for anti-CD20 monoclonal antibody therapy)
 - ▶ Lenalidomide + obinutuzumab
 - ▶ Obinutuzumab
 - ▶ PI3K inhibitors (relapsed/refractory after 2 prior therapies)
 - ◊ Copanlisib^m
 - ◊ Duvelisib^m
 - ◊ Idelalisib^m
 - ▶ Rituximab
 - ▶ Tazemetostat
 - ◊ *EZH2* mutation positive relapsed/refractory disease after 2 prior therapies
 - ◊ *EZH2* wild type relapsed/refractory disease in patients who have no satisfactory alternative treatment options
 - ▶ [See Second-line Therapy for DLBCL \(BCEL-C 2 of 4\)](#) without regard to transplantability

^a See references for regimens on FOLL-B 3 of 4 and [FOLL-B 4 of 4](#).

^b The choice of initial therapy requires consideration of many factors, including age, comorbidities, and future treatment possibilities (eg, HDT with ASCR). Therefore, treatment selection is highly individualized.

^c Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^g Selection of patients requires adequate marrow cellularity >15% and <25% involvement of lymphoma in bone marrow, and platelets >100,000. In patients with prior autologous stem cell rescue, referral to a tertiary care center is highly recommended for ibritumomab tiuxetan. If ibritumomab tiuxetan is considered, bilateral cores are recommended and the pathologist should provide the percent of overall cellular elements and the percent of cellular elements involved in the marrow. As of 2010, updates suggest a trend towards an increased risk of MDS with RIT. Cytogenetics/FISH assessment for MDS markers is recommended for patients receiving RIT.

Second-line and Subsequent Therapy for Elderly or Infirm

(if none of the therapies are expected to be tolerable in the opinion of treating physician)

- Preferred regimen
 - ▶ Rituximab (375 mg/m² weekly for 4 doses)
- Other recommended regimens
 - ▶ Chlorambucil ± rituximab
 - ▶ Cyclophosphamide ± rituximab
 - ▶ Ibritumomab tiuxetan^g (category 2B)

Second-line Consolidation or Extended Dosing (optional)

- Preferred regimen
 - ▶ Rituximab maintenance 375 mg/m² one dose every 12 weeks for 2 years (category 1)
 - ▶ Obinutuzumab maintenance for rituximab-refractory disease (1 g every 8 weeks for total of 12 doses)
- Other recommended regimens
 - ▶ High-dose therapy with autologous stem cell rescue
 - ▶ Allogeneic hematopoietic cell transplant for highly selected patients

Histologic Transformation to DLBCL

- Anti-CD19 CAR T-cell therapy (only after ≥2 prior chemoimmunotherapy regimens)^{n,o}
 - ▶ Axicabtagene ciloleucel
 - ▶ Tisagenlecleucel

Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

^j Generally, a first-line regimen is not repeated.

^k Prophylaxis for PJP and VZV should be administered; [see NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections](#).

^l The clinical trial evaluating this regimen included obinutuzumab maintenance. The use without maintenance was an extrapolation of the data. Obinutuzumab is preferred in patients with rituximab refractory disease, which includes disease progressing on or within 6 months of prior rituximab therapy.

^m [See Special Considerations for the Use of Small-Molecule Inhibitors \(NHODG-E\)](#).

ⁿ [See Guidance for Treatment of Patients with Chimeric Antigen Receptor \(CAR\) T-Cell Therapy \(BCEL-D\)](#).

^o Patients should have received at least one anthracycline or anthracenedione-based regimen, unless contraindicated.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Follicular Lymphoma (grade 1–2)

SUGGESTED TREATMENT REGIMENS

REFERENCES

First-line Therapy

Bendamustine + rituximab

Rummel MJ, Niederle N, Maschmeyer G, et al. Bendamustine plus rituximab versus CHOP plus rituximab as first-line treatment for patients with indolent and mantle-cell lymphomas: an open-label, multicentre, randomised, phase 3 non-inferiority trial. *Lancet* 2013;381:1203-1210.

Flinn IW, van der Jagt R, Kahl BS, et al. Open-label, randomized, noninferiority study of bendamustine-rituximab or R-CHOP/R-CVP in first-line treatment of advanced indolent NHL or MCL: the BRIGHT study. *Blood* 2014;123:2944-2952.

Bendamustine + obinutuzumab

Marcus R, Davies A, Ando K, et al. Obinutuzumab for the first-line treatment of follicular lymphoma. *N Engl J Med* 2017;377:1331-1344.

RCHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone)

Czuczman MS, Weaver R, Alkuzweny B, et al. Prolonged clinical and molecular remission in patients with low-grade or follicular non-Hodgkin's lymphoma treated with rituximab plus CHOP chemotherapy: 9-year follow-up. *J Clin Oncol* 2004;22:4711-4716.

Hiddemann W, Kneba M, Dreyling M, et al. Frontline therapy with rituximab added to the combination of cyclophosphamide, doxorubicin, vincristine, and prednisone (CHOP) significantly improves the outcome for patients with advanced-stage follicular lymphoma compared with therapy with CHOP alone: results of a prospective randomized study of the German Low-Grade Lymphoma Study Group. *Blood* 2005;106:3725-3732.

CHOP + obinutuzumab

Marcus R, Davies A, Ando K, et al. Obinutuzumab for the first-line treatment of follicular lymphoma. *N Engl J Med* 2017;377:1331-1344.

RCVP (rituximab, cyclophosphamide, vincristine, prednisone)

Marcus R, Imrie K, Solal-Celigny P, et al. Phase III study of R-CVP compared with cyclophosphamide, vincristine, and prednisone alone in patients with previously untreated advanced follicular lymphoma. *J Clin Oncol* 2008;26:4579-4586.

CVP + obinutuzumab

Marcus R, Davies A, Ando K, et al. Obinutuzumab for the first-line treatment of follicular lymphoma. *N Engl J Med* 2017;377:1331-1344.

Rituximab

Hainsworth JD, Litchy S, Burris HA, III, et al. Rituximab as first-line and maintenance therapy for patients with indolent Non-Hodgkin's lymphoma. *J Clin Oncol* 2002;20:4261-4267.

Colombat P, Salles GA, Brousse N, et al. Rituximab (anti-CD20 monoclonal antibody) as single first-line therapy for patients with follicular lymphoma with a low tumor burden: Clinical and molecular evaluation. *Blood* 2001;97:101-106.

Martinelli G, Schmitz SF, Utiger U, et al. Long-term follow-up of patients with follicular lymphoma receiving single-agent rituximab at two different schedules in trial SAKK 35/98. *J Clin Oncol* 2010;28:4480-4484.

Ardeshtna KM, Qian W, Smith P, et al. Rituximab versus a watch-and-wait approach in patients with advanced-stage, asymptomatic, non-bulky follicular lymphoma: an open-label randomised phase 3 trial. *Lancet Oncol* 2014;15:424-435.

Lenalidomide + rituximab

Martin P, Jung SH, Pitcher B, et al. A phase II trial of lenalidomide plus rituximab in previously untreated follicular non-Hodgkin's lymphoma (NHL): CALGB 50803 (Alliance). *Ann Oncol* 2017;28:2806-2812.

Fowler N, Davis R, Rawal S, et al. Safety and activity of lenalidomide and rituximab in untreated indolent lymphoma: an open-label, phase 2 trial. *Lancet Oncol* 2014;15:1311-1318.

Morschhauser F, Fowler NH, Feugier P, et al. Rituximab plus Lenalidomide in Advanced Untreated Follicular Lymphoma. *N Engl J Med* 2018;379:934-947.

Ibritumomab tiuxetan

Scholz CW, Pinto A, Linkesch W, et al. (90)Yttrium-ibritumomab-tiuxetan as first-line treatment for follicular lymphoma: 30 months of follow-up data from an international multicenter phase II clinical trial. *J Clin Oncol* 2013;31:308-313.

First-line Consolidation or Extended Dosing

Chemoimmunotherapy followed by rituximab maintenance

Salles GA, Seymour JF, Offner F, et al. Rituximab maintenance for 2 years in patients with high tumour burden follicular lymphoma responding to rituximab plus chemotherapy (PRIMA): A phase 3, randomised controlled trial. *Lancet* 2011;377:42-51.

Salles GA, Seymour JF, Feugier P, et al. Long term follow-up of the PRIMA Study: Half of patients receiving rituximab maintenance remain progression free at 10 years [abstract]. *Blood* 2017;130 (Suppl 1):Abstract 486.

Extended dosing with rituximab

Ghielmini M, Schmitz SH, Cogliatti SB, et al. Prolonged treatment with rituximab in patients with follicular lymphoma significantly increases event-free survival and response duration compared with the standard weekly x 4 schedule. *Blood* 2004;103:4416-4423.

Obinutuzumab-based chemoimmunotherapy followed by obinutuzumab maintenance

Marcus R, Davies A, Ando K, et al. Obinutuzumab for the first-line treatment of follicular lymphoma. *N Engl J Med* 2017;377:1331-1344.

Ibritumomab tiuxetan

Morschhauser F, Radford J, Van Hoof A, et al. 90Yttrium-ibritumomab tiuxetan consolidation of first remission in advanced-stage follicular non-Hodgkin lymphoma: Updated results after a median follow-up of 7.3 years from the international, randomized, phase III first-line indolent trial. *J Clin Oncol* 2013;31:1977-1983.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

FOLL-B
3 OF 4



NCCN Guidelines Version 2.2020

Follicular Lymphoma (grade 1–2)

SUGGESTED TREATMENT REGIMENS REFERENCES

Second-line and Subsequent Therapy

Bendamustine + obinutuzumab

Sehn LH, Chua N, Mayer J, et al. Obinutuzumab plus bendamustine versus bendamustine monotherapy in patients with rituximab-refractory indolent non-Hodgkin lymphoma (GADOLIN): a randomised, controlled, open-label, multicentre, phase 3 trial. *Lancet Oncol* 2016;17:1081-1093.

Copanlisib

Dreyling M, Santoro A, Mollica L, et al. Long-term efficacy and safety from the copanlisib CHRONOS-1 study in patients with relapsed or refractory indolent B-cell lymphoma. *Blood* 2018;132:1595.

Dreyling M, Santoro A, Mollica L, et al. Phosphatidylinositol 3-kinase inhibition by copanlisib in relapsed or refractory indolent lymphoma. *J Clin Oncol* 2017;35:3898-3905.

Duvelisib

Flinn I, Miller C, Ardeshta K, et al. DYNAMO: A phase II study of duvelisib (IPI-145) in patients with refractory indolent non-Hodgkin lymphoma. *J Clin Oncol* 2019;37:912-922.

Idelalisib

Gopal A, Kahl B, De Vos S, et al. PI3Kδ inhibition by idelalisib in patients with relapsed indolent lymphoma. *N Engl J Med* 2014;370:1008-1018.

Salles G, Schuster SJ, de Vos S, et al. Efficacy and safety of idelalisib in patients with relapsed, rituximab- and alkylating agent-refractory follicular lymphoma: a subgroup analysis of a phase 2 study. *Haematologica* 2017;102:e156-e159.

Lenalidomide ± rituximab

Leonard JP, Jung SH, Johnson J, et al. Randomized trial of lenalidomide alone versus lenalidomide plus rituximab in patients with recurrent follicular lymphoma: CALGB 50401 (Alliance). *J Clin Oncol* 2015;33:3635-3640.

Witzig TE, Wiernik PH, Moore T, et al. Lenalidomide oral monotherapy produces durable responses in relapsed or refractory indolent non-Hodgkin's Lymphoma. *J Clin Oncol* 2009;27:5404-5409.

Leonard JP, Trneny M, Izutsu K, et al. AUGMENT: A Phase III Study of Lenalidomide Plus Rituximab Versus Placebo Plus Rituximab in Relapsed or Refractory Indolent Lymphoma. *J Clin Oncol* 2019;37:1188-1199.

Ibritumomab tiuxetan

Witzig TE, Flinn IW, Gordon LI, et al. Treatment with ibritumomab tiuxetan radioimmunotherapy in patients with rituximab-refractory follicular non-Hodgkin's lymphoma. *J Clin Oncol* 2002;20:3262-3269.

Witzig TE, Gordon LI, Cabanillas F, et al. Randomized controlled trial of yttrium-90-labeled ibritumomab tiuxetan radioimmunotherapy versus rituximab immunotherapy for patients with relapsed or refractory low-grade, follicular, or transformed B-cell non-Hodgkin's lymphoma. *J Clin Oncol* 2002;20:2453-2463.

Rituximab

McLaughlin P, Grillo-Lopez AJ, Link BK, et al. Rituximab chimeric anti-CD20 monoclonal antibody therapy for relapsed indolent lymphoma: half of patients respond to a four-dose treatment program. *J Clin Oncol* 1998;16:2825-2833.

Ghielmini M, Schmitz SH, Cogliatti SB, et al. Prolonged treatment with rituximab in patients with follicular lymphoma significantly increases event-free survival and response duration compared with the standard weekly x 4 schedule. *Blood* 2004;103:4416-4423.

Tazemetostat

Salles G, Adib D, Navia SB, et al. Phase 2 Multicenter Study of Tazemetostat, an EZH2 Inhibitor, in Patients with Relapsed or Refractory Follicular Lymphoma [abstract]. *Blood* 2019;134:Abstract 123.

CAR T-Cell Therapy

Axicabtagene ciloleucel

Neelapu SS, Locke FL, Bartlett NL, et al. Axicabtagene ciloleucel CAR T-cell therapy in refractory large B-cell lymphoma. *N Engl J Med* 2017;377:2531-2544.

Locke FL, Ghobadi A, Jacobson CA, et al. Long-term safety and activity of axicabtagene ciloleucel in refractory large B-cell lymphoma (ZUMA-1): a single-arm, multicentre, phase 1-2 trial. *Lancet Oncol* 2019;20:31-42.

Tisagenlecleucel

Schuster SJ, Bishop MR, Tam CS, et al. Tisagenlecleucel in adult relapsed or refractory diffuse large B-cell lymphoma. *N Engl J Med* 2019;380:45-56.

Second-line Consolidation or Extended Dosing

Rituximab maintenance

van Oers MHJ, Van Glabbeke M, Giurgea L, et al. Rituximab maintenance treatment of relapsed/resistant follicular non-hodgkin's lymphoma: Long-term outcome of the EORTC 20981 Phase III randomized Intergroup Study. *J Clin Oncol* 2010;28:2853-2858.

Obinutuzumab maintenance for rituximab refractory disease

Sehn LH, Chua N, Mayer J, et al. Obinutuzumab plus bendamustine versus bendamustine monotherapy in patients with rituximab-refractory indolent non-Hodgkin lymphoma (GADOLIN): a randomised, controlled, open-label, multicentre, phase 3 trial. *Lancet Oncol* 2016;17:1081-1093.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Marginal Zone Lymphomas



Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Extranodal Marginal Zone B-Cell Lymphoma

Gastric MALT Lymphoma

ADDITIONAL DIAGNOSTIC TESTING^{a,b}

ESSENTIAL:

- Diagnosis of gastric MALT lymphoma requires an endoscopic biopsy and an FNA is never adequate.
- Adequate immunophenotyping to establish diagnosis^{c,d}
 - IHC panel: CD20, CD3, CD5, CD10, BCL2, kappa/lambda, CD21 or CD23, cyclin D1,^f BCL6 with or without
 - Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: kappa/lambda, CD19, CD20, CD5, CD23, CD10
- Helicobacter pylori (H. pylori) stain (gastric), if positive, then PCR or FISH for t(11;18)^e

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- Molecular analysis to detect: antigen receptor gene rearrangements; *MYD88* mutation status to differentiate WM versus MZL if plasmacytic differentiation present
- Karyotype or FISH: t(1;14); t(3;14); t(11;14);^f t(11;18)
- FISH or PCR: t(14;18)

WORKUP

ESSENTIAL:

- Physical exam
- Performance status
- CBC with differential
- Comprehensive metabolic panel
- LDH
- If H. pylori negative by histopathology, then use noninvasive H. pylori testing (stool antigen test or urea breath test)
- Hepatitis B testing^g if rituximab contemplated
- Hepatitis C testing
- C/A/P CT with contrast of diagnostic quality
- Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)

USEFUL IN SELECTED CASES:

- Bone marrow biopsy ± aspirate
- PET/CT scan (especially if ISRT anticipated)
- Echocardiogram or MUGA scan if anthracycline or anthracenedione-based regimen is indicated
- Endoscopy with ultrasound (if available) with multiple biopsies of anatomical sites^h
- Discussion of fertility issues and sperm banking
- SPEP

→ [See Initial Therapy \(MALT-2\)](#)

^a Nondiagnostic atypical lymphoid infiltrates that are H. pylori positive should be rebiopsied to confirm or exclude lymphoma prior to treatment of H. pylori.

^b Any area of DLBCL should be treated according to the [NCCN Guidelines for Diffuse Large B-Cell Lymphoma \(BCL-1\)](#).

^c Typical immunophenotype: CD10-, CD5-, CD20+, cyclin D1-, BCL2- follicles.

^d See [Use of Immunophenotyping/Genetic Testing in Differential Diagnosis of Mature B-Cell and NK/T-Cell Neoplasms \(NHODG-A\)](#).

^e Locally advanced disease is more likely in patients with gastric MALT lymphoma with t(11;18), which is less likely to respond to antibiotics.

^f If IHC for cyclin D1 is positive, FISH for t(11;14) is not necessary; see [MANT-1](#).

^g Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen ([See NHODG-B](#)). If positive, check viral load and consider consult with gastroenterologist.

^h This is particularly useful for H. pylori-positive cases because the likelihood of tumor response is related to depth of tumor invasion.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Extranodal Marginal Zone B-Cell Lymphoma

Gastric MALT Lymphoma

STAGEⁱ**INITIAL THERAPY**

Stage I₁, or I₂^j
or Stage II₁^j
H. pylori positive
t(11;18) negative
or t(11;18)
unknown

Currently accepted antibiotic
therapy for H. pylori

Evaluate with endoscopy ([MALT-4](#))

See monoclonal antibody and
viral reactivation ([NHODG-B](#))

Stage I₁, or I₂^j
or Stage II₁^j
H. pylori positive,
t(11;18) positive^k

Currently accepted antibiotic
therapy for H. pylori + ISRT^m
(preferred)
or
Rituximab (if ISRT is
contraindicated)

Evaluate with endoscopy (MALT-5)

Stage I₁, or I₂^j
or Stage II₁^j
H. pylori negative

ISRT^{l,m} (preferred)
or
Rituximab (if ISRT is contraindicated)

Stage IIE, or II2
or Stage IV
(distant nodal,
advanced stage)

[See MALT-3](#)

ⁱ See Lugano Staging System for Gastrointestinal Lymphomas ([MALT-A](#)).

^j Involvement of submucosa or regional lymph nodes are much less likely to respond to antibiotic therapy. If there is persistent disease after evaluation, RT may be considered earlier in the course.

^k t(11;18) is a predictor for lack of tumor response (<5%) to antibiotics. Antibiotics are used in these patients to eradicate the H. pylori infection. These patients should be considered for alternative therapy of the lymphoma. Liu H, Ye H, Ruskone-Fourmestreaux A, et al. t(11;18) is a marker for all stage gastric MALT lymphomas that will not respond to H. pylori eradication. Gastroenterology 2002;122:1286-1294.

^l If H. pylori negative by both histology and serum antibodies, RT is recommended.

^m [See Principles of Radiation Therapy \(NHODG-D\)](#).

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Extranodal Marginal Zone B-Cell Lymphoma

Gastric MALT Lymphoma



ⁱ See Lugano Staging System for Gastrointestinal Lymphomas ([MALT-A](#)).

^m See [Principles of Radiation Therapy \(NHODG-D\)](#).

ⁿ Given incurability with conventional therapy, consider investigational therapy as first line of treatment.

^o Surgical resection is generally limited to specific clinical situations (ie, life-threatening hemorrhage).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020


Extranodal Marginal Zone B-Cell Lymphoma

Gastric MALT Lymphoma

3-MONTH RESTAGING AND FOLLOW-UP ENDOSCOPY

AFTER ANTIBIOTICS

ADDITIONAL THERAPY



^m See Principles of Radiation Therapy (NHODG-D).

^p Reassessment to rule out H. pylori by institutional standards. Biopsy to rule out large cell lymphoma. Any area of DLBCL should be treated according to the [NCCN Guidelines for Diffuse Large B-Cell Lymphoma \(BCEL-1\)](#).

^q If re-evaluation suggests slowly responding disease or asymptomatic nonprogression, continued observation may be warranted. Complete responses may be observed as early as 3 months after antibiotic treatment but can take longer to achieve (up to 18 months) (category 2B).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020


Extranodal Marginal Zone B-Cell Lymphoma

Gastric MALT Lymphoma

3- TO 6-MONTH RESTAGING AND FOLLOW-UP ENDOSCOPY

AFTER ISRT OR RITUXIMAB

ADDITIONAL THERAPY



^P Reassessment to rule out H. pylori by institutional standards. Biopsy to rule out large cell lymphoma. Any area of DLBCL should be treated according to the [NCCN Guidelines for Diffuse Large B-Cell Lymphoma \(BCEL-1\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 2.2020

Extranodal Marginal Zone B-Cell Lymphoma

Gastric MALT Lymphoma

FOLLOW-UP ENDOSCOPY

^m [See Principles of Radiation Therapy \(NHODG-D\).](#)^P Reassessment to rule out *H. pylori* by institutional standards. Biopsy to rule out large cell lymphoma. Any area of DLBCL should be treated according to the [NCCN Guidelines for Diffuse Large B-Cell Lymphoma \(BCEL-1\)](#).^r Optimal interval for follow-up endoscopy and imaging is not known. At NCCN Member Institutions, follow-up endoscopy and imaging using the modalities performed during workup is driven by symptoms.**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Extranodal Marginal Zone B-Cell Lymphoma

Gastric MALT Lymphoma

STAGING OF GASTRIC MALT LYMPHOMA: COMPARISON OF DIFFERENT SYSTEMS

Lugano Staging System for Gastrointestinal Lymphomas		Lugano Modification of Ann Arbor Staging System	TNM Staging System Adapted for Gastric Lymphoma	Tumor Extension
Stage I	Confined to GI tract ^a			
	I ₁ = mucosa, submucosa	I _E	T1 N0 M0	Mucosa, submucosa
	I ₂ = muscularis propria, serosa	I _E	T2 N0 M0	Muscularis propria
		I _E	T3 N0 M0	Serosa
Stage II	Extending into abdomen			
	II ₁ = local nodal involvement	II _E	T1-3 N1 M0	Perigastric lymph nodes
	II ₂ = distant nodal involvement	II _E	T1-3 N2 M0	More distant regional lymph nodes
Stage IIE	Penetration of serosa to involve adjacent organs or tissues	II _E	T4 N0 M0	Invasion of adjacent structures
Stage IV ^b	Disseminated extranodal involvement or concomitant supradiaphragmatic nodal involvement		T1-4 N3 M0	Lymph nodes on both sides of the diaphragm/ distant metastases (eg, bone marrow or additional extranodal sites)
		IV	T1-4 N0-3 M1	

Zucca E, Bertoni F, Yahalom J, Isaacson P. Extranodal Marginal Zone B-cell Lymphoma of Mucosa-Associated Lymphoid Tissue (MALT lymphoma) in Armitage et al eds. Non-Hodgkin's Lymphomas. Philadelphia: Lippincott, 2010:242. (<http://www.com>)

^a Single primary or multiple, noncontiguous.

^b Involvement of multiple extranodal sites in MALT lymphoma appears to be biologically distinct from multiple extranodal involvement in other lymphomas, and these patients may be managed by treating each site separately with excision or RT. In contrast, cases with disseminated nodal involvement appear to behave more like nodal marginal zone lymphoma or like disseminated follicular lymphoma.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

National
Comprehensive
Cancer
Network®**NCCN Guidelines Version 2.2020****Extranodal Marginal Zone B-Cell Lymphoma****Nongastric MALT Lymphoma (Noncutaneous)**[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)**ADDITIONAL DIAGNOSTIC TESTING^{a,b}****ESSENTIAL:**

- Adequate immunophenotyping to establish diagnosis^{c,d}
 - ▶ IHC panel: CD20, CD3, CD5, CD10, BCL2, kappa/lambda, CD21 or CD23, cyclin D1 with or without
 - ▶ Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: kappa/lambda, CD19, CD20, CD5, CD23, CD10

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- Molecular analysis to detect: antigen receptor gene rearrangements; *MYD88* mutation status to differentiate WM versus MZL if plasmacytic differentiation present; PCR for t(11;18)
- Karyotype or FISH: t(11;18), t(11;14), t(3;14)
- FISH or PCR: t(14;18)

WORKUP**ESSENTIAL:**

- Physical exam with attention to nongastric sites^a
- Performance status
- CBC with differential
- Comprehensive metabolic panel
- LDH
- Hepatitis B testing^e if rituximab contemplated
- Hepatitis C testing
- C/A/P CT and other suspected sites with contrast of diagnostic quality
- Whole-body PET/CT scan essential if RT for stage I, II disease or systemic therapy is planned
- Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)

USEFUL IN SELECTED CASES:

- Echocardiogram or MUGA scan if anthracycline or anthracenedione-based regimen is indicated
- Bone marrow biopsy ± aspirate
- Endoscopy with multiple biopsies of anatomical sites^f
- MRI with contrast for neurologic evaluation or if CT with contrast is contraindicated
- MRI of head/neck, cranial, and ocular adnexa
- Autoimmune disease testing, particularly Sjogren's
- Discussion of fertility issues and sperm banking
- SPEP

[See Initial
Therapy
\(NGMLT-2\)](#)

^a Typical non-gastric sites include the following: bowel (small and large), breast, head and neck, lung, dural, ocular adnexa, ovary, parotid, prostate, and salivary gland. Infectious agents have been reported to be associated with many nongastric sites, but testing for these infectious organisms is not required for management in the United States.

^b This guideline pertains to noncutaneous; for primary cutaneous marginal zone lymphoma, [see NCCN Guidelines for Primary Cutaneous B-Cell Lymphomas](#).

^c Typical immunophenotype: CD10-, CD5-, CD20+, CD23-/+, CD43-/+, cyclin D1-, BCL2- follicles.

^d [See Use of Immunophenotyping/Genetic Testing in Differential Diagnosis of Mature B-Cell and NK/T-Cell Neoplasms \(NHODG-A\)](#).

^e Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen ([See NHODG-B](#)). If positive, check viral load and consider consult with gastroenterologist.

^f In cases where primary site is thought to be in head/neck or lungs, upper GI endoscopy should be considered.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2020


Extranodal Marginal Zone B-Cell Lymphoma

Nongastric MALT Lymphoma (Noncutaneous)

[NCCN Guidelines Index](#)

[Table of Contents](#)

[Discussion](#)



^g Based on anecdotal responses to antibiotics in ocular and cutaneous marginal zone lymphomas, some physicians will give an empiric course of doxycycline prior to initiating other therapy.

^h [See Principles of Radiation Therapy \(NHODG-D\)](#).

ⁱ Surgical excision for adequate diagnosis may be appropriate treatment for disease.

^j Observation may be considered for patients whose diagnostic biopsy was excisional, or where RT could result in significant morbidity.

^k Definitive treatment of multiple sites may be indicated (eg, bilateral orbital disease without evidence of disease elsewhere) or palliative treatment of symptomatic sites.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 2.2020

Extranodal Marginal Zone B-Cell Lymphoma

Nongastric MALT Lymphoma (Noncutaneous)

STAGE FOLLOW-UP



^h See Principles of Radiation Therapy (NHODG-D).

^l Follow-up includes diagnostic tests and imaging previously used as clinically indicated.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Nodal Marginal Zone Lymphoma

ADDITIONAL DIAGNOSTIC TESTING^a

ESSENTIAL:

- Adequate immunophenotyping to establish diagnosis^{b,c}
 - ▶ IHC panel: CD20, CD3, CD5, CD10, BCL2, kappa/lambda, CD21 or CD23, cyclin D1 with or without
 - ▶ Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: kappa/lambda, CD19, CD20, CD5, CD23, CD10
- Pediatric NMZL should be considered with localized disease in a young patient.

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- Molecular analysis to detect: antigen receptor gene rearrangements; *MYD88* mutation status to differentiate WM versus MZL if plasmacytic differentiation present; PCR for t(11;18)
- Karyotype or FISH: t(11;18), t(1;14), del(13q), del(7q)
- FISH or PCR: t(14;18)

WORKUP

ESSENTIAL:

- Physical exam with performance status
- CBC with differential
- Comprehensive metabolic panel
- LDH
- Hepatitis B testing^d if rituximab contemplated
- Hepatitis C testing
- C/A/P CT or other suspected sites with contrast of diagnostic quality
- Whole-body PET/CT scan essential if RT for stage I, II disease or systemic therapy is planned
- Bone marrow biopsy + aspirate to document clinical stage I–II disease^e
- Evaluation to rule out extranodal primary sites
 - ▶ Neck nodes: ocular, parotid, thyroid, and salivary gland
 - ▶ Axillary nodes: lung, breast, and skin
 - ▶ Mediastinal/hilar nodes: lung
 - ▶ Abdominal nodes: splenic and GI
 - ▶ Inguinal/iliac nodes: GI and skin
- Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)

USEFUL IN SELECTED CASES:

- Echocardiogram or MUGA scan if anthracycline or anthracenedione-based regimen is indicated
- Additional imaging as appropriate
- Discussion of fertility issues and sperm banking
- SPEP

Stage I, II
See [NODE-2](#)

Stage III, IV
See [NODE-3](#)

^a NMZL is rare and occurs most commonly as spread from extranodal MALT; must also be distinguished from nodal FL, MCL, lymphoplasmacytic lymphoma, and CLL, all of which are more common.

^b Typical immunophenotype: CD10-, CD5-, CD20+, CD23-/+, CD43-/+, and cyclin D1-, BCL2- follicles.

^c See [Use of Immunophenotyping/Genetic Testing in Differential Diagnosis of Mature B-Cell and NK/T-Cell Neoplasms \(NHODG-A\)](#).

^d Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen ([See NHODG-B](#)). If positive, check viral load and consider consult with gastroenterologist.

^e Bilateral or unilateral provided core biopsy is >2 cm. If ibritumomab tiuxetan is considered, bilateral cores are recommended and the pathologist should provide the percent of overall cellular elements and the percent of cellular elements involved in the marrow. If observation is initial therapy, bone marrow biopsy may be deferred.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Nodal Marginal Zone Lymphoma



^f See [Principles of Radiation Therapy \(NHODG-D\)](#).

^g Observation may be appropriate in circumstances where potential toxicity of ISRT or systemic therapy outweighs potential clinical benefit in consultation with a radiation oncologist.

^h Anti-CD20 monoclonal antibodies include rituximab or obinutuzumab. Obinutuzumab is not indicated as single-agent therapy.

ⁱ Initiation of chemotherapy or more extended RT can improve FFS, but has not been shown to improve overall survival. These are options for therapy.

^j See [Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#). PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS).

^k Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated. Imaging should be performed whenever there are clinical indications. For surveillance imaging, see [Discussion](#) for consensus imaging recommendations.

^l Consider possibility of histologic transformation in patients with progressive disease, especially if LDH levels are rising, single site is growing disproportionately, extranodal disease develops, or there are new B symptoms. If clinical suspicion of transformation, FDG-PET may help identify areas suspicious for transformation. FDG-PET scan demonstrating marked heterogeneity or sites of intense FDG avidity may indicate transformation, and biopsy should be directed biopsy at the most FDG-avid area. Functional imaging does not replace biopsy to diagnose transformation. If transformation is histologically confirmed, treat with anthracycline-based therapy. See Management of Transformation (NODE-5).


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020


Nodal Marginal Zone Lymphoma

STAGE**MANAGEMENT AND FOLLOW-UP^k**^f See Principles of Radiation Therapy (NHODG-D).^j See Lugano Response Criteria for Non-Hodgkin Lymphoma (NHODG-C). PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS).^k Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated. Imaging should be performed whenever there are clinical indications. For surveillance imaging, see Discussion for consensus imaging recommendations.^l Consider possibility of histologic transformation in patients with progressive disease, especially if LDH levels are rising, single site is growing disproportionately, extranodal disease develops, or there are new B symptoms. If clinical suspicion of transformation, FDG-PET may help identify areas suspicious for transformation. FDG-PET scan demonstrating marked heterogeneity or sites of intense FDG avidity may indicate transformation, and biopsy should be directed biopsy at the most FDG-avid area. Functional imaging does not replace biopsy to diagnose transformation. If transformation is histologically confirmed, treat with anthracycline-based therapy. See Management of Transformation (NODE-5).^m See GELF criteria (FOLL-A).**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Nodal Marginal Zone Lymphoma



^f See Principles of Radiation Therapy (NHODG-D).

^j See Lugano Response Criteria for Non-Hodgkin Lymphoma (NHODG-C). PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS).

^k Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated. Imaging should be performed whenever there are clinical indications. For surveillance imaging, see Discussion for consensus imaging recommendations.

^l Consider possibility of histologic transformation in patients with progressive disease, especially if LDH levels are rising, single site is growing disproportionately, extranodal disease develops, or there are new B symptoms. If clinical suspicion of transformation, FDG-PET may help identify areas suspicious for transformation. FDG-PET scan demonstrating marked heterogeneity or sites of intense FDG avidity may indicate transformation, and biopsy should be directed biopsy at the most FDG-avid area. Functional imaging does not replace biopsy to diagnose transformation. If transformation is histologically confirmed, treat with anthracycline-based therapy. See Management of Transformation (NODE-5).

^m See GELF criteria (FOLL-A).

ⁿ A PET-positive PR is associated with a shortened PFS (see Discussion); however, additional treatment at this juncture has not been shown to change outcome.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Nodal Marginal Zone Lymphoma

HISTOLOGIC TRANSFORMATION TO DIFFUSE LARGE B-CELL LYMPHOMA


^f [See Principles of Radiation Therapy \(NHODG-D\)](#).^j [See Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#). PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS).^o Perform FISH for *BCL6* rearrangement and *MYC* rearrangements.^p ISRT alone or one course of single-agent therapy including rituximab.^q NGS may be useful for treatment selection.^r This includes ≥2 of chemoimmunotherapy regimens for indolent or transformed disease. For example, prior treatment with BR and RCHOP.^s Consider ISRT for localized presentations, bulky disease, and/or limited osseous disease.^t If transformation is coexisting with extensive MZL, consider maintenance ([see NODE-4, Optional Extended Therapy](#)).^u If proceeding to an autologous stem cell rescue, consider additional systemic therapy ± ISRT to induce CR prior to transplant.^v Repeat biopsy should be strongly considered if PET-positive prior to additional therapy, because PET positivity may represent post-treatment inflammation. Patients with a durable response for transformed disease may recur with the original indolent lymphoma. In that case, the management should be as per [NODE-4](#). If biopsy negative, follow CR pathway.**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



NCCN Guidelines Version 2.2020

Nodal Marginal Zone Lymphoma

HISTOLOGIC TRANSFORMATION TO DIFFUSE LARGE B-CELL LYMPHOMA

^f See Principles of Radiation Therapy (NHODG-D).^j See Lugano Response Criteria for Non-Hodgkin Lymphoma (NHODG-C).^o PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS).^r FISH for *BCL6* rearrangement and *MYC* rearrangements.^r This includes ≥2 of chemoimmunotherapy regimens for indolent or transformed disease. For example, prior treatment with BR and RCHOP.^s Consider ISRT for localized presentations, bulky disease, and/or limited osseous disease.^u If proceeding to an autologous stem cell rescue, consider additional systemic therapy ± ISRT to induce CR prior to transplant.^v Repeat biopsy should be strongly considered if PET-positive prior to additional therapy because PET positivity may represent post-treatment inflammation. Patients with a durable response for transformed disease may recur with the original indolent lymphoma. In that case, the management should be as per [NODE-4](#). If biopsy negative, follow CR pathway.^w See Guidance for Treatment of Patients with Chimeric Antigen Receptor (CAR) T-Cell Therapy (BCEL-D).^x Patients should have received at least one anthracycline or anthracenedione-based regimen, unless contraindicated.**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



NCCN Guidelines Version 2.2020

Splenic Marginal Zone Lymphoma

ADDITIONAL DIAGNOSTIC TESTING^a

ESSENTIAL:

- Adequate immunophenotyping to establish diagnosis^{b,c}
 - IHC panel: CD20, CD3, CD5, CD10, BCL2, kappa/lambda, CD21 or CD23, cyclin D1, IgD, CD43, annexin A1; with or without
 - Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: kappa/lambda, CD19, CD20, CD5, CD23, CD10, CD43, CD103

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- Molecular analysis to detect: antigen receptor gene rearrangements; *MYD88* mutation status to differentiate WM versus MZL if plasmacytic differentiation present; *BRAF* mutation status to differentiate MZL from HCL by IHC or sequencing; PCR for t(11;18)
- Karyotype or FISH: CLL panel; t(11;18), t(11;14), del(7q)
- FISH or PCR: t(14;18)

WORKUP

ESSENTIAL:

- Physical exam with performance status
- CBC with differential
- Comprehensive metabolic panel
- LDH
- Hepatitis B testing^d if rituximab contemplated
- Hepatitis C testing
- C/A/P CT or other suspected sites with contrast of diagnostic quality
- Bone marrow biopsy ± aspirate
- SPEP and/or quantitative immunoglobulin levels
- Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)

USEFUL IN SELECTED CASES:

- Whole-body PET/CT scan
- Additional imaging as appropriate
- Discussion of fertility issues and sperm banking
- Immunofixation of blood (for elevated immunoglobulins or positive SPEP)
- Cryoglobulins
- Direct Coombs testing

→ [See Management \(SPLN-2\)](#)

^a SMZL is most definitively diagnosed at splenectomy, since the immunophenotype is nonspecific and morphologic features on the bone marrow may not be diagnostic. However, the diagnosis of SMZL may be made on the basis of bone marrow ± peripheral blood involvement by small lymphoid cells with immunoglobulin (Ig) light chain restriction that lack characteristic features of other small B-cell neoplasms (CD5, CD10, cyclin D1). Plasmacytoid differentiation with cytoplasmic Ig detectable on paraffin sections may occur. In such cases, the differential diagnosis may include lymphoplasmacytic lymphoma. With a characteristic intrasinusoidal lymphocytic infiltration of the bone marrow, the diagnosis can strongly be suggested on bone marrow biopsy alone, if the immunophenotype is consistent.

^b Typical immunophenotype: CD10-, CD5-, CD20+, CD23-/+, CD43-/+, and cyclin D1-, BCL2- follicles, annexin A1, and CD103- (distinction from hairy cell leukemia) with expression of both IgM and IgD.

^c [See Use of Immunophenotyping/Genetic Testing in Differential Diagnosis of Mature B-Cell and NK/T-Cell Neoplasms \(NHODG-A\).](#)

^d Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen ([See NHODG-B](#)). If positive, check viral load and consider consult with gastroenterologist.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Splenic Marginal Zone Lymphoma

CLINICAL PRESENTATION

MANAGEMENT


FOLLOW-UP

Asymptomatic,
without progressive
cytopenia, no
splenomegaly

Observe

Consider prophylaxis for tumor
lysis syndrome ([See NHODG-B](#))

See monoclonal antibody and viral
reactivation ([NHODG-B](#))



[See Follow-up \(SPLN-3\)](#)

^e If there is hepatic involvement and hepatitis C positive, treat with an appropriate regimen for hepatitis C.

^f Tsimberidou AM, Catovsky D, Schlette E, et al. Outcomes in patients with splenic marginal zone lymphoma and marginal zone lymphoma treated with rituximab with or without chemotherapy or chemotherapy alone. Cancer 2006;107:125-135.

^g Pneumococcal, meningococcal, and hepatitis B vaccinations should be given at least 2 weeks before splenectomy.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Splenic Marginal Zone Lymphoma

FOLLOW-UP



^h Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Marginal Zone Lymphomas

SUGGESTED TREATMENT REGIMENS^{a,b,c}

An FDA-approved biosimilar is an appropriate substitute for rituximab.

First-line Therapy

- Preferred regimens (in alphabetical order)
 - Bendamustine + rituximab
 - RCHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone)
 - RCVP (rituximab, cyclophosphamide, vincristine, prednisone)
 - Rituximab (375 mg/m² weekly for 4 doses) for SMZL
- Other recommended regimens (in alphabetical order)
 - Ibritumomab tiuxetan^d (category 2B)
 - Lenalidomide + rituximab (category 2B)
 - Rituximab (375 mg/m² weekly for 4 doses) for extranodal (MALT) and nodal MZL

First-line Extended Therapy (optional)

- If initially treated with single-agent rituximab, consolidation with rituximab 375 mg/m² one dose every 12 weeks for up to 2 years

[See Second-line and Subsequent Therapy on MZL-A 2 of 3](#)

First-line Therapy for Elderly or Infirm (if none of the above is expected to be tolerable in the opinion of treating physician)

- Preferred regimen
 - Rituximab (preferred) (375 mg/m² weekly for 4 doses)
- Other recommended regimens
 - Chlorambucil ± rituximab
 - Cyclophosphamide ± rituximab

Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

^a See references for regimens ([MZL-A 3 of 3](#)).

^b The choice of initial therapy requires consideration of many factors, including age, comorbidities, and future treatment possibilities (eg, HDT with ASCR). Therefore, treatment selection is highly individualized.

^c Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^d Selection of patients requires adequate marrow cellularity >15% and <25% involvement of lymphoma in bone marrow, and platelets >100,000. In patients with prior autologous stem cell rescue, referral to a tertiary care center is highly recommended for ibritumomab tiuxetan. If ibritumomab tiuxetan is considered, bilateral cores are recommended and the pathologist should provide the percent of overall cellular elements and the percent of cellular elements involved in the marrow. As of 2010, updates suggest a trend towards an increased risk of MDS with RIT. Cytogenetics/FISH assessment for MDS markers is recommended for patients receiving RIT.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Marginal Zone Lymphomas

SUGGESTED TREATMENT REGIMENS^{a,b,c}

An FDA-approved biosimilar is an appropriate substitute for rituximab.

Second-line and Subsequent Therapy

- Preferred regimens (in alphabetical order)
 - Bendamustine + obinutuzumab (not recommended if treated with prior bendamustine)
 - Bendamustine + rituximab (not recommended if treated with prior bendamustine) (may be considered for both nodal MZL and extranodal [MALT] lymphoma)
 - Ibrutinib^e
 - Lenalidomide + rituximab (may be considered for both nodal MZL and extranodal [MALT] lymphoma)
 - RCHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone)
 - RCVP (rituximab, cyclophosphamide, vincristine, prednisone)
- Other recommended regimens (in alphabetical order)
 - Ibritumomab tiuxetan^d (category 2B)
 - PI3K inhibitors (relapsed/refractory after 2 prior therapies)^e
 - ◊ Copanlisib
 - ◊ Duvelisib
 - ◊ Idelalisib
 - Rituximab (if longer duration of remission)

Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

^a See references for regimens ([MZL-A 3 of 3](#)).

^b The choice of initial therapy requires consideration of many factors, including age, comorbidities, and future treatment possibilities (eg, HDT with ASCR). Therefore, treatment selection is highly individualized.

^c Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^d Selection of patients requires adequate marrow cellularity >15% and <25% involvement of lymphoma in bone marrow, and platelets >100,000. In patients with prior autologous stem cell rescue, referral to a tertiary care center is highly recommended for ibritumomab tiuxetan. If ibritumomab tiuxetan is considered, bilateral cores are recommended and the pathologist should provide the percent of overall cellular elements and the percent of cellular elements involved in the marrow. As of 2010, updates suggest a trend towards an increased risk of MDS with RIT. Cytogenetics/FISH assessment for MDS markers is recommended for patients receiving RIT.

^e [See Special Considerations for the Use of Small-Molecule Inhibitors \(NHODG-E\).](#)

^f [See Guidance for Treatment of Patients with Chimeric Antigen Receptor \(CAR\) T-Cell Therapy \(BCEL-D\).](#)

^g Patients should have received at least one anthracycline or anthracenedione-based regimen, unless contraindicated.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

Second-line and Subsequent Therapy for Elderly or Infirm (if none of the above is expected to be tolerable in the opinion of treating physician)

- Preferred regimens (in alphabetical order)
 - Ibrutinib^e
 - Lenalidomide + rituximab
 - Rituximab (375 mg/m² weekly for 4 doses)
- Other recommended regimens
 - Chlorambucil ± rituximab
 - Cyclophosphamide ± rituximab

Second-line Consolidation or Extended Dosing (optional)

- Preferred regimen
 - If treated with bendamustine + obinutuzumab for recurrent disease then obinutuzumab maintenance for rituximab-refractory disease (1 g every 8 weeks for total of 12 doses)
- Other recommended regimens
 - High-dose therapy with autologous stem cell rescue
 - Allogeneic hematopoietic cell transplant for highly selected patients

Histologic Transformation of Nodal MZL to DLBCL

- Anti-CD19 CAR T-Cell therapy (only after ≥2 prior chemoimmunotherapy regimens)^{f,g}
 - Axicabtagene ciloleucel
 - Tisagenlecleucel



NCCN Guidelines Version 2.2020

Marginal Zone Lymphomas

SUGGESTED TREATMENT REGIMENS

REFERENCES

First-line Therapy

Chlorambucil ± rituximab

Zucca E, Conconi A, Martinelli G, et al. Final results of the IELSG-19 randomized trial of mucosa-associated lymphoid tissue lymphoma: Improved event-free and progression-free survival with rituximab plus chlorambucil versus either chlorambucil or rituximab monotherapy. *J Clin Oncol* 2017;35:1905-1912.

RCHOP/RCVP/BR

Rummel MJ, Niederle N, Maschmeyer G, et al. Bendamustine plus rituximab versus CHOP plus rituximab as first-line treatment for patients with indolent and mantle-cell lymphomas: an open-label, multicentre, randomised, phase 3 non-inferiority trial. *Lancet* 2013;381:1203-1210.

Flinn IW, van der Jagt R, Kahl BS, et al. Randomized trial of bendamustine-rituximab or R-CHOP/R-CVP in first-line treatment of indolent NHL or MCL: the BRIGHT study. *Blood* 2014;123:2944-2952.

Salar A, Domingo-Domenech E, Panizo C, et al. Long-term results of a phase 2 study of rituximab and bendamustine for mucosa-associated lymphoid tissue lymphoma. *Blood* 2017;130:1772-1774.

Ibritumomab tiuxetan

Lossos IS, Fabregas JC, Koru-Sengul T, et al. Phase II study of (90)Y Ibritumomab tiuxetan in patients with previously untreated marginal zone lymphoma. *Leuk Lymphoma* 2015;56:1750-1755.

Lenalidomide + rituximab

Fowler NH, Davis RE, Rawal S, et al. Safety and activity of lenalidomide and rituximab in untreated indolent lymphoma: an open-label, phase 2 trial. *Lancet Oncol* 2014;15:1311-1318.

Rituximab (preferred for SMZL)

Tsimberidou AM, Catovsky D, Schlette E, et al. Outcomes in patients with splenic marginal zone lymphoma and marginal zone lymphoma treated with rituximab with or without chemotherapy or chemotherapy alone. *Cancer* 2006;107:125-135.

Else M, Marin-Niebla A, de la Cruz F, et al. Rituximab, used alone or in combination, is superior to other treatment modalities in splenic marginal zone lymphoma. *Br J Haematol* 2012;159:322-328.

Kalpadakis C, Pangalis GA, Angelopoulou MK, et al. Treatment of splenic marginal zone lymphoma with rituximab monotherapy: progress report and comparison with splenectomy. *Oncologist* 2013;18:190-197.

First-line Extended Therapy (optional)

Extended dosing with rituximab

Williams ME, Hong F, Gascoyne RD, et al. Rituximab extended schedule or retreatment trial for low tumour burden non-follicular indolent B-cell non-Hodgkin lymphomas: Eastern Cooperative Oncology Group Protocol E4402. *Br J Haematol* 2016;173:867-875.

Second-line and Subsequent Therapy

Bendamustine + obinutuzumab

Sehn LH, Chua N, Mayer J, et al. Obinutuzumab plus bendamustine versus bendamustine monotherapy in patients with rituximab-refractory indolent non-Hodgkin lymphoma (GADOLIN): a randomised, controlled, open-label, multicentre, phase 3 trial. *Lancet Oncol* 2016;17:1081-1093.

Copanlisib

Dreyling M, Panayiotidis P, Egyed M, et al. Efficacy of copanlisib monotherapy in patients with relapsed or refractory marginal zone lymphoma: subset analysis from the CHRONOS-1 trial [abstract]. *Blood* 2017;130:Abstract 4053.

Duvelisib

Flinn IW, Miller CB, Ardeschna KM, et al. DYNAMO: A phase II study of duvelisib (IPI-145) in patients with refractory indolent non-Hodgkin lymphoma. *J Clin Oncol* 2019; 37:912 922.

Ibritumomab tiuxetan

Vanazzi A, Grana C, Crosta C, et al. Efficacy of (9)(0)Yttrium-ibritumomab tiuxetan in relapsed/refractory extranodal marginal-zone lymphoma. *Hematol Oncol* 2014;32:10-15.

Ibrutinib

Noy A, de Vos S, Thieblemont C, et al. Targeting Bruton tyrosine kinase with ibrutinib in relapsed/refractory marginal zone lymphoma. *Blood* 2017;129:2224-2232.

Idelalisib

Gopal A, Kahl B, De Vos S, et al. PI3Kδ inhibition by idelalisib in patients with relapsed indolent lymphoma. *N Engl J Med* 2014;370:1008-1018.

Lenalidomide + rituximab

Witzig TE, Wiernik PH, Moore T, et al. Lenalidomide oral monotherapy produces durable responses in relapsed or refractory indolent non-Hodgkin's Lymphoma. *J Clin Oncol* 2009;27:5404-5409.

Sacchi S, Marcheselli R, Bari A, et al. Safety and efficacy of lenalidomide in combination with rituximab in recurrent indolent non-follicular lymphoma: final results of a phase II study conducted by the Fondazione Italiana Linfomi. *Haematologica* 2016;101:e196.

Second-line Consolidation or Extended Dosing (optional)

Obinutuzumab maintenance for rituximab refractory disease

Sehn LH, Chua N, Mayer J, et al. Obinutuzumab plus bendamustine versus bendamustine monotherapy in patients with rituximab-refractory indolent non-Hodgkin lymphoma (GADOLIN): a randomised, controlled, open-label, multicentre, phase 3 trial. *Lancet Oncol* 2016;17:1081-1093.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Mantle Cell Lymphoma

ADDITIONAL DIAGNOSTIC TESTING

ESSENTIAL:

- Adequate immunophenotyping to establish diagnosis^{a,b}
 - ▶ IHC panel: CD20, CD3, CD5, cyclin D1, CD10, CD21, CD23, BCL2, BCL6, TP53, Ki-67^c with or without
 - ▶ Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: kappa/lambda, CD19, CD20, CD5, CD23, CD10
- TP53 sequencing (patients with typical MCL with an expected aggressive clinical course or particularly if upfront transplant anticipated)^e

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- IHC: LEF1 may help distinguish from variant CLL; SOX11 or IGHV sequencing may be useful for determination of clinically indolent^d MCL; may also help in diagnosis of CCND1- MCL.
- Karyotype or FISH: t(11;14), t(14;18), CLL panel
- Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: CD200

WORKUP

ESSENTIAL:

- Physical exam: Attention to node-bearing areas, including Waldeyer's ring, and to size of liver and spleen
- Performance status
- B symptoms
- CBC with differential
- Comprehensive metabolic panel
- LDH
- C/A/P CT with contrast of diagnostic quality
- Whole-body PET/CT scan essential if RT for stage I, II disease or systemic therapy is planned
- Hepatitis B testing^f if rituximab contemplated
- Echocardiogram or MUGA scan if anthracycline or anthracenedione-based regimen is indicated
- Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- Endoscopy/colonoscopy^g
- Bone marrow biopsy ± aspirate
- Neck CT with contrast
- Uric acid
- Beta-2-microglobulin
- Hepatitis C testing
- Lumbar puncture (for blastic variant or CNS symptoms)
- Discussion of fertility issues and sperm banking

[See Stage I, II Induction Therapy \(MANT-2\)](#)

[See Stage II bulky, III, IV Induction Therapy \(MANT-3\)](#)

^a Typical immunophenotype: CD5+, CD20+, CD43+, CD23-/+ , cyclin D1+, CD10-/+.

Note: Some cases of MCL may be CD5- or CD23+. If the diagnosis is suspected, cyclin D1 staining or FISH for t(11;14) should be done. There are rare cases of CCND1- MCL (<5%) with an otherwise typical immunophenotype.

^b [See Use of Immunophenotyping/Genetic Testing in Differential Diagnosis of Mature B-Cell and NK/T-Cell Neoplasms \(NHODG-A\).](#)

^c Ki-67 proliferation fraction of <30% in lymph nodes is associated with a more favorable prognosis.

^d Most common biomarker for indolent disease: SOX11- [IGHV mutated]. Typical clinical presentation: leukemic non-nodal CLL-like with splenomegaly, low tumor burden, Ki-67 proliferation fraction <10%.

^e TP53 mutation has been associated with poor prognosis in patients treated with conventional therapy, including transplant. Clinical trial is strongly suggested for these patients.

^f Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen ([See NHODG-B](#)). If positive, check viral load and consider consult with gastroenterologist.

^g Essential for confirmation of stage I–II disease. [See Discussion](#) for details.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Mantle Cell Lymphoma



Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))

See monoclonal antibody and viral reactivation ([NHODG-B](#))

^h Localized presentation is extremely rare.

ⁱ Early referral for high-dose therapy with stem cell rescue is advisable for planning purposes.

^j [See Principles of Radiation Therapy \(NHODG-D\)](#).

^k Leitch HA, Gascoyne RD, Chhanabhai M, et al. Limited-stage mantle-cell lymphoma. Ann Oncol 2003;14:1555-1561.

^l [See Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#).

^m Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Mantle Cell Lymphoma



^e *TP53* mutation has been associated with poor prognosis in patients treated with conventional therapy, including transplant. Clinical trial is strongly suggested for these patients.

ⁱ Early referral for high-dose therapy with stem cell rescue is advisable for planning purposes.

^j See [Principles of Radiation Therapy \(NHODG-D\)](#).

^l See [Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#).

^m Follow-up includes diagnostic tests and imaging using the same modalities performed during workup as clinically indicated.

ⁿ Patients who have achieved near CR can proceed to HDT/ASCR. Patients who have achieved minimal PR with substantial disease should be treated as having stable, refractory disease. Patients who have achieved a very good PR may be treated with additional therapy to achieve CR with the goal of proceeding to HDT/ASCR.

^o Patients who have achieved a very good PR or better can be observed or consider rituximab maintenance. Patients who have achieved minimal PR with substantial disease should be treated as having stable, refractory disease.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Mantle Cell Lymphoma



^e *TP53* mutation has been associated with poor prognosis in patients treated with conventional therapy, including transplant. Clinical trial is strongly suggested for these patients.

^P The description represents the most common indolent presentation; however, there are some patients with GI or blood/bone marrow involvement only, which may express SOX11 and have an indolent course.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



SUGGESTED TREATMENT REGIMENS^{a,b}

An FDA-approved biosimilar is an appropriate substitute for rituximab.

Induction Therapy

• Aggressive therapy

▶ Preferred regimens

- ◊ RDHA (rituximab, dexamethasone, cytarabine) + platinum (carboplatin, cisplatin, or oxaliplatin)
- ◊ Alternating RCHOP/RDHAP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone)/(rituximab, dexamethasone, cytarabine, cisplatin)
- ◊ NORDIC regimen (dose-intensified induction immunochemotherapy with rituximab + cyclophosphamide, vincristine, doxorubicin, prednisone [maxi-CHOP]) alternating with rituximab + high-dose cytarabine
- ◊ HyperCVAD (cyclophosphamide, vincristine, doxorubicin, and dexamethasone alternating with high-dose methotrexate and cytarabine) + rituximab^c (NOTE: There are conflicting data regarding the need for consolidation with HDT/ASCR.)

▶ Other recommended regimen

- ◊ Bendamustine^d + rituximab (category 2B)

Consolidation After Aggressive Therapy

- High-dose therapy followed by autologous stem cell rescue

Maintenance After HDT/ASCR

- Maintenance rituximab every 8 weeks x 3 y (category 1)

• Less aggressive therapy

▶ Preferred

- ◊ Bendamustine + rituximab
 - ◊ VR-CAP (bortezomib, rituximab, cyclophosphamide, doxorubicin, and prednisone)
 - ◊ RCHOP^e
 - ◊ Lenalidomide + rituximab
- ###### ▶ Other recommended regimen
- ◊ Modified rituximab-HyperCVAD in patients older than 65 y
 - ◊ RBAC500 (rituximab, bendamustine, cytarabine)

Maintenance After Less Aggressive Therapy

- Rituximab every 8 weeks until progression or intolerance (category 1 following RCHOP; 2–5 y following modified rituximab-HyperCVAD)
 - ▶ Prospective trial data suggest no benefit after BR
 - ▶ Untested after VR-CAP, RBAC500

[See Second-line Therapy on MANT-A 2 of 4](#)

Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

^a See references for regimens [MANT-A 3 of 4](#) and [MANT-A 4 of 4](#).

^b Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^c Rituximab + ibrutinib can be used as a pre-treatment to limit the number of cycles of RHyperCVAD/rituximab maintenance. Wang ML, Lee H, Thirumurthi S, et al. Hematol Oncol 2017;35:142-143.

^d In patients intended to receive HDT/ASCR, bendamustine should be used with caution as there are conflicting data regarding ability to collect peripheral progenitor cell collection.

^e There is a randomized trial that demonstrated that RCHOP was not superior to CHOP.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Mantle Cell Lymphoma

SUGGESTED TREATMENT REGIMENS^{a,b}

An FDA-approved biosimilar is an appropriate substitute for rituximab.

Second-line Therapy

- Partial response^f with intention to proceed to transplant
 - ▶ Preferred regimens (in alphabetical order)
 - ◊ Bendamustine^d ± rituximab (if not previously given)
 - ◊ Bortezomib ± rituximab
 - ◊ Lenalidomide ± rituximab
 - ◊ RCHOP or VRCAP (if not previously given)
- Short response duration to prior chemoimmunotherapy (< expected median PFS)
 - ▶ Preferred regimens (in alphabetical order)
 - ◊ BTK inhibitors^g
 - Acalabrutinib^h
 - Ibrutinib ± rituximab
 - Zanubrutinib
 - ◊ Lenalidomide ± rituximab
 - ◊ Venetoclax^g
 - ▶ Other recommended regimens
 - ◊ Ibrutinib,^g lenalidomide, rituximab (category 2B)
 - ◊ Venetoclax + ibrutinib^g (category 2B)

- Extended response duration to prior chemoimmunotherapy (> expected median PFS)
 - ▶ Preferred regimens (in alphabetical order)
 - ◊ Bendamustine ± rituximab (if not previously given)
 - ◊ Bortezomib ± rituximab
 - ◊ BTK inhibitors^g
 - Acalabrutinib^h
 - Ibrutinib ± rituximab
 - Zanubrutinib
 - ◊ Lenalidomide ± rituximab
 - ▶ Other recommended regimens (in alphabetical order by category)
 - ◊ Venetoclax^g
 - ◊ Bendamustine, bortezomib, and rituximab (category 2B)
 - ◊ PEPC (prednisone, etoposide, procarbazine, cyclophosphamide) ± rituximab (category 2B)
 - ◊ RCHOP or VRCAP (if not previously given) (category 2B)
 - ◊ [See Second-line Therapy for DLBCL \(BCEL-C 2 of 4\)](#) without regard to transplantability

Second-line Consolidation

- Allogeneic hematopoietic cell transplant (nonmyeloablative or myeloablative)

Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

^a See references for regimens [MANT-A 3 of 4](#) and [MANT-A 4 of 4](#).

^b Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^d In patients intended to receive HDT/ASCR, bendamustine should be used with caution as there are conflicting data regarding ability to collect peripheral progenitor cell collection.

^f Patients who have achieved near CR can proceed to HDT/ASCR. Patients who have achieved minimal PR with substantial disease should be treated as having stable, refractory disease. Patients who have achieved a very good PR may be treated with additional therapy to achieve CR with the goal of proceeding to HDT/ASCR.

^g [See Special Considerations for Use of Small-Molecule Inhibitors \(NHODG-E\)](#).

^h The phase 2 ACE-LY-004 study excluded patients treated with Bruton's tyrosine kinase (BTK) or BCL-2 inhibitor and concomitant warfarin or equivalent vitamin K antagonists.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**SUGGESTED TREATMENT REGIMENS**
REFERENCES**Induction Therapy****Aggressive therapy****HyperCVAD (cyclophosphamide, vincristine, doxorubicin, dexamethasone alternating with methotrexate and cytarabine) + rituximab**

Romaguera JE, Fayad L, Rodriguez MA, et al. High rate of durable remissions after treatment of newly diagnosed aggressive mantle-cell lymphoma with rituximab plus hyper-CVAD alternating with rituximab plus high-dose methotrexate and cytarabine. *J Clin Oncol* 2005;23:7013-7023.

Merli F, Luminari S, Ilariucci F, et al. Rituximab plus HyperCVAD alternating with high dose cytarabine and methotrexate for the initial treatment of patients with mantle cell lymphoma, a multicentre trial from Gruppo Italiano Studio Linfomi. *Br J Haematol* 2012;156:346-353.

Nordic trial regimen (Dose-intensified induction immunochemotherapy with rituximab + cyclophosphamide, vincristine, doxorubicin, prednisone [maxi-CHOP] alternating with rituximab + high-dose cytarabine)

Geisler CH, Kolstad A, Laurell A, et al. Long-term progression-free survival of mantle cell lymphoma following intensive front-line immunochemotherapy with in vivo-purged stem cell rescue: A non-randomized phase-II multicenter study by the Nordic Lymphoma Group. *Blood* 2008;112:2687-2693.

Eskelund CW, Kolstad A, Jerkeman M, et al. 15-year follow-up of the Second Nordic Mantle Cell Lymphoma trial (MCL2): prolonged remissions without survival plateau. *Br J Haematol* 2016;175:410-418.

RCHOP/RDHAP

Hermine O, Hoster E, Walewski J, et al. Addition of high-dose cytarabine to immunochemotherapy before autologous stem-cell transplantation in patients aged 65 years or younger with mantle cell lymphoma (MCL Younger): a randomised, open-label, phase 3 trial of the European Mantle Cell Lymphoma Network. *Lancet* 2016;388:565-575.

Delarue R, Haioun C, Ribrag V, et al. CHOP and DHAP plus rituximab followed by autologous stem cell transplantation (ASCT) in mantle cell lymphoma (MCL): a phase II study from the GELA. *Blood* 2013;121:48-53.

RDHAP (rituximab, dexamethasone, cisplatin [oxaliplatin or carboplatin], cytarabine)

Le Gouill S, Thieblemont C, Oberic L, et al. Rituximab after autologous stem-cell transplantation in mantle-cell lymphoma. *N Engl J Med* 2017;377:1250-1260.

Le Gouill S, Thieblemont C, Oberic L, et al. R-DHA-Oxaliplatin before autologous stem cell transplantation prolongs PFS and OS as compared to R-DHA-carboplatin and R-DHA-cisplatin in patients with mantle cell lymphoma, a subgroup analysis of the LyMa trial [abstract]. *Blood* 2017;130 (Suppl 1):Abstract 1496.

Less aggressive therapy**Bendamustine + rituximab**

Rummel MJ, Niederle N, Maschmeyer G, et al. Bendamustine plus rituximab versus CHOP plus rituximab as first-line treatment for patients with indolent and mantle-cell lymphomas: an open-label, multicentre, randomised, phase 3 non-inferiority trial. *Lancet* 2013;381:1203-1210.

Flinn IW, van der Jagt R, Kahl BS, et al. Open-label, randomized, noninferiority study of bendamustine-rituximab or R-CHOP/R-CVP in first-line treatment of advanced indolent NHL or MCL: the BRIGHT study. *Blood* 2014;123:2944-2952.

Bendamustine + rituximab + maintenance rituximab

Rummel MJ, Knauf W, Goerner M, et al. Two years rituximab maintenance vs. observation after first-line treatment with bendamustine plus rituximab (B-R) in patients with mantle cell lymphoma: First results of a prospective, randomized, multicenter phase II study (a subgroup study of the StiL NHL7-2008 MAINTAIN trial) [abstract]. *J Clin Oncol* 2016;34:Abstract 7503.

VR-CAP (bortezomib, rituximab, cyclophosphamide, doxorubicin, and prednisone)

Robak T, Jin J, Pylypenko H, et al. Frontline bortezomib, rituximab, cyclophosphamide, doxorubicin, and prednisone (VR-CAP) versus rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisone (R-CHOP) in transplantation-ineligible patients with newly diagnosed mantle cell lymphoma: final overall survival results of a randomised, open-label, phase 3 study. *Lancet Oncol* 2018;19:1449-1458.

CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) + rituximab

Lenz G, Dreyling M, Hoster E, et al. Immunochemotherapy with rituximab and cyclophosphamide, doxorubicin, vincristine, and prednisone significantly improves response and time to treatment failure, but not long-term outcome in patients with previously untreated mantle cell lymphoma: results of a prospective randomized trial of the German Low Grade Lymphoma Study Group (GLSG). *J Clin Oncol* 2005;23:1984-1992.

Kluin-Nelemans HC, Hoster E, Hermine O, et al. Treatment of older patients with mantle-cell lymphoma. *N Engl J Med* 2012;367:520-531.

Modified HyperCVAD with rituximab maintenance

Kahl BS, Long WL, Eickhoff JC, et al. Maintenance rituximab following induction chemoimmunotherapy may prolong progression-free survival in mantle cell lymphoma: A pilot study from the Wisconsin Oncology Network. *Ann Oncol* 2006;17:1418-1423.

Lenalidomide + rituximab

Ruan J, Martin P, Shah B, et al. Lenalidomide plus rituximab as initial treatment for mantle-cell lymphoma. *N Engl J Med* 2015;373:1835-44.

Ruan J, Martin P, Christos P, et al. Five-year follow-up of lenalidomide plus rituximab as initial treatment for mantle cell lymphoma. *Blood* 2018;132:2016-2025.

RBAC500 (rituximab, bendamustine, cytarabine)

Visco C, Finotto S, Zambello R, et al. Combination of rituximab, bendamustine, and cytarabine for patients with mantle-cell non-Hodgkin lymphoma ineligible for intensive regimens or autologous transplantation. *J Clin Oncol* 2013;31:1442-1449.

Visco C, Chiappella A, Nassi L, et al. Rituximab, bendamustine, and low-dose cytarabine as induction therapy in elderly patients with mantle cell lymphoma: a multicentre, phase 2 trial from Fondazione Italiana Linfomi. *Lancet Haematol* 2017;4:e15-e23.

Note: All recommendations are category 2A unless otherwise indicated.**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.****Continued****MANT-A**
3 OF 4

**SUGGESTED TREATMENT REGIMENS**
REFERENCES**First-line Consolidation****High-dose therapy with autologous stem cell rescue**

Dreyling M, Lenz G, Hoster E, et al. Early consolidation by myeloablative radiochemotherapy followed by autologous stem cell transplantation in first remission significantly prolongs progression-free survival in mantle cell lymphoma: results of a prospective randomized trial of the European MCL Network. *Blood* 2005;105:2677-2684.

Thieblemont C, Antal D, Lacotte-Thierry L, et al. Chemotherapy with rituximab followed by high-dose therapy and autologous stem cell transplantation in patients with mantle cell lymphoma. *Cancer* 2005;104:1434-1441.

Ritchie D, Seymour J, Grigg A, et al. The hyper-CVAD—rituximab chemotherapy programme followed by high-dose busulfan, melphalan and autologous stem cell transplantation produces excellent event-free survival in patients with previously untreated mantle cell lymphoma. *Ann Hematol* 2007;86:101-105.

van 't Veer MB, de Jong D, MacKenzie M, et al. High-dose Ara-C and beam with autograft rescue in R-CHOP responsive mantle cell lymphoma patients. *Br J Haematol* 2009;144:524-530.

Rituximab maintenance

Graf S, Stevenson P, Holmberg LA, et al. Maintenance rituximab after autologous stem cell transplantation in patients with mantle cell lymphoma. *Ann Oncol* 2015;26:2323-2328.

Le Gouill S, Thieblemont C, Oberic L, et al. Rituximab after autologous stem-cell transplantation in mantle-cell lymphoma. *N Engl J Med* 2017;377:1250-1260.

Kluin-Nelemans HC, Hoster E, Hermine O, et al. Treatment of Older Patients With Mantle Cell Lymphoma (MCL): Long-Term Follow-Up of the Randomized European MCL Elderly Trial. *J Clin Oncol* 2020;38:248-256.

Second-line Therapy**Acalabrutinib**

Wang M, Rule S, Zinzani PL, et al. Acalabrutinib in relapsed or refractory mantle cell lymphoma (ACE-LY-004): a single-arm, multicentre, phase 2 trial. *Lancet* 2018;391:659-667.

Bendamustine

Robinson KS, Williams ME, van der Jagt RH, et al. Phase II multicenter study of bendamustine plus rituximab in patients with relapsed indolent B-cell and mantle cell Non-Hodgkin's Lymphoma. *J Clin Oncol* 2008; 26:4473-4479.

Rummel MJ, Al-Batran SE, Kim S-Z, et al. Bendamustine plus rituximab is effective and has a favorable toxicity profile in the treatment of mantle cell and low-grade non-hodgkin's lymphoma. *J Clin Oncol* 2005;23:3383-3389.

Bendamustine, bortezomib, and rituximab

Friedberg JW, Vose JM, Kelly JL, et al. The combination of bendamustine, bortezomib, and rituximab for patients with relapsed/refractory indolent and mantle cell non-Hodgkin lymphoma. *Blood* 2011;117:2807-2812.

Bortezomib ± rituximab

Goy A, Bernstein SH, Kahl BS, et al. Bortezomib in patients with relapsed or refractory mantle cell lymphoma: updated time-to-event analyses of the multicenter phase 2 PINNACLE study. *Ann Oncol* 2009;20:520-525.

Baiocchi RA, Alinari L, Lustberg ME, et al. Phase 2 trial of rituximab and bortezomib in patients with relapsed or refractory mantle cell and follicular lymphoma. *Cancer* 2011;117:2442-2451.

Lamm W, Kaufmann H, Raederer M, et al. Bortezomib combined with rituximab and dexamethasone is an active regimen for patients with relapsed and chemotherapy-refractory mantle cell lymphoma. *Haematologica* 2011;96:1008-1014.

Ibrutinib ± rituximab

Wang ML, Blum KA, Martin P, et al. Long-term follow-up of MCL patients treated with single-agent ibrutinib: updated safety and efficacy results. *Blood* 2015;126:739-745.

Dreyling M, Jurczak W, Jerkeman M, et al. Ibrutinib versus temsirolimus in patients with relapsed or refractory mantle-cell lymphoma: an international, randomised, open-label, phase 3 study. *Lancet* 2016;387:770-778.

Rule S, Jurczak W, Jerkeman M, et al. Ibrutinib versus temsirolimus: 3-year follow-up of patients with previously treated mantle cell lymphoma from the phase 3, international, randomized, open-label RAY study. *Leukemia* 2018;32:1799-1803.

Wang ML, Lee H, Chuang H, et al. Ibrutinib in combination with rituximab in relapsed or refractory mantle cell lymphoma: a single-centre, open-label, phase 2 trial. *Lancet Oncol* 2016;7:48-56.

Ibrutinib, lenalidomide, and rituximab

Jerkeman M, Eskelund C, Hutchings M, et al. Ibrutinib, lenalidomide, and rituximab in relapsed or refractory mantle cell lymphoma (PHILEMON): a multicentre, open-label, single-arm, phase 2 trial. *Lancet Haematol* 2018(3):e109-e116.

Lenalidomide ± rituximab

Habermann TM, Lossos IS, Justice G, et al. Lenalidomide oral monotherapy produces a high response rate in patients with relapsed or refractory mantle cell lymphoma. *Br J Haematol* 2009;145:344-349.

Goy A, Sinha R, Williams ME, et al. Single-agent lenalidomide in patients with mantle-cell lymphoma who relapsed or progressed after or were refractory to bortezomib: phase II MCL-001 (EMERGE) study. *J Clin Oncol* 2013;31:3688-3695.

Zinzani PL, Vose JM, Czuczman MS, et al. Long-term follow-up of lenalidomide in relapsed/refractory mantle cell lymphoma: subset analysis of the NHL-003 study. *Ann Oncol* 2013;24:2892-2897.

Trneny M, Lamy T, Walewski J, et al. Lenalidomide versus investigator's choice in relapsed or refractory mantle cell lymphoma (MCL-002; SPRINT): a phase 2, randomised, multicentre trial. *Lancet Oncol* 2016;17:319-331.

Wang M, Fayad L, Wagner-Bartak N, et al. Lenalidomide in combination with rituximab for patients with relapsed or refractory mantle-cell lymphoma: a phase 1/2 clinical trial. *Lancet Oncol* 2012;13:716-723.

PEP-C (prednisone, etoposide, procarbazine, cyclophosphamide) ± rituximab

Coleman M, Martin P, Ruan J, et al. Low-dose metronomic, multidrug therapy with the PEP-C oral combination chemotherapy regimen for mantle cell lymphoma. *Leuk Lymphoma* 2008;49:447-450.

Venetoclax

David M, Roberts A, Seymour J, et al. Phase I first-in-human study of venetoclax in patients with relapsed or refractory non-Hodgkin lymphoma. *J Clin Oncol* 2017;35:826-833.

David M, von Keudell G, Portell G, et al. Revised dose ramp-up to mitigate the risk of tumor lysis syndrome when initiating venetoclax in patients with mantle cell lymphoma. *J Clin Oncol* 2018;36:3525-3527.

Venetoclax + ibrutinib

Tam C, Anderson M, Pott C, et al. Ibrutinib plus venetoclax for the treatment of mantle-cell lymphoma. *N Engl J Med* 2018;378:1211-1223.

Zanubrutinib

Song Y, Zhou K, Zou D, et al. Zanubrutinib in patients with relapsed/refractory mantle cell lymphoma [abstract]. *Hematol Oncol* 2019;37:45-46.

Tam CS, Wang M, Simpson D, et al. Updated safety and efficacy data in the phase 1 trial of patients with mantle cell lymphoma (mcl) treated with bruton tyrosine kinase (BTK) inhibitor zanubrutinib (BGB-3111) [abstract]. *Hematol Oncol* 2019;37:245-247.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

ADDITIONAL DIAGNOSTIC TESTING^a

ESSENTIAL:

- Adequate immunophenotyping to establish diagnosis and GCB versus non-GCB origin^{b,c}
 - ▶ IHC panel: CD20, CD3, CD5, CD10, CD45, BCL2, BCL6, Ki-67, IRF4/MUM1, MYC with or without
 - ▶ Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: kappa/lambda, CD45, CD3, CD5, CD19, CD10, CD20
- Karyotype or FISH for *MYC*

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- Additional immunohistochemical studies to establish lymphoma subtype
 - ▶ IHC panel: cyclin D1, kappa/lambda, CD30, CD138, Epstein-Barr virus in situ hybridization (EBV-ISH), ALK, HHV8, SOX11
- Karyotype or FISH for *BCL2*, *BCL6* rearrangements if *MYC* positive

SUBTYPES

• Subtypes included:

- ▶ DLBCL, NOS^{d,e}
- ▶ DLBCL coexistent with follicular lymphoma of any grade
- ▶ DLBCL coexistent with gastric MALT lymphoma
- ▶ DLBCL coexistent with nongastric MALT lymphoma
- ▶ Follicular lymphoma grade 3^f
- ▶ Intravascular large B-cell lymphoma
- ▶ DLBCL associated with chronic inflammation
- ▶ ALK-positive DLBCL^g
- ▶ EBV-positive DLBCL, NOS
- ▶ T-cell-/histiocyte-rich large B-cell lymphoma
- ▶ DLBCL with *IRF4/MUM1* rearrangement

• Subtypes *not* included:

- ▶ Primary cutaneous marginal zone lymphoma (PCMZL) and primary cutaneous follicle center lymphoma (PCFCL) ([See NCCN Guidelines for Primary Cutaneous B-Cell Lymphomas](#))
- ▶ Primary DLBCL of the CNS ([See NCCN Guidelines for CNS](#))

- Primary mediastinal large B-cell lymphoma (PMBL); [see BCEL-B 1 of 3](#).
- Grey zone lymphoma; [see BCEL-B 2 of 3](#).
- High-grade B-cell lymphomas with translocations of *MYC* and *BCL2* and/or *BCL6* (double-/triple-hit lymphoma); [see HGBL-1](#).
- High-grade B-cell lymphomas, NOS; [see HGNOS-1](#).
- Primary cutaneous DLBCLs, leg type; [see BCEL-B 3 of 3](#).
- DLBCL arising from CLL (Richter's transformation) ([See NCCN Guidelines for CLL/SLL](#))

^e In the 2017 revised WHO classification of lymphomas, DLBCL, double hit has been designated in a unique category called high-grade B-cell lymphomas with translocations of *MYC* and *BCL2* and/or *BCL6*.

^f FL, grade 3b is commonly treated as DLBCL ([BCEL-1](#)). The management of FL, grade 3a is controversial and treatment should be individualized.

^g These are most often CD20 negative and rituximab is not necessary.

→ [See
Workup
\(BCEL-2\)](#)

^a [See International Prognostic Index \(BCEL-A\)](#).

^b Typical immunophenotype: CD20+, CD45+, CD3-; other markers used for subclassification.

^c [See Use of Immunophenotyping/Genetic Testing in Differential Diagnosis of Mature B-Cell and NK/T-Cell Neoplasms \(NHODG-A\)](#).

^d Germinal center (or follicle center) phenotype is not equivalent to follicular lymphoma and can occur in DLBCL and Burkitt lymphoma. Morphology is required to establish diagnosis.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

WORKUP

ESSENTIAL:

- Physical exam: attention to node-bearing areas, including Waldeyer's ring, and to size of liver and spleen
- Performance status
- B symptoms
- CBC with differential
- LDH
- Comprehensive metabolic panel
- Uric acid
- Whole-body PET/CT scan ± C/A/P CT with contrast of diagnostic quality
- Adequate bone marrow biopsy (>1.6 cm) ± aspirate; bone marrow biopsy is not necessary if PET/CT scan demonstrates bone disease. Bone marrow biopsy with a negative PET/CT scan may reveal discordant lymphoma
- Calculation of International Prognostic Index (IPI) (See [BCEL-A 1 of 2](#))
- Hepatitis B testing^h
- Echocardiogram or MUGA scan if anthracycline or anthracenedione-based regimen is indicated
- Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)

→ [See Induction Therapy \(BCEL-3\)](#)

USEFUL IN SELECTED CASES:

- Head CT/MRI with contrast or neck CT/MRI with contrast
- Discussion of fertility issues and sperm banking
- HIV testing
- Hepatitis C testing
- Beta-2-microglobulin
- Lumbar puncture for patients at risk for CNS involvement, see [BCEL-A 2 of 2](#)

^h Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen ([See NHODG-B](#)). If positive, check viral load and consider consult with gastroenterologist.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma



ⁱ In testicular lymphoma, after completion of chemotherapy, scrotal RT should be given (25–30 Gy).

^j In patients who are not candidates for chemotherapy, ISRT is recommended.

^k [See Prognostic Model to Assess the Risk of CNS Disease \(BCEL-A 2 of 2\)](#).

^l Patients with systemic disease with concurrent CNS disease, [see BCEL-C](#).

^m Recommendations are for HIV-negative lymphoma only.

For HIV-positive DLBCL, [see AIDS-2](#).

ⁿ [See BCEL-C](#) for regimens used in patients with poor left ventricular function, very frail patients, and patients >80 years of age with comorbidities.

^o [See Principles of Radiation Therapy \(NHODG-D\)](#).

^p If RT is not used, interim staging after 3–4 cycles of RCHOP is appropriate to confirm response.

^q Based on current clinical trials, RCHOP is preferable due to reduced toxicities, but other comparable anthracycline-based regimens are also acceptable ([see BCEL-C](#)).

^r In selected cases, RT to initially bulky sites of disease may be beneficial (category 2B).

^s PET/CT scan at interim restaging can lead to increased false positives and should be carefully considered in select cases. If PET/CT scan performed and positive, rebiopsy before changing course of treatment.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

PRE RT EVALUATION

(End of first-line chemoimmunotherapy)

FOLLOW-UP THERAPY



^o See [Principles of Radiation Therapy \(NHODG-D\)](#).

^t Repeat biopsy should be strongly considered if PET-positive prior to additional therapy. If biopsy negative, follow PET-negative guideline.

^v See [Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#).

^v PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS) ([See NHODG-C 3 of 3](#)).

^w Chemosensitive partial response.

^x The optimum timing of end-of-treatment PET/CT is unknown; however, waiting a minimum of 8 weeks after RT to repeat PET/CT scan is suggested. False positives may occur due to posttreatment changes.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

END-OF-TREATMENT RESTAGING

END-OF-TREATMENT RESPONSE

FOLLOW-UP

Stage I, II:
At completion of treatment, repeat all positive studies^{t,x}

Complete response^{u,y}
(PET negative [5-PS 1–3])^v

Clinical
• H&P and labs, every 3–6 mo for 5 y and then yearly or as clinically indicated
Imaging^z
• Repeat C/A/P CT with contrast scan only as clinically indicated

→ Relapse →

[See Relapse or Refractory Disease \(BCEL-7\)](#)

Partial response^{t,u}
(PET positive [5-PS 4])^v

[See Relapse or Refractory Disease \(BCEL-7\)](#)
or
ISRT^o

Progressive disease^{t,u}
(PET positive [5-PS 5])^v

[See Relapse or Refractory Disease \(BCEL-7\)](#)

^o See Principles of Radiation Therapy (NHODG-D).

^t Repeat biopsy should be strongly considered if PET-positive prior to additional therapy. If biopsy negative, follow PET-negative guideline.

^u See Lugano Response Criteria for Non-Hodgkin Lymphoma (NHODG-C).

^v PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS) (See NHODG-C 3 of 3).

^x The optimum timing of end-of-treatment PET/CT is unknown; however, waiting a minimum of 8 weeks after RT to repeat PET/CT scan is suggested. False positives may occur due to posttreatment changes.

^y Patients in first remission may be candidates for consolidation trials including high-dose therapy with autologous stem cell rescue.

^z When disease (eg, bone) is only imaged on PET/CT scan, it is appropriate to be used in follow-up.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma


INTERIM RESTAGING

FOLLOW-UP THERAPY

END-OF- TREATMENT RESTAGING

END-OF-TREATMENT RESPONSE

FOLLOW-UP



^o See [Principles of Radiation Therapy \(NHODG-D\)](#).

^s PET/CT scan at interim restaging can lead to increased false positives and should be carefully considered in select cases. If PET/CT scan performed and positive, rebiopsy before changing course of treatment.

^t Repeat biopsy should be strongly considered in PET-positive prior to additional therapy. If biopsy negative, follow PET-negative guideline.

^u See [Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#).

^v PET/CT scan should be interpreted via the PET Five-Point Scale (5-PS) ([See NHODG-C 3 of 3](#)).

^z When disease (eg, bone) is only imaged on PET/CT scan, it is appropriate to be used in follow-up.

^{aa} A 5-PS = 4 response can reflect post-treatment inflammation as well as active disease. If there is uncertainty regarding interpretation of the response, consider brief interval restaging to clarify post-treatment inflammation response.

^{bb} For other regimens, [see BCEL-C](#).


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma



^o See Principles of Radiation Therapy (NHODG-D).

^u See Lugano Response Criteria for Non-Hodgkin Lymphoma (NHODG-C).

^{cc} Some NCCN Member Institutions require a complete metabolic response in order to proceed to high-dose therapy with autologous stem cell rescue.

^{dd} Additional RT can be given before or after transplant to sites of previous positive disease.

^{ee} Selected cases include mobilization failures and persistent bone marrow involvement.

^{ff} See Guidance for Treatment of Patients with Chimeric Antigen Receptor (CAR) T-Cell Therapy (BCEL-D).

^{gg} Tisagenlecleucel is not FDA-approved for relapsed/refractory primary mediastinal large B-cell lymphoma.

^{hh} Repeat biopsy should be strongly considered if PET-positive prior to additional therapy, because PET positivity may represent post-treatment inflammation. If biopsy negative, follow CR pathway.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

FOLLOW-UP

RELAPSE #2 OR GREATER



Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))

See monoclonal antibody and viral reactivation ([NHODG-B](#))

^o See Principles of Radiation Therapy (NHODG-D).

^u See Lugano Response Criteria for Non-Hodgkin Lymphoma (NHODG-C).

^{ff} See Guidance for Treatment of Patients with Chimeric Antigen Receptor (CAR) T-Cell Therapy (BCEL-D).

^{gg} Tisagenlecleucel is not FDA-approved for relapsed/refractory primary mediastinal large B-cell lymphoma.

^{hh} Repeat biopsy should be strongly considered if PET-positive prior to additional therapy, because PET positivity may represent post-treatment inflammation. If biopsy negative, follow CR pathway.

ⁱⁱ Patients who progress after three successive regimens are unlikely to derive additional benefit from currently utilized combination chemotherapy regimens, except for patients with a long disease-free interval.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

INTERNATIONAL PROGNOSTIC INDEX^a

ALL PATIENTS:

- Age >60 years
- Serum LDH > normal
- Performance status 2–4
- Stage III or IV
- Extranodal involvement >1 site

INTERNATIONAL INDEX, ALL PATIENTS:

- Low 0 or 1
- Low-intermediate 2
- High-intermediate 3
- High 4 or 5

AGE-ADJUSTED INTERNATIONAL PROGNOSTIC INDEX^a

PATIENTS ≤60 YEARS:

- Stage III or IV
- Serum LDH > normal
- Performance status 2–4

**INTERNATIONAL INDEX,
PATIENTS ≤60 YEARS:**

- Low 0
- Low-intermediate 1
- High-intermediate 2
- High 3

STAGE-MODIFIED INTERNATIONAL PROGNOSTIC INDEX^b

STAGE I OR II PATIENTS:

- Age >60 years
- Serum LDH > normal
- Performance status 2–4
- Stage II or IIE

**INTERNATIONAL INDEX,
STAGE I OR II PATIENTS:**

- Low 0 or 1
- High 2–4

NCCN-IPI^c

Risk Group**Age, years**

- >40 to ≤60 1
- >60 to <75 2
- ≥75 3

LDH, normalized

- >1 to ≤3 1
- >3 2

Ann Arbor stage III-IV 1**Extranodal disease* 1****Performance status ≥2 1**

- Low 0–1
- Low-intermediate 2–3
- High-intermediate 4–5
- High ≥6

*Disease in bone marrow, CNS, liver/GI tract, or lung.

^a The International Non-Hodgkin's Lymphoma Prognostic Factors Project. A predictive model for aggressive non-hodgkin's lymphoma. N Engl J Med 1993;329:987-994.^b Miller TP, Dahlberg S, Cassady JR. Chemotherapy alone compared with chemotherapy plus radiotherapy for localized intermediate- and high-grade non-Hodgkin's lymphoma. N Engl J Med 1998;339:21-26.^c This research was originally published in *Blood*. Zhou Z, Sehn LH, Rademaker AW, et al. An enhanced International Prognostic Index (NCCN-IPI) for patients with diffuse large B-cell lymphoma treated in the rituximab era. Blood 2014;123:837-842. © the American Society of Hematology**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**[Back to Workup](#)
[\(BCEL-2\)](#)**BCEL-A**
1 OF 2



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

Prognostic Model to Assess the Risk of CNS Disease^d

- | | | |
|----------------------------------------------------------------------------------------------------------------------|-------------------|--------------------------------------------|
| • Age >60 years | Low risk | 0–1 |
| • Serum LDH > normal | Intermediate-risk | 2–3 |
| • Performance status >1 | High-risk | 4–6 or kidney or adrenal gland involvement |
| • Stage III or IV | | |
| • Extranodal involvement >1 site | | |
| • Kidney or adrenal gland involvement | | |
| • Additional indications for CNS prophylaxis independent of CNS risk score | | |
| ▶ HIV lymphoma | | |
| ▶ Testicular lymphoma | | |
| ▶ High-grade B-cell lymphomas (HGBLs) with translocations of <i>MYC</i> and <i>BCL2</i> and/or <i>BCL6</i> HGBL, NOS | | |
| ▶ Primary cutaneous DLBCL, leg type | | |
| ▶ Stage IE DLBCL of the breast | | |
| • Suggested CNS prophylactic therapy (Optimal management is uncertain) | | |
| ▶ Intrathecal methotrexate and/or cytarabine (4–8 doses) during the course of treatment | | |
| or | | |
| ▶ Systemic methotrexate (3–3.5 g/m ² for 2–4 cycles) during the course of treatment | | |

^d Schmitz N, Zeynalova S, Nickelsen M, et al. CNS International Prognostic Index: A risk model for CNS relapse in patients with diffuse large B-cell lymphoma treated with R-CHOP. J Clin Oncol 2016;34:3150-3156.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Back to Workup](#)
[\(BCEL-2\)](#)

BCEL-A
2 OF 2



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

PRIMARY MEDIASTINAL LARGE B-CELL LYMPHOMA

- **Primary mediastinal large B-cell lymphoma (PMBL) can be defined as a clinical entity presenting with primary site of disease in mediastinum with or without other sites and has histology of DLBCL. PMBL overlaps with grey zone lymphomas that have intermediate features between Hodgkin lymphoma and PMBL and have unique diagnostic characteristics.**
[See Grey Zone Lymphoma \(BCEL-B 2 of 3\).](#)
- **Clinical pathologic correlation is required to establish diagnosis.**
- **Optimal first-line therapy is more controversial than other subtypes of NHL; however, treatment regimens include (in order of preference):^{a,b}**
 - ▶ **Dose-adjusted EPOCH-R ([etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin] + rituximab)^c x 6 cycles**
 - ◊ **For persistent focal disease, RT can be added.**
 - ▶ **RCHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone) x 6 cycles + RT**
 - ▶ **RCHOP x 4 cycles followed by ICE (ifosfamide, carboplatin, etoposide) ± rituximab^d x 3 cycles ± RT (category 2B)**
- **Residual mediastinal masses are common. PET/CT scan is essential post-treatment. Biopsy of PET/CT scan positive mass is recommended if additional systemic treatment is contemplated.**
- **Relapsed/refractory therapy**
 - ▶ [See BCEL-7](#)
 - ▶ **Pembrolizumab^e**
 - ▶ **Nivolumab ± brentuximab vedotin (category 2B)^f**

^a Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^b An FDA-approved biosimilar is an appropriate substitute for rituximab.

^c Dunleavy K, Pittaluga S, Maeda LS, et al. Dose-adjusted EPOCH-rituximab therapy in primary mediastinal B-cell lymphoma. N Engl J Med 2013;368:1408-1416.

^d Moskowitz C, Hamlin PA, Jr., Maragulia J, et al. Sequential dose-dense RCHOP followed by ICE consolidation (MSKCC protocol 01-142) without radiotherapy for patients with primary mediastinal large B-cell lymphoma [abstract]. Blood 2010;116:Abstract 420.

^e Armand P, Rodig S, Melnichenko V, et al. Pembrolizumab in Relapsed or Refractory Primary Mediastinal Large B-Cell Lymphoma. J Clin Oncol 2019;37:3291-3299.

^f Zinzani P, Santoro A, Gritti G, et al. Nivolumab combined with brentuximab vedotin for relapsed/refractory primary mediastinal large b-cell lymphoma: efficacy and safety from the phase II Checkmate 436 study. J Clin Oncol 2019;37:3081-3089.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

GREY ZONE LYMPHOMA^a (intermediate between DLBCL and classical HL)

Synonyms

- B-cell lymphoma, unclassifiable, with features intermediate between DLBCL and classical Hodgkin lymphoma (CHL)
- Large B-cell lymphoma with Hodgkin features
- Hodgkin-like anaplastic large cell lymphoma

Clinical Presentation

- Present with large anterior mediastinal mass with or without supraclavicular lymph nodes
 - ▶ More common in males, presenting between 20–40 y
- Non-mediastinal grey zone lymphoma is more likely compared to mediastinal cases to occur in older individuals and typically have higher risk features, more advanced-stage disease, and higher IPI.^b

Morphology

- Pleomorphic cells in a diffusely fibrous stroma
- Typically larger and more pleomorphic than in PMBL, sometimes resembling lacunar or Hodgkin-like cells
- Necrosis without neutrophilic infiltrate is frequent

Immunophenotype

- Often transitional features between CHL and PMBL
- CD45 often positive; CD30, CD15, CD20, CD79a frequently positive
- EBV-
- PAX5, BOB.1, OCT-2 are often positive, BCL6 is variable
- CD10, ALK are negative
- If morphology closer to PMBL, or absence of CD20, CD15+ would suggest the diagnosis of grey zone lymphoma
- If morphology closer to CHL, CD20 strong positivity and other B-cell markers and absence of CD15 would suggest grey zone lymphoma.

Prognosis and Treatment^c

- A worse prognosis than either CHL or PMBL has been suggested.
- While there is no consensus on the treatment, aggressive large B-cell lymphoma regimens are preferred.
- If the tumor cells are CD20+, the addition of rituximab to chemotherapy should be considered.^d
- Data suggest that the use of rituximab and anthracycline-based chemotherapy as in other B-cell lymphomas ([See BCEL-C](#)) is helpful. If localized disease, then consolidative RT is preferred.
- There is no ostensible difference in outcome between mediastinal and non-mediastinal grey zone lymphoma.

^a Pilichowska M, Pittaluga S, Ferry JA, et al. Clinicopathologic consensus study of gray zone lymphoma with features intermediate between DLBCL and classical HL. *Blood Adv* 2017;1:2600-2609; Wilson WH, Pittaluga S, Nicolae A, et al. A prospective study of mediastinal gray-zone lymphoma. *Blood* 2014;124:1563-1569; Swerdlow SH CE, Harris NL, Jaffe ES, Pileri SA, Stein H, Thiele J, ed. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Revised 4th ed. Lyon: IARC; 2017.; Quintanilla-Martinez L, de Jong D, de Mascarel A, et al. Gray zones around diffuse large B cell lymphoma. Conclusions based on the workshop of the XIV meeting of the European Association for Hematopathology and the Society of Hematopathology in Bordeaux, France. *J Hematop* 2009;2:211-236.

^b Evens AM, Kanakry JA, Sehn LH, et al. Gray zone lymphoma with features intermediate between classical Hodgkin lymphoma and diffuse large B-cell lymphoma: Characteristics, outcomes, and prognostication among a large multicenter cohort. *Am J Hematol* 2015;90:778-783.

^c Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^d An FDA-approved biosimilar is an appropriate substitute for rituximab.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

PRIMARY CUTANEOUS DIFFUSE LARGE B-CELL LYMPHOMA, LEG TYPE^a



^a Expert hematopathologist review is essential to confirm the diagnosis of primary cutaneous DLBCL, Leg type.

^b [For TNM Classification of Cutaneous Lymphoma other than MF/SS \(See NCCN Guidelines for Primary Cutaneous Lymphomas\).](#)

^c Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^d These patients are at higher risk for CNS involvement ([See \[BCEL-A 2 of 2\]\(#\)](#)); consider CNS prophylaxis according to institutional standards.

^e For patients who cannot tolerate anthracyclines, see [BCEL-C](#) for regimens for patients with poor left ventricular function.

^f [See Principles of Radiation Therapy \(NHODG-D\).](#)

^g For patients not able to tolerate chemotherapy.

^h PET/CT (strongly preferred) or C/A/P CT with contrast at the end of treatment to assess response. It can be repeated if there is clinical suspicion of progressive disease.

ⁱ An FDA-approved biosimilar is an appropriate substitute for rituximab.

^j Selection of patients requires adequate marrow cellularity >15% and <25% involvement of lymphoma in bone marrow, and platelets >100,000. In patients with prior autologous stem cell rescue, referral to a tertiary care center is highly recommended for ibritumomab tiuxetan. If ibritumomab tiuxetan is considered, bilateral cores are recommended and the pathologist should provide the percent of overall cellular elements and the percent of cellular elements involved in the marrow. As of 2010, updates suggest a trend towards an increased risk of MDS with RIT. Cytogenetics/FISH assessment for MDS markers is recommended for patients receiving RIT.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

SUGGESTED TREATMENT REGIMENS^{a,b}

An FDA-approved biosimilar is an appropriate substitute for rituximab.

First-line Therapy

- Preferred regimen
 - ▶ RCHOP (rituximab,^c cyclophosphamide, doxorubicin, vincristine, prednisone) (category 1)
- Other recommended regimens
 - ▶ Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin) + rituximab
 - ▶ Dose-dense RCHOP 14^c (category 3)

First-line Therapy for Patients with Poor Left Ventricular Function^{d,e,f}

- Other recommended regimens (in alphabetical order)
 - ▶ DA-EPOCH^g (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin) + rituximab
 - ▶ RCDOP (rituximab, cyclophosphamide, liposomal doxorubicin, vincristine, prednisone)
 - ▶ RCEPP (rituximab, cyclophosphamide, etoposide, prednisone, procarbazine)
 - ▶ RCEOP (rituximab, cyclophosphamide, etoposide, vincristine, prednisone)
 - ▶ RGCVP (rituximab, gemcitabine, cyclophosphamide, vincristine, prednisolone)

First-line Therapy for Very Frail Patients and Patients >80 Years of Age with Comorbidities^{e,f}

- Other recommended regimens (in alphabetical order)
 - ▶ RCEPP (rituximab, cyclophosphamide, etoposide, prednisone, procarbazine)
 - ▶ RCDOP (rituximab, cyclophosphamide, liposomal doxorubicin, vincristine, prednisone)
 - ▶ R-mini-CHOP
 - ▶ RGCVP (rituximab, gemcitabine, cyclophosphamide, vincristine, prednisolone)

First-line Consolidation (optional)

- Lenalidomide maintenance (category 2B) for patients 60–80 y of age

Concurrent Presentation with CNS Disease^h

- Parenchymal: systemic methotrexate (≥ 3 g/m² or more of given on Day 15 of a 21-day RCHOP cycle that has been supported by growth factors)
- Leptomeningeal: IT methotrexate/cytarabine, consider Ommaya reservoir placement. Systemic methotrexate (3–3.5 g/m²) can be given in combination with RCHOP or as consolidation after RCHOP + IT methotrexate/cytarabine

See Second-line Therapy on [BCEL-C 2 of 4](#).

Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))

See monoclonal antibody and viral reactivation ([NHODG-B](#))

^a See references for regimens [BCEL-C 3 of 4](#) and [BCEL-C 4 of 4](#).

^b Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^c In RCHOP-14 and RCHOP-21, may consider increasing dose of rituximab to 500 mg/m² in men >60 y.

^d Inclusion of any anthracycline or anthracenedione in patients with impaired cardiac functioning should have more frequent cardiac monitoring.

^e There are limited published data regarding the use of these regimens; however, they are used at NCCN Member Institutions for the first-line treatment of DLBCL for patients with poor left ventricular function, very frail patients, and patients >80 years of age with comorbidities.

^f There are limited data for treatment of early-stage disease with these regimens; however, short-course chemotherapy + RT for stage I–II disease is practiced at NCCN Member Institutions.

^g If upward dose adjustment is necessary, doxorubicin should be maintained at base dose and not increased.

^h Concurrent high-dose methotrexate with dose-adjusted EPOCH can result in unacceptable toxicities.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

SUGGESTED TREATMENT REGIMENS^{a,b}

An FDA-approved biosimilar is an appropriate substitute for rituximab.

Second-line and Subsequent Therapy^{d,i,j} (intention to proceed to transplant)

- Preferred regimens (in alphabetical order)
 - ▶ DHAP (dexamethasone, cisplatin, cytarabine) ± rituximab
 - ▶ DHAX (dexamethasone, cytarabine, oxaliplatin) ± rituximab
 - ▶ GDP (gemcitabine, dexamethasone, cisplatin) ± rituximab or (gemcitabine, dexamethasone, carboplatin) ± rituximab
 - ▶ ICE (ifosfamide, carboplatin, etoposide) ± rituximab
- Other recommended regimens (in alphabetical order)
 - ▶ ESHAP (etoposide, methylprednisolone, cytarabine, cisplatin) ± rituximab
 - ▶ GemOx (gemcitabine, oxaliplatin) ± rituximab
 - ▶ MINE (mesna, ifosfamide, mitoxantrone, etoposide) ± rituximab

Anti-CD19 CAR T-cell therapy^{n,o}

- Axicabtagene ciloleucel
- Tisagenlecleucel

See First-line Therapy on [BCEL-C 1 of 4](#).

Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

^a See references for regimens [BCEL-C 3 of 4](#) and [BCEL-C 4 of 4](#).

^b Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^d Inclusion of any anthracycline or anthracenedione in patients with impaired cardiac functioning should have more frequent cardiac monitoring.

ⁱ If additional anthracycline is administered after a full course of therapy, careful cardiac monitoring is essential. Dexrazoxane may be added as a cardioprotectant.

^j Rituximab should be included in second-line therapy if there is relapse after a reasonable remission (>6 mo); however, rituximab should often be omitted in patients with primary refractory disease.

^k In patients intended to receive CAR T-cell therapy, bendamustine should be used with caution unless immediately prior to CAR T-cell therapy, since it could impact the success of the patient's T-cell collection.

Second-line and Subsequent Therapy^{d,i,j} (non-candidates for transplant)

- Preferred regimens (in alphabetical order)
 - ▶ GemOx ± rituximab
 - ▶ Polatuzumab vedotin ± bendamustine ± rituximab (after ≥2 prior therapies)^{k,l}
- Other recommended regimens (in alphabetical order)
 - ▶ CEPP (cyclophosphamide, etoposide, prednisone, procarbazine) ± rituximab - PO and IV
 - ▶ CEOP (cyclophosphamide, etoposide, vincristine, prednisone) ± rituximab
 - ▶ DA-EPOCH ± rituximab
 - ▶ GDP ± rituximab or (gemcitabine, dexamethasone, carboplatin) ± rituximab
 - ▶ Gemcitabine, vinorelbine ± rituximab (category 3)
 - ▶ Rituximab
- Useful in certain circumstances
 - ▶ Brentuximab vedotin for CD30+ disease
 - ▶ Bendamustine^k ± rituximab (category 2B)
 - ▶ Ibrutinib^m (non-GCB DLBCL)
 - ▶ Lenalidomide ± rituximab (non-GCB DLBCL)

Third-line and Subsequent Therapy (including patients with disease progression after transplant or CAR T-cell therapy)

- Selinexor (only after at least two lines of systemic therapy)^p

^l Bendamustine, rituximab, and polatuzumab vedotin-piiq is indicated for the treatment of adult patients with relapsed or refractory DLBCL and HGBL with translocations of *MYC* and *BCL2* and/or *BCL6* after ≥2 prior therapies.

^m [See Special Considerations for Use of Small-Molecule Inhibitors \(NHODG-E\)](#).

ⁿ [See Guidance for Treatment of Patients with Chimeric Antigen Receptor \(CAR\) T-Cell Therapy \(BCEL-D\)](#).

^o Tisagenlecleucel is not FDA-approved for relapsed/refractory primary mediastinal large B-cell lymphoma.

^p Selinexor is FDA approved only for DLBCL and transformed DLBCL arising from FL.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

SUGGESTED TREATMENT REGIMENS

REFERENCES

First-line Therapy

CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) + rituximab with RT

Miller TP, Dahlberg S, Cassady JR, et al. Chemotherapy alone compared with chemotherapy plus radiotherapy for localized intermediate- and high-grade non-hodgkin's lymphoma. *N Engl J Med* 1998;339:21-26.

Horning SJ, Weller E, Kim K, et al. Chemotherapy with or without radiotherapy in limited-stage diffuse aggressive non-hodgkin's lymphoma: Eastern Cooperative Oncology Group Study 1484. *J Clin Oncol* 2004;22:3032-3038.

Persky DO, Unger JM, Spier CM, et al. Phase II study of rituximab plus three cycles of CHOP and involved-field radiotherapy for patients with limited-stage aggressive B-cell lymphoma: Southwest Oncology Group Study 0014. *J Clin Oncol* 2008;26:2258-2263.

CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) + rituximab

Coiffier B, Thieblemont C, Van Den Neste E, et al. Long-term outcome of patients in the LNH-98.5 trial, the first randomized study comparing rituximab-CHOP to standard CHOP chemotherapy in DLBCL patients: a study by the Groupe d'Etudes des Lymphomes de l'Adulte. *Blood* 2010;116:2040-2045.

Feugier P, Van Hoof A, Sebban C, et al. Long-term results of the R-CHOP study in the treatment of elderly patients with diffuse large B-cell lymphoma: a study by the Groupe d'Etude des Lymphomes de l'Adulte. *J Clin Oncol* 2005;23:4117-4126.

Pfreundschuh M, Trumper L, Osterborg A, et al. CHOP-like chemotherapy plus rituximab versus CHOP-like chemotherapy alone in young patients with good-prognosis diffuse large-B-cell lymphoma: a randomised controlled trial by the MabThera International Trial (MInT) Group. *Lancet Oncol* 2006;7:379-391.

Poeschel V, Held G, Ziepert M, et al. Excellent outcome of young patients (18-60 years) with favorable-Prognosis diffuse large B-cell lymphoma (DLBCL) treated with 4 cycles CHOP plus 6 applications of rituximab: Results of the 592 patients of the Flyer trial of the DSHNHL/GLA. *Blood* 2018;132: Abstract 781

Dose-dense CHOP 14 + rituximab

Pfreundschuh M, Schubert J, Ziepert M, et al. Six versus eight cycles of bi-weekly CHOP-14 with or without rituximab in elderly patients with aggressive CD20+ B-cell lymphomas: a randomised controlled trial (RICOVER-60). *Lancet Oncol* 2008;9:105-116.

Cunningham D, Hawkes EA, Jack A, et al. Rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisolone in patients with newly diagnosed diffuse large B-cell non-Hodgkin lymphoma: a phase 3 comparison of dose intensification with 14-day versus 21-day cycles. *Lancet* 2013;381:1817-1826.

Lamy T, Damaj G, Soubeyran P, et al. R-CHOP 14 with or without radiotherapy in nonbulky limited-stage diffuse large B-cell lymphoma. *Blood* 2018;131:174-181.

Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin) + rituximab

Purroy N, Bergua J, Gallur L, et al. Long-term follow-up of dose-adjusted EPOCH plus rituximab (DA-EPOCH-R) in untreated patients with poor prognosis large B-cell lymphoma. A phase II study conducted by the Spanish PETHEMA Group. *Br J Haematol* 2015;169:188-198.

Wilson WH, Dunleavy K, Pittaluga S, et al. Phase II study of dose-adjusted EPOCH and rituximab in untreated diffuse large B-cell lymphoma with analysis of germinal center and post-germinal center biomarkers. *J Clin Oncol* 2008;26:2717-2724.

Wilson WH, Jung SH, Porcu P, et al. A Cancer and Leukemia Group B multi-center study of DA-EPOCH-rituximab in untreated diffuse large B-cell lymphoma with analysis of outcome by molecular subtype. *Haematologica* 2012;97:758-765.

First-line Therapy for Patients with Poor Left Ventricular Function

CDOP (cyclophosphamide, liposomal doxorubicin, vincristine, and prednisone) + rituximab

Martino R, Perea G, Caballero MD, et al. Cyclophosphamide, pegylated liposomal doxorubicin (Caelyx), vincristine and prednisone (CCOP) in elderly patients with diffuse large B-cell lymphoma: Results from a prospective phase II study. *Haematologica* 2002;87:822-827.

Zaja F, Tomadini V, Zaccaria A, et al. CHOP-rituximab with pegylated liposomal doxorubicin for the treatment of elderly patients with diffuse large B-cell lymphoma. *Leuk Lymphoma* 2006;47:2174-2180.

RCEOP (rituximab, cyclophosphamide, etoposide, vincristine, prednisone)

Moccia A, Schaff K, Hoskins P, et al. R-CHOP with etoposide substituted for doxorubicin (R-CEOP): Excellent outcome in diffuse large B cell lymphoma for patients with a contraindication to anthracyclines [abstract]. *Blood* 2009;114:Abstract 408.

RGCVP (rituximab, gemcitabine, cyclophosphamide, vincristine, and prednisolone)

Fields PA, Townsend W, Webb A, et al. De novo treatment of diffuse large B-cell lymphoma with rituximab, cyclophosphamide, vincristine, gemcitabine, and prednisolone in patients with cardiac comorbidity: a United Kingdom National Cancer Research Institute trial. *J Clin Oncol* 2014;32:282-287.

First-line Therapy for Elderly Patients (age >80 years)

R-mini-CHOP

Peyrade F, Jardin F, Thieblemont C, et al. Attenuated immunochemotherapy regimen (R-miniCHOP) in elderly patients older than 80 years with diffuse large B-cell lymphoma: a multicentre, single-arm, phase 2 trial. *Lancet Oncol* 2011;12:460-468.

Peyrade F, Fain O, Fabiani B, et al. Long-term follow-up of the GELA LNH 03-7B study: A prospective phase II study of 150 patients over 80 years with diffuse large B-cell lymphoma (DLBCL) treated with RminiCHOP [abstract]. *J Clin Oncol* 2013;31(15_suppl):Abstract 8536.

First-line Consolidation

Thieblemont C, Tilly H, Gomes da Silva M, et al. Lenalidomide maintenance compared with placebo in responding elderly patients with diffuse large B-cell lymphoma treated with first-line rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisone. *J Clin Oncol* 2017;35:2473-2481.

Second-line and Subsequent Therapy

Bendamustine ± rituximab

Weidmann E, Kim SZ, Rost A, et al. Bendamustine is effective in relapsed or refractory aggressive non-Hodgkin's lymphoma. *Ann Oncol* 2002;13:1285-1289.

Vacirca JL, Acs PI, Tabbara IA, et al. Bendamustine combined with rituximab for patients with relapsed or refractory diffuse large B cell lymphoma. *Ann Hematol* 2014;93:403-409.

Ohmachi K, Niitsu N, Uchida T, et al. Multicenter phase II study of bendamustine plus rituximab in patients with relapsed or refractory diffuse large B-cell lymphoma. *J Clin Oncol* 2013;31:2103-2109.

Bendamustine, rituximab and polatuzumab vedotin-piiq

Sehn LH, Herrera AF, Matasar MJ, et al. Polatuzumab vedotin (Pola) plus bendamustine (B) with rituximab (R) or obinutuzumab (G) in relapsed/refractory (R/R) Diffuse Large B-Cell Lymphoma (DLBCL): Updated results of a phase (Ph) Ib/II study [abstract]. *Blood* 2018;132: Abstract 1683.

Brentuximab vedotin

Jacobsen ED, Sharman JP, Oki Y, et al. Brentuximab vedotin demonstrates objective responses in a phase 2 study of relapsed/refractory DLBCL with variable CD30 expression. *Blood* 2015;125:1394-1402.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

BCEL-C
3 OF 4



NCCN Guidelines Version 2.2020

Diffuse Large B-Cell Lymphoma

SUGGESTED TREATMENT REGIMENS

REFERENCES

Second-line and Subsequent Therapy (continued)

CAR T-Cell Therapy

Axicabtagene ciloleucel

Neelapu SS, Locke FL, Bartlett NL, et al. Axicabtagene ciloleucel CAR T-cell therapy in refractory large B-cell lymphoma. *N Engl J Med* 2017;377:2531-2544.

Locke FL, Ghobadi A, Jacobson CA, et al. Long-term safety and activity of axicabtagene ciloleucel in refractory large B-cell lymphoma (ZUMA-1): a single-arm, multicentre, phase 1-2 trial. *Lancet Oncol* 2019;20:31-42.

Tisagenlecleucel

Schuster SJ, Bishop MR, Tam CS, et al. Tisagenlecleucel in adult relapsed or refractory diffuse large B-cell lymphoma. *N Engl J Med* 2019;380:45-56.

CEPP (cyclophosphamide, etoposide, prednisone, procarbazine) ± rituximab

Chao NJ, Rosenberg SA, and Horning SJ. CEPP(B): An effective and well-tolerated regimen in poor-risk, aggressive non-Hodgkin's lymphoma. *Blood* 1990;76:1293-1298.

DHAP (dexamethasone, cisplatin, cytarabine) ± rituximab

Mey UJ, Orloff KS, Flieger D, et al. Dexamethasone, high-dose cytarabine, and cisplatin in combination with rituximab as salvage treatment for patients with relapsed or refractory aggressive non-Hodgkin's lymphoma. *Cancer Invest* 2006;24:593-600.

Gisselbrecht C, Schmitz N, Mounier N, et al. Rituximab maintenance therapy after autologous stem-cell transplantation in patients with relapsed CD20+ diffuse large B-cell lymphoma: Final analysis of the collaborative trial in relapsed aggressive lymphoma. *J Clin Oncol* 2012;30:4462-4469.

DHAX (dexamethasone, cytarabine, oxaliplatin) ± rituximab

Lignon J, Sibon D, Madelaine I, et al. Rituximab, dexamethasone, cytarabine, and oxaliplatin (R-DHAX) is an effective and safe salvage regimen in relapsed/refractory B-cell non-Hodgkin lymphoma. *Clin Lymphoma Myeloma Leuk* 2010;10:262-269.

Rigacci L, Fabbri A, Puccini B, et al. Oxaliplatin-based chemotherapy (dexamethasone, high-dose cytarabine, and oxaliplatin) +/-rituximab is an effective salvage regimen in patients with relapsed or refractory lymphoma. *Cancer* 2010;116:4573-4579.

EPOCH + rituximab

Gutierrez M, Chabner BA, Pearson D, et al. Role of a doxorubicin-containing regimen in relapsed and resistant lymphomas: An 8-year follow-up study of EPOCH. *J Clin Oncol* 2000;18:3633-3642.

Jermann M, Jost LM, Taverna C, et al. Rituximab-EPOCH, an effective salvage therapy for relapsed, refractory or transformed B-cell lymphomas: Results of a phase II study. *Ann Oncol* 2004;15:511-516.

ESHAP (etoposide, methylprednisolone, cytarabine, cisplatin) ± rituximab

Velasquez WS, McLaughlin P, Tucker S, et al. ESHAP - an effective chemotherapy regimen in refractory and relapsing lymphoma: a 4-year follow-up study. *J Clin Oncol* 1994;12:1169-1176.

Martin A, Conde E, Arnan M, et al. R-ESHAP as salvage therapy for patients with relapsed or refractory diffuse large B-cell lymphoma: the influence of prior exposure to rituximab on outcome. *A GEL/TAMO study. Haematologica* 2008;93:1829-1836.

GDP (gemcitabine, dexamethasone, cisplatin or carboplatin) ± rituximab

Crump M, Kuruvilla J, Couban S, et al. Randomized comparison of gemcitabine, dexamethasone, and cisplatin versus dexamethasone, cytarabine, and cisplatin chemotherapy before autologous stem-cell transplantation for relapsed and refractory aggressive lymphomas: NCIC-CTG LY.12. *J Clin Oncol* 2014;32:3490-3496.

Gopal AK, Press OW, Shustov AR, et al. Efficacy and safety of gemcitabine, carboplatin, dexamethasone, and rituximab in patients with relapsed/refractory lymphoma: a prospective multi-center phase II study by the Puget Sound Oncology Consortium. *Leuk Lymphoma* 2010;51:1523-1529.

GemOX (gemcitabine, oxaliplatin) ± rituximab

Lopez A, Gutierrez A, Palacios A, et al. GEMOX-R regimen is a highly effective salvage regimen in patients with refractory/relapsing diffuse large-cell lymphoma: a phase II study. *Eur J Haematol* 2008;80:127-132.

Corazzelli G, Capobianco G, Arcamone M, et al. Long-term results of gemcitabine plus oxaliplatin with and without rituximab as salvage treatment for transplant-ineligible patients with refractory/relapsing B-cell lymphoma. *Cancer Chemother Pharmacol* 2009;64:907-916.

Mounier N, El Gnaoui T, Tilly H, et al. Rituximab plus gemcitabine and oxaliplatin in patients with refractory/relapsed diffuse large B-cell lymphoma who are not candidates for high-dose therapy. A phase II Lymphoma Study Association trial. *Haematologica* 2013;98:1726-1731.

Gemcitabine, vinorelbine, rituximab

Papageorgiou ES, Tsirigotis P, Dimopoulos M, et al. Combination chemotherapy with gemcitabine and vinorelbine in the treatment of relapsed or refractory diffuse large B-cell lymphoma: a phase-II trial by the Hellenic Cooperative Oncology Group. *Eur J Hematol* 2005;75:124-129.

Xiros N, Economopoulos T, Valsami S, et al. Rituximab in combination with vinorelbine/gemcitabine chemotherapy in patients with primary refractory or early relapsed T cell rich B cell lymphoma. A pilot study. *Leuk Res* 2003;27:1097-1099.

Ibrutinib

Wilson WH, Young RM, Schmitz R, et al. Targeting B cell receptor signaling with ibrutinib in diffuse large B cell lymphoma. *Nat Med* 2015;21:922-926.

ICE (ifosfamide, carboplatin, etoposide) ± rituximab

Kewalramani T, Zelenetz AD, Nimer SD, et al. Rituximab and ICE (RICE) as second-line therapy prior to autologous stem cell transplantation for relapsed or primary refractory diffuse large B-cell lymphoma. *Blood* 2004;103:3684-8.

Gisselbrecht C, Schmitz N, Mounier N, et al. Rituximab maintenance therapy after autologous stem-cell transplantation in patients with relapsed CD20+ diffuse large B-cell lymphoma: Final analysis of the collaborative trial in relapsed aggressive lymphoma. *J Clin Oncol* 2012;30:4462-4469.

Lenalidomide ± rituximab

Wang M, Fowler N, Wagner-Bartak N, et al. Oral lenalidomide with rituximab in relapsed or refractory diffuse large cell, follicular, and transformed lymphoma: a phase II clinical trial. *Leukemia* 2013;27:1902-1909.

Czuczman MS, Trneny M, Davies A, et al. A phase 2/3 multicenter, randomized, open-label study to compare the efficacy and safety of lenalidomide versus investigator's choice in patients with relapsed or refractory diffuse large B-cell lymphoma. *Clin Cancer Res* 2017;23:4127-4137.

Third-line and Subsequent Therapy

Neste, Andre Goy, Brian T. Hill, Catherine Thieblemont, et al. Single Agent Oral Selinexor Demonstrates Deep and Durable Responses in Relapsed/Refractory Diffuse Large B-Cell Lymphoma (DLBCL) in Both GCB and Non-GCB Subtypes: The Phase 2b Sadal Study. *Blood* 2018; 132 (Supplement 1): Abstract 1677.

Federica Cavallo, George A Follows, Andre Goy, et al. Effect of Prior Therapy on the Efficacy and Safety of Oral Selinexor in Patients with Relapsed/Refractory (R/R) Diffuse Large B-Cell Lymphoma (DLBCL): A Post-Hoc Analysis of the Sadal Study. *Blood* 2019; 134 (Supplement_1): Abstract 5333.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



GUIDANCE FOR TREATMENT OF PATIENTS WITH CHIMERIC ANTIGEN RECEPTOR (CAR) T-CELL THERAPY

Axicabtagene ciloleucel^a

• Patient selection

- ▶ Axicabtagene ciloleucel is indicated for the treatment of adult patients with relapsed or refractory large B-cell lymphoma after two or more lines of systemic therapy, including DLBCL, NOS; primary mediastinal large B-cell lymphoma; high-grade B-cell lymphoma; and DLBCL arising from follicular lymphoma.^a
- ▶ Health care facilities that dispense and administer axicabtagene ciloleucel must be enrolled and comply with the Risk Evaluation and Mitigation Strategies (REMS) requirements.^a [See REMS for axicabtagene ciloleucel.](#)

• Cytokine release syndrome (CRS) management - See CAR T-Cell-Related Toxicities in the [NCCN Guidelines for Management of Immunotherapy-Related Toxicities](#)

• Neurologic toxicity management - See CAR T-Cell-Related Toxicities in the [NCCN Guidelines for Management of Immunotherapy-Related Toxicities](#)

• Prolonged cytopenias

- ▶ Patients may exhibit cytopenias for several weeks following lymphodepleting chemotherapy and axicabtagene ciloleucel infusion.

• Hypogammaglobulinemia

- ▶ B-cell aplasia and hypogammaglobulinemia can occur in patients receiving treatment with axicabtagene ciloleucel.

Tisagenlecleucel^b

• Patient selection

- ▶ Tisagenlecleucel is indicated for the treatment of adult patients with relapsed or refractory large B-cell lymphoma after two or more lines of systemic therapy including DLBCL, NOS and high grade B-cell lymphoma and DLBCL arising from follicular lymphoma.^b
- ▶ Health care facilities that dispense and administer tisagenlecleucel must be enrolled and comply with the REMS requirements.^b [See REMS for tisagenlecleucel.](#)

• CRS management - See CAR T-Cell-Related Toxicities in the [NCCN Guidelines for Management of Immunotherapy-Related Toxicities](#)

• Neurologic toxicity management - See CAR T-Cell-Related Toxicities in the [NCCN Guidelines for Management of Immunotherapy-Related Toxicities](#)

• Prolonged cytopenias

- ▶ Patients may exhibit cytopenias for several weeks following lymphodepleting chemotherapy and tisagenlecleucel infusion.

• Hypogammaglobulinemia

- ▶ B-cell aplasia and hypogammaglobulinemia can occur in patients with a complete remission after tisagenlecleucel infusion.

^a Prescribing information for axicabtagene ciloleucel is available at: <https://www.fda.gov/media/108377/download>

^b Prescribing information for tisagenlecleucel is available at: <https://www.fda.gov/media/107296/download>

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

High-Grade B-Cell Lymphomas with Translocations of MYC and BCL2 and/or BCL6 (Double-/Triple-Hit Lymphoma)

Definition^a

- High-grade B-cell lymphomas (HGBLs) with translocations of *MYC* and *BCL2* and/or *BCL6* as detected by FISH or standard cytogenetics are known as "double-hit" lymphomas. If all three are rearranged, they are referred to as "triple-hit" lymphomas.
- The vast majority are germinal center B-cell–like lymphomas.

Clinical Presentation

- Often present with poor prognostic parameters, such as elevated LDH, bone marrow and CNS involvement, and a high IPI score.

Treatment^{b,c}

- Clinical trial is recommended.
- While the standard of care is not established, the following induction regimens have been used at NCCN Member Institutions:
 - DA-EPOCH-R
 - Potentially toxic regimens; performance status and comorbidities should be considered
 - ◊ RHyper-CVAD (rituximab, cyclophosphamide, vincristine, doxorubicin, and dexamethasone alternating with high-dose methotrexate and cytarabine)
 - ◊ R-CODOX-M/R-IVAC (rituximab-cyclophosphamide, vincristine, doxorubicin with methotrexate/ifosfamide, etoposide, and cytarabine)
- RCHOP has been associated with inferior outcomes.
- Consider consolidation with high-dose therapy with autologous stem cell rescue. While its role is not established, this is done at some NCCN Member Institutions.
- These patients are at higher risk for CNS involvement ([See BCEL-A 2 of 2](#))
- If localized disease, then consolidative RT is preferred.
- Relapsed/refractory disease, [see BCEL-7](#).

References:

- Petrich A, Gandhi M, Jovanovic B, et al. Impact of induction regimen and stem cell transplantation on outcomes in double-hit lymphoma: a multicenter retrospective analysis. *Blood* 2014;124:2354-2361.
- Dunleavy K, Fanale MA, Abramson JS, et al. Dose-adjusted EPOCH-R (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin, and rituximab) in untreated aggressive diffuse large B-cell lymphoma with MYC rearrangement: a prospective, multicentre, single-arm phase 2 study. *Lancet Haematol* 2018;5:e609-e617.
- Johnson NA, Slack GW, Savage KJ et al. Concurrent expression of MYC and BCL2 in diffuse large B-cell lymphoma treated with rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisone. *J Clin Oncol* 2012;30:3452-3459.
- Green TM, Young KH, Visco C, et al. Immunohistochemical double-hit score is a strong predictor of outcome in patients with diffuse large B-cell lymphoma treated with rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisone. *J Clin Oncol* 2012;30:3460-3467.

^a In the 2017 revised WHO classification of lymphomas, DLBCL, double hit has been designated in a unique category called HGBL with translocations of *MYC* and *BCL2* and/or *BCL6*.

^b Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^c An FDA-approved biosimilar is an appropriate substitute for rituximab.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

High-Grade B-Cell Lymphomas, NOS

Definition

- HGBL, NOS include cases that appear blastoid or cases intermediate between DLBCL and BL, but which lack an *MYC* and *BCL2* and/or *BCL6* rearrangement.
- They exclude patient with translocations of *MYC* and *BCL2* and/or *BCL6* or clear DLBCL.

Clinical Presentation

- HGBCLs, NOS often present with poor prognostic parameters, such as elevated LDH, bone marrow and CNS involvement, and a high IPI score.

Treatment^{a,b}

- Clinical trial is recommended.
- For early-stage disease, consider consolidative RT for localized disease.
- While the standard of care is not established, the following induction regimens have been used at NCCN Member Institutions:
 - ▶ RCHOP
 - ▶ DA-EPOCH-R
 - ▶ Potentially toxic regimens; performance status and comorbidities should be considered
 - ◊ RHyper-CVAD (rituximab, cyclophosphamide, vincristine, doxorubicin, and dexamethasone alternating with high-dose methotrexate and cytarabine)
 - ◊ R-CODOX-M/R-IVAC (rituximab, cyclophosphamide, vincristine, doxorubicin with methotrexate/ifosfamide, etoposide, and cytarabine)
- Consider consolidation with high-dose therapy with autologous stem cell rescue. While its role is not established, this is done at some NCCN Member Institutions.
- These patients are at higher risk for CNS involvement ([See BCEL-A 2 of 2](#))
- For Relapse/Refractory Disease, [see BCEL-7](#).

^a Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^b An FDA-approved biosimilar is an appropriate substitute for rituximab.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**ADDITIONAL DIAGNOSTIC TESTING^{a,b}****ESSENTIAL:**

- Adequate immunophenotyping to establish diagnosis^{c,d,e}
 - ▶ IHC panel: CD45, CD20, CD3, CD10, Ki-67, BCL2, BCL6, TdT with or without
 - ▶ Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: kappa/lambda, CD45, CD20, CD3, CD5, CD19, CD10, TdT
- Karyotype ± FISH: t(8;14) or variants; *MYC* rearrangement^f

USEFUL UNDER CERTAIN CIRCUMSTANCES

- FISH: *BCL2*; *BCL6* rearrangements
- EBER-ISH

WORKUP**ESSENTIAL:**

- Physical exam: attention to node-bearing areas, including Waldeyer's ring, and to size of liver and spleen
 - Performance status
 - B symptoms
 - CBC with differential
 - LDH
 - Comprehensive metabolic panel
 - Uric acid
 - C/A/P CT with contrast of diagnostic quality
 - Lumbar puncture
 - Flow cytometry of cerebrospinal fluid
 - Unilateral or bilateral bone marrow biopsy ± aspirate
 - HIV testing (if positive, [see AIDS-1](#))
 - Hepatitis B testing^g
 - Echocardiogram or MUGA scan if anthracycline or anthracenedione-based regimen is indicated
 - Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)
- USEFUL IN SELECTED CASES:**
- Hepatitis C testing
 - Neck CT with contrast
 - Discussion of fertility issues and sperm banking
 - Brain MRI with and without contrast
 - Whole-body PET/CT scan^h

[See Risk Assessment and Induction Therapy \(BURK-2\)](#)

^a For treatment of double- or triple-hit tumors, [see HGBL-1](#). In other cases where it is not possible to distinguish between BL and high-grade lymphoma, therapy per this guideline may be appropriate.

^b This disease is complex and curable; it is preferred that treatment occur at centers with expertise in the management of the disease.

^c Typical immunophenotype: sIg+, CD10+, Cd20+, TdT-, Ki-67+ (≥95%), BCL2-, BCL6+.

^d [See Use of Immunophenotyping/Genetic Testing in Differential Diagnosis of Mature B-Cell and NK/T-Cell Neoplasms \(NHODG-A\)](#).

^e If flow cytometry initially performed, IHC for selected markers (BCL2 and Ki-67) can supplement the flow results.

^f Most common karyotype is *MYC* rearrangement as a sole abnormality. There is an uncommon variant of BL without *MYC* rearrangement but with 11q aberration. Optimum management of this rare subtype is undefined, though it is most often treated like typical BL.

^g Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen ([See NHODG-B](#)). If positive, check viral load and consider consult with gastroenterologist.

^h Initiation of therapy should not be delayed in order to obtain a PET/CT scan.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2020

Burkitt Lymphoma

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

RISK ASSESSMENT

INDUCTION THERAPY

INITIAL RESPONSE

RELAPSE

Low risk
Normal LDH
or
Stage I and
Completely resected
abdominal lesion
or
Single extra-
abdominal mass
<10 cm

Clinical trial
or
See Suggested
Regimensⁱ ([BURK-A](#))

Complete
response^j

Follow-up after complete
response:^k
every 2–3 mo for 1 y,
then every 3 mo for 1 y,
then every 6 mo^l

→ Relapse → [See \(BURK-3\)](#)

< Complete
response^j

Clinical trial^m
or
Palliative ISRTⁿ

Prophylaxis for tumor lysis syndrome
is mandatory ([See NHODG-B](#))

See monoclonal antibody and viral
reactivation ([NHODG-B](#))

High risk
Stage I
and
Abdominal mass
or
Extra-abdominal mass
>10 cm
or
Stage II–IV

Clinical trial
or
See Suggested
Regimensⁱ ([BURK-A](#))

Complete
response^j

Follow-up after complete
response:^k
every 2–3 mo for 1 y,
then every 3 mo for 1 y,
then every 6 mo^l

→ Relapse → [See \(BURK-3\)](#)

< Complete
response^j

Clinical trial^m
or
Palliative ISRTⁿ

ⁱ All regimens for Burkitt lymphoma include CNS prophylaxis/therapy.

^j [See Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#).

^k Repeat C/A/P CT scan with contrast only as clinically indicated.

^l Relapse after 2 y is rare; therefore, follow-up should be individualized according to patient characteristics.

^m If there is no clinical trial available, some NCCN Member Institutions use non-cross resistant second-line therapy regimens listed on [BURK-A](#) in selected cases.

ⁿ [See Principles of Radiation Therapy \(NHODG-D\)](#).


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Burkitt Lymphoma



Prophylaxis for tumor lysis syndrome is mandatory ([See NHODG-B](#))

See monoclonal antibody and viral reactivation ([NHODG-B](#))

ⁱ All regimens for Burkitt lymphoma include CNS prophylaxis/therapy.

^j [See Lugano Response Criteria for Non-Hodgkin Lymphoma \(NHODG-C\)](#).

^m If there is no clinical trial available, some NCCN Member Institutions use non-cross resistant second-line therapy regimens listed on [BURK-A](#) in selected cases.

ⁿ [See Principles of Radiation Therapy \(NHODG-D\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



SUGGESTED TREATMENT REGIMENS^{a,b}

An FDA-approved biosimilar is an appropriate substitute for rituximab.

CHOP is not an adequate therapy.

Induction Therapy <60 y

- **Low Risk, Preferred Regimens (alphabetical order)**
 - ▶ **CODOX-M (original or modified) (cyclophosphamide, doxorubicin, vincristine with intrathecal methotrexate and cytarabine followed by high-dose systemic methotrexate) + rituximab (3 cycles)**
 - ▶ **Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin) + rituximab (minimum 3 cycles with one additional cycle beyond CR) (regimen includes intrathecal methotrexate) (Data included patients with leptomeningeal CNS disease; patients with parenchymal CNS disease were excluded in the clinical trials of this regimen.)**
 - ▶ **HyperCVAD (cyclophosphamide, vincristine, doxorubicin, and dexamethasone) alternating with high-dose methotrexate and cytarabine + rituximab (regimen includes intrathecal therapy)**
- **High Risk, Preferred Regimens (alphabetical order)**
 - ▶ **High-risk patients presenting with symptomatic CNS disease should be started with the portion of the systemic therapy that contains CNS penetrating drugs.**
 - ▶ **CODOX-M (original or modified) (cyclophosphamide, doxorubicin, vincristine with intrathecal methotrexate and cytarabine followed by high-dose systemic methotrexate) alternating with IVAC (ifosfamide, cytarabine, etoposide, and intrathecal methotrexate) + rituximab**
 - ▶ **Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin) + rituximab (for high-risk patients not able to tolerate aggressive treatments) (regimen includes intrathecal methotrexate) (Data included patients with leptomeningeal CNS disease; patients with parenchymal CNS disease were excluded in the clinical trials of this regimen.)**
 - ▶ **HyperCVAD (cyclophosphamide, vincristine, doxorubicin, and dexamethasone) alternating with high-dose methotrexate and cytarabine + rituximab (regimen includes intrathecal therapy)**

Induction Therapy ≥60 y

- **Low Risk and High Risk, Preferred Regimen**
 - ▶ **Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin) + rituximab (minimum 3 cycles with one additional cycle beyond CR) (regimen includes intrathecal methotrexate) (Data included patients with leptomeningeal CNS disease; patients with parenchymal CNS disease were excluded in the clinical trials of this regimen.)**
 - ◊ **In high-risk patients presenting with symptomatic CNS disease, the management of the CNS disease should be addressed with the initial regimen.**

^a See references for regimens [BURK-A 3 of 3](#).

^b All regimens for Burkitt lymphoma include CNS prophylaxis/therapy.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



SUGGESTED TREATMENT REGIMENS^{a,b} (in alphabetical order)

An FDA-approved biosimilar is an appropriate substitute for rituximab.

Second-line Therapy

- Patients with disease relapse >6–18 mo after appropriate first-line therapy should be treated with alternate regimens. While no definitive second-line therapies exist, there are limited data for the following regimens:
- Other recommended regimens
 - ▶ Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin) + rituximab (minimum 3 cycles with one additional cycle beyond CR) (regimen includes intrathecal methotrexate) (Data included patients with leptomeningeal CNS disease; patients with parenchymal CNS disease were excluded in the clinical trials of this regimen.)
 - ▶ RICE (rituximab, ifosfamide, carboplatin, etoposide); intrathecal methotrexate if have not received previously
 - ▶ RIVAC (rituximab, ifosfamide, cytarabine, etoposide); intrathecal methotrexate if have not received previously
- Useful in certain circumstances
 - ▶ RGDP (rituximab, gemcitabine, dexamethasone, cisplatin)
 - ▶ High-dose cytarabine + rituximab

Prophylaxis for tumor lysis syndrome is mandatory ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

^a See references for regimens [BURK-A 3 of 3](#).

^b All regimens for Burkitt lymphoma include CNS prophylaxis/therapy.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



SUGGESTED TREATMENT REGIMENS REFERENCES

Induction Therapy

Low- and High-Risk Combination Regimens

CODOX-M (original or modified) (cyclophosphamide, doxorubicin, vincristine with intrathecal methotrexate and cytarabine followed by high-dose systemic methotrexate) with (for high-risk) or without (for low-risk) alternating IVAC (ifosfamide, cytarabine, etoposide, and intrathecal methotrexate ± rituximab)

LaCasce A, Howard O, Lib S, et al. Modified magrath regimens for adults with Burkitt and Burkitt-like lymphoma: preserved efficacy with decreased toxicity. *Leuk Lymphoma* 2004;45:761-767.

Mead GM, Sydes MR, Walewski J, et al. An international evaluation of CODOX-M and CODOX-M alternating with IVAC in adult Burkitt's lymphoma: results of United Kingdom Lymphoma Group LY06 study. *Ann Oncol* 2002;13:1264-1274.

Barnes JA, Lacasce AS, Feng Y, et al. Evaluation of the addition of rituximab to CODOX-M/IVAC for Burkitt's lymphoma: a retrospective analysis. *Ann Oncol* 2011;22:1859-1864.

Evens AM, Carson KR, Kolesar J, et al. A multicenter phase II study incorporating high-dose rituximab and liposomal doxorubicin into the CODOX-M/IVAC regimen for untreated Burkitt's lymphoma. *Ann Oncol* 2013;24:3076-3081.

Dose-adjusted EPOCH plus rituximab (regimen includes IT methotrexate)

Dunleavy K, Pittaluga S, Shovlin M, et al. Low-intensity therapy in adults with Burkitt's lymphoma. *N Engl J Med* 2013;369:1915-1925.

Roschewski M, Dunleavy K, Abramson JS, et al. Risk-adapted therapy in adults with Burkitt lymphoma: Results of NCI 9177, a multicenter prospective phase II study of DA-EPOCH-R [abstract]. *Blood* 2017;130 (Suppl 1):Abstract 188.

HyperCVAD (cyclophosphamide, vincristine, doxorubicin, and dexamethasone) alternating with high-dose methotrexate and cytarabine + rituximab

Thomas DA, Faderl S, O'Brien S, Bueso-Ramos C, et al. Chemoimmunotherapy with hyper-CVAD plus rituximab for the treatment of adult Burkitt and Burkitt-type lymphoma or acute lymphoblastic leukemia. *Cancer* 2006;106:1569-1580.

Thomas DA, Kantarjian HM, Faderl S, et al. Hyper-CVAD and rituximab for de novo Burkitt lymphoma/leukemia [abstract]. *Blood* 2011;118:Abstract 2698.

Second-line Therapy

RICE (rituximab, ifosfamide, carboplatin, etoposide)

Griffin TC, Weitzman S, Weinstein H, et al. A study of rituximab and ifosfamide, carboplatin, and etoposide chemotherapy in children with recurrent/refractory B-cell (CD20+) non-Hodgkin lymphoma and mature B-cell acute lymphoblastic leukemia: A report from the Children's Oncology Group. *Pediatr Blood Cancer* 2009;52:177-181.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



ADDITIONAL DIAGNOSTIC TESTING

ESSENTIAL:

- Adequate immunophenotyping to establish diagnosis^a and subclassification (eg, DLBCL, Burkitt lymphoma, plasmablastic lymphoma, primary effusion lymphoma [PEL])
 - ▶ IHC panel: CD45, CD20, CD3, CD10, BCL2, BCL6, Ki-67, CD138, kappa/lambda, HHV8 LANA,^b CD30 for PEL, with or without
 - ▶ Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: kappa/lambda, CD45, CD3, CD5, CD19, CD10, CD20
- EBER-ISH

→ [See Workup \(AIDS-2\)](#)

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- Molecular analysis to detect: antigen receptor gene rearrangements; *BCL2*; *BCL6*; *MYC* rearrangements
- Karyotype or FISH: *BCL2*; *BCL6*; *MYC*

^a [See Use of Immunophenotyping/Genetic Testing in Differential Diagnosis of Mature B-Cell and NK/T-Cell Neoplasms \(NHODG-A\).](#)

^b HHV8 can also be detected by PCR.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




WORKUP

ESSENTIAL

- Physical exam: attention to node-bearing areas, including Waldeyer's ring, and to size of liver and spleen
- Performance status
- B symptoms
- CBC with differential
- LDH
- Comprehensive metabolic panel
- Uric acid, phosphate
- C/A/P CT with contrast of diagnostic quality and/or whole-body PET/CT scan
- CD4 count
- Lumbar puncture, except for PEL
- HIV viral load
- Hepatitis B testing^c
- Hepatitis C testing^d
- Echocardiogram or MUGA scan if anthracycline or anthracenedione-based regimen is indicated
- Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)

USEFUL IN SELECTED CASES:

- UGI/barium enema/endoscopy
- Adequate bone marrow biopsy (>1.6 cm) ± aspirate; bone marrow biopsy is not necessary if PET/CT scan demonstrates bone disease. Bone marrow biopsy with a negative PET/CT scan may reveal discordant lymphoma
- Neck CT with contrast
- Plain bone radiographs and bone scan
- Brain MRI with contrast, or head CT with contrast
- Beta-2-microglobulin
- EBV viral load
- Quantitative immunoglobulins
- Discussion of fertility issues and sperm banking



^c Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen ([See NHODG-B](#)). If positive, check viral load and consider consult with gastroenterologist.

^d Hepatitis C antibody and if positive, viral load and consult with hepatologist.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

AIDS-Related B-Cell Lymphomas

TREATMENT

Antiretroviral therapy (ART) can be administered safely with chemotherapy but consultation with an HIV specialist or pharmacist is important to optimize compatibility. With continued development of new ARTs, effective alternatives are often available to patients when the existing ARTs are expected to affect metabolism of or share toxicities with chemotherapy. In general, avoidance of zidovudine, cobicistat, and ritonavir is strongly recommended. Concurrent ART is associated with higher CR rates (Barta et al. Blood 2013;122:3251-3262). For principles of concurrent HIV management and supportive care, see the [NCCN Guidelines for Cancer in People Living with HIV](#).^e

Burkitt lymphoma

- Suggested regimens:^{f,g}
 - Preferred regimens
 - ◇ CODOX-M/IVAC (modified): cyclophosphamide, vincristine, doxorubicin, high-dose methotrexate alternating with ifosfamide, etoposide, high-dose cytarabine + rituximab^h
 - ◇ DA-EPOCH-R: etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin + rituximabⁱ
 - Other recommended regimen
 - ◇ R-HyperCVAD rituximab, cyclophosphamide, vincristine, doxorubicin, and dexamethasone alternating with high-dose methotrexate and cytarabine^h
- If CD4 <50, maximize supportive care and monitor closely for cytopenias and infections while administering lymphoma therapy
- GCSF for all patients

For relapse, see [second-line regimens \(BURK-A\)](#)

Primary CNS lymphoma

- Initiate ART, if not already receiving
- Even with poorly controlled HIV and/or marginal performance status, consider high-dose methotrexate^j
- For select patients with good performance status on ART, see the [NCCN Guidelines for CNS - Primary CNS Lymphoma](#)
- Consider RT alone as initial management of patients who are not candidates for systemic therapy at presentation, then reassess eligibility for systemic therapy after RT
- Best supportive care (See [NCCN Guidelines for Palliative Care](#))

For relapse, see the [NCCN Guidelines for CNS - Primary CNS Lymphoma](#)

^e In the [NCCN Guidelines for Cancer in People Living with HIV](#), see the Principles of HIV Management While Undergoing Cancer Therapy; Principles of Systemic Therapy and Drug-Drug Interactions; and Principles of Supportive Care.

^f See references for regimens ([AIDS-A](#)).

^g An FDA-approved biosimilar is an appropriate substitute for rituximab.

^h High-risk patients presenting with symptomatic CNS disease should be started with the portion of the systemic therapy that contains CNS penetrating drugs.

ⁱ For R-EPOCH dosing, see Dunleavy K, Pittaluga S, Shovlin M, et al. Low-intensity therapy in adults with Burkitt's lymphoma. N Engl J Med 2013;369:1915-1925. Roschewski M, Dunleavy K, Abramson JS, et al. Risk-adapted therapy in adults with Burkitt lymphoma: Results of NCI 9177, a multicenter prospective phase II study of DA-EPOCH-R [abstract]. Blood 2017;130 (Suppl 1):Abstract 188.

^j Gupta N, Nolan A, Omuro A, et al. Long-term survival in AIDS-related primary central nervous system lymphoma. Neuro Oncol 2017;19:99-108.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 2.2020

AIDS-Related B-Cell Lymphomas

TREATMENT

ART can be administered safely with chemotherapy but consultation with an HIV specialist or pharmacist is important to optimize compatibility. With continued development of new ARTs, effective alternatives are often available to patients when the existing ARTs are expected to affect metabolism of or share toxicities with chemotherapy. In general, avoidance of zidovudine, cobicistat, and ritonavir is strongly recommended. Concurrent ART is associated with higher CR rates (Barta et al. Blood 2013;122:3251-3262). For principles of supportive care, see the [NCCN Guidelines for Cancer in People Living with HIV](#).^e



Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

^e In the [NCCN Guidelines for Cancer in People Living with HIV](#), see the Principles of HIV Management While Undergoing Cancer Therapy; Principles of Systemic Therapy and Drug-Drug Interactions; and Principles of Supportive Care.

^f See references for regimens ([AIDS-A](#)).

^g An FDA-approved biosimilar is an appropriate substitute for rituximab.

^k [See Recommendations for EPOCH ± Rituximab Dose Adjustments for Non-Burkitt Lymphomas \(AIDS-B\)](#).

^l [See Prognostic Model to Assess the Risk of CNS Disease \(BCEL-A 2 of 2\)](#).

^m Management can also apply to HIV-negative plasmablastic lymphoma.

ⁿ High-risk features include an age-adjusted IPI higher than 2, presence of MYC gene rearrangement, or TP53 gene deletion. Note that HIV-negative patients with plasmablastic lymphoma are generally considered to have higher risk disease. Optimization of HIV control with antiretroviral therapy is important.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**SUGGESTED TREATMENT REGIMENS**
REFERENCES**CODOX-M/IVAC (cyclophosphamide, vincristine, doxorubicin, high-dose methotrexate alternating with ifosfamide, etoposide, high-dose cytarabine) ± rituximab**

Wang ES, Straus DJ, Teruya-Feldstein J, et al. Intensive chemotherapy with cyclophosphamide, doxorubicin, high-dose methotrexate/ifosfamide, etoposide, and high-dose cytarabine (CODOX-M/IVAC) for human immunodeficiency virus-associated Burkitt lymphoma. *Cancer* 2003;98:1196-1205.

Barnes JA, LaCasce AS, Feng Y, et al. Evaluation of the addition of rituximab to CODOX-M/IVAC for Burkitt's lymphoma: A retrospective analysis. *Ann Oncol* 2011; 22:1859-1864.

Noy A, Lee JY, Cesarman E, et al. AMC 048: modified CODOX-M/IVAC-rituximab is safe and effective for HIV-associated Burkitt lymphoma. *Blood* 2015;126:160-166.

Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin)

Little RF, Pittaluga S, Grant N, et al. Highly effective treatment of acquired immunodeficiency syndrome-related lymphoma with dose-adjusted EPOCH: impact of antiretroviral therapy suspension and tumor biology. *Blood* 2003;101:4653-4659.

Roschewski M, Dunleavy K, Abramson JS, et al. Risk-adapted therapy in adults with Burkitt lymphoma: Results of NCI 9177, a multicenter prospective phase II study of DA-EPOCH-R [abstract]. *Blood* 2017;130 (Suppl 1):Abstract 188.

EPOCH + rituximab

Barta SK, Lee JY, Kaplan LD, et al. Pooled analysis of AIDS malignancy consortium trials evaluating rituximab plus CHOP or infusional EPOCH chemotherapy in HIV-associated non-Hodgkin lymphoma. *Cancer* 2012;118:3977-3983.

Bayraktar UD, Ramos JC, Petrich A, et al. Outcome of patients with relapsed/refractory acquired immune deficiency syndrome-related lymphoma diagnosed 1999-2008 and treated with curative intent in the AIDS Malignancy Consortium. *Leuk Lymphoma* 2012;53:2383-2389.

Dunleavy K, Pittaluga S, Shovlin M, et al. Low-intensity therapy in adults with Burkitt's lymphoma. *N Engl J Med* 2013;369:1915-1925.

Ramos J, Sparano J, Rudek M, et al. Safety and preliminary efficacy of vorinostat with R-EPOCH in high-risk HIV-associated non-Hodgkin's lymphoma (AMC-075). *Clin Lymphoma Myeloma Leuk* 2018;18:180-190.

HyperCVAD (cyclophosphamide, vincristine, doxorubicin, and dexamethasone alternating with high-dose methotrexate and cytarabine) ± rituximab

Cortes J, Thomas D, Rios A, et al. Hyperfractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone and highly active antiretroviral therapy for patients with acquired immunodeficiency syndrome-related Burkitt lymphoma/leukemia. *Cancer* 2002;94:1492-1499.

Thomas DA, Faderl S, O'Brien S, et al. Chemoimmunotherapy with hyper-CVAD plus rituximab for the treatment of adult Burkitt and Burkitt-type lymphoma or acute lymphoblastic leukemia. *Cancer* 2006;106:1569-1580.

Thomas DA, Kantarjian HM, Faderl S, et al. Hyper-CVAD and rituximab for de novo Burkitt lymphoma/leukemia [abstract]. *Blood* 2011;118:Abstract 2698.

CHOP + rituximab

Boue F, Gabarre J, Gisselbrecht C, et al. Phase II trial of CHOP plus rituximab in patients with HIV-associated non-Hodgkin's lymphoma. *J Clin Oncol* 2006;24:4123-4128.

Ribera JM, Oriol A, Morgades M, et al. Safety and efficacy of cyclophosphamide, adriamycin, vincristine, prednisone and rituximab in patients with human immunodeficiency virus-associated diffuse large B-cell lymphoma: results of a phase II trial. *Br J Haematol* 2008;140:411-419.

Rituximab and CD4 counts

Sparano JA, Lee JY, Kaplan LD et al. Rituximab plus concurrent infusional EPOCH chemotherapy is highly effective in HIV-associated B-cell non-Hodgkin lymphoma. *Blood* 2010;115:3008-3016.

Kaplan LD, Lee JY, Ambinder RF, et al. Rituximab does not improve clinical outcome in a randomized phase 3 trial of CHOP with or without rituximab in patients with HIV-associated non-Hodgkin lymphoma: AIDS-Malignancies Consortium Trial 010. *Blood* 2005;106:1538-1543.

Barta SK, Xue X, Wang D, et al. Treatment factors affecting outcomes in HIV-associated non-Hodgkin lymphomas: a pooled analysis of 1546 patients. *Blood* 2013;122:3251-3262.

Barta SK, Lee JY, Kaplan LD, et al. Pooled analysis of AIDS malignancy consortium trials evaluating rituximab plus CHOP or infusional EPOCH chemotherapy in HIV-associated non-Hodgkin lymphoma. *Cancer* 2012;118:3977-3983.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

AIDS-Related B-Cell Lymphomas

RECOMMENDATIONS FOR EPOCH ± RITUXIMAB DOSE ADJUSTMENTS FOR NON-BURKITT LYMPHOMAS¹

EPOCH ± rituximab dosing recommendations for cycle 1:

- Rituximab (if CD20-positive) 375 mg/m² IV on Day 1
- Etoposide 50 mg/m²/day continuous IV infusion for 4 days (96 hours)
- Doxorubicin 10 mg/m²/day continuous IV infusion for 4 days (96 hours)
- Vincristine 0.4 mg/m²/day continuous IV infusion for 4 days (96 hours)
- Day 5 cyclophosphamide dosing
 - ▶ If baseline CD4 count is >200 cells/mm³, start cyclophosphamide at 750 mg/m²
 - ▶ If baseline CD4 count is 50–200 cells/mm³, start cyclophosphamide at 375 mg/m²
 - ▶ For baseline CD4 counts <50 cells/mm³, doses of cyclophosphamide above 187.5 mg/m² have not been published.
- Prednisone 60 mg/m²/day for 5 days.

EPOCH Dose Modifications for Subsequent Cycles Based on Cytopenias (Non-Burkitt Lymphomas)¹	
Event	Action
ANC nadir on any cycle <500/mm ³ on 2 nonconsecutive days at least 3 days apart and/or platelet nadir <25,000/mm ³ in the previous cycle	Reduce cyclophosphamide dose by 187 mg/m ²
ANC nadir <500/mm ³ x ≥3 days or platelets <25,000/mm ³ x ≥3 days, AND patient was receiving no cyclophosphamide in the previous cycle	Reduce doxorubicin and etoposide by 25% of the full dose
ANC nadir ≥500/mm ³ AND platelet nadir ≥50,000/mm ³ in the previous cycle	Increase cyclophosphamide dose by 187 mg/m ² each cycle to a maximum dose of 750 mg/m ²

¹ For R-EPOCH dosing for non-Burkitt lymphomas, see Ramos J, Sparano J, Rudek M, et al. Safety and preliminary efficacy of vorinostat with R-EPOCH in high-risk HIV-associated non-Hodgkin's lymphoma (AMC-075). Clin Lymphoma Myeloma Leuk 2018;18:180-190. This is an ongoing clinical trial and the utility of adding vorinostat to R-EPOCH has not yet been established.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Lymphoblastic Lymphoma

ADDITIONAL DIAGNOSTIC TESTING^{a,b}

ESSENTIAL:

- Adequate immunophenotyping to establish diagnosis^c
 - ▶ IHC panel: CD45, CD19, CD20, CD79a, CD3, CD2, CD5, CD7, TdT, CD1a, CD10, cyclin D1 with or without
 - ▶ Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: kappa/lambda, CD45, CD3, CD5, CD4, CD7, CD8, CD19, CD20, CD10, TdT, CD13, CD33, CD1a, cytoplasmic CD3, CD22, myeloperoxidase
- Karyotype ± FISH: *MYC*; t(9;22); t(8;14); and variants or PCR for *BCR-ABL*

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- Additional immunohistochemical studies to establish lymphoma subtype
 - ▶ Paraffin panel: CD22, CD4, CD8, cyclin D1
- Molecular analysis to detect: antigen receptor gene rearrangements

WORKUP

ESSENTIAL:

- Physical exam: attention to node-bearing areas, including Waldeyer's ring, and to size of liver and spleen
- Performance status
- B symptoms
- CBC with differential
- LDH
- Comprehensive metabolic panel
- Uric acid, phosphate
- C/A/P CT with contrast of diagnostic quality
- Lumbar puncture
- Flow cytometry of cerebrospinal fluid
- Bilateral or unilateral bone marrow biopsy ± aspirate with flow and cytogenetics
- Hepatitis B testing^d
- Echocardiogram or MUGA scan if anthracycline or anthracenedione-based regimen is indicated
- Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)

USEFUL IN SELECTED CASES:

- Brain MRI with contrast
- Discussion of fertility issues and sperm banking
- Beta-2-microglobulin
- Whole-body PET/CT scan^e

→ See NCCN
Guidelines for
Acute
Lymphoblastic
Leukemia

^a The lymphoblastic lymphoma (LL) category comprises two diseases, T-cell LL (LL-T; 90%) and B-cell LL (LL-B; 10%), which corresponds to T-ALL and B-ALL, respectively, with presentations in extramedullary sites.

^b This disease is complex and curable; it is preferred that treatment occur at centers with expertise in the management of the disease.

^c Typical immunophenotype: LL-B: slg-, CD10+/-, CD19+, CD20+/-, TdT+.

LL-T: slg-, CD10-, CD19/20-, CD3-/+, CD4/8+/-, CD1a+/-, TdT+, CD2+, CD7+ cytoplasmic CD3+, sCD3+/-.

^d Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen ([See NHODG-B](#)). If positive, check viral load and consider consult with gastroenterologist.

^e Initiation of therapy should not be delayed in order to obtain a PET/CT scan.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Post-Transplant Lymphoproliferative Disorders

ADDITIONAL DIAGNOSTIC TESTING

ESSENTIAL:

- Adequate immunophenotyping to establish diagnosis
 - IHC panel: CD3, CD5, CD10, BCL6, BCL2, IRF4/MUM1, CD20, CD79a, PAX5, Ki-67, kappa, lambda with or without
 - Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: CD3, CD5, CD7, CD4, CD8, CD19, CD20, CD10, kappa, lambda
- Epstein-Barr virus (EBV) evaluation by EBV-LMP1 or EBER-ISH (if EBV-LMP1 negative, EBER-ISH is recommended)

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- Additional immunophenotyping
 - IHC panel: CD15, CD30, CD45, CD7, CD4, CD8, ALK, TIA-1, Granzyme B, CD57, CD56, CD138
 - Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: CD138, cytoplasmic kappa and lambda, CD30, CD57, CD56, CD16, CD25, CD52
- Molecular analysis to detect: *IGHV* gene rearrangements
- *BCL6* gene mutation analysis
- EBV by southern blot

WORKUP

ESSENTIAL:

- Performance status
- Albumin
- History of therapy for transplant
- LDH, electrolytes, BUN, creatinine
- CBC with differential
- Hepatitis B testing^a
- EBV PCR for cell-free plasma EBV DNA marker^b
- C/A/P CT with contrast and/or whole-body PET/CT scan
- Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)

USEFUL IN SELECTED CASES:

- Echocardiogram or MUGA scan if treatment includes regimens containing anthracyclines or anthracenediones
- Bone marrow evaluation
- Brain MRI with and without contrast
- CMV PCR
- EBV serology for primary versus reactivation

PTLD SUBTYPE^c

Non-destructive lesions^d

Monomorphic PTLD (B-cell type)

Polymorphic PTLD

Monomorphic PTLD (T-cell type)

Classical Hodgkin lymphoma PTLD

[See First-line Therapy \(PTLD-2\)](#)[See First-line Therapy \(PTLD-3\)](#)

- Other than reduction of immunosuppression (RI), there are no established treatments.
- Consider treatment with multiagent regimens ([see Suggested Regimens PTLD-A](#)); autologous HCT may not be appropriate.

[See NCCN Guidelines for Hodgkin Lymphoma](#)

^a Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen ([See NHODG-B](#)). If positive, check viral load and consider consult with gastroenterologist.

^b If EBV-negative, should not be used as response marker.

^c Indolent small B-cell lymphomas arising in transplant recipients are not included among PTLDs, with the exception of EBV-positive marginal zone lymphomas.

^d Non-destructive lesions are of B-cell type and include plasmacytic hyperplasia, infectious mononucleosis, and florid follicular hyperplasia.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Post-Transplant Lymphoproliferative Disorders



^d Non-destructive lesions are of B-cell type and include plasmacytic hyperplasia, infectious mononucleosis, and florid follicular hyperplasia.

^e Treatment is based on the unique histology.

^f Response to RI is variable and patients need to be closely monitored; RI should be coordinated with the transplant team. RI: Reduction in calcineurin inhibition (cyclosporin and tacrolimus), discontinuation of antimetabolic agents (azathioprine and mycophenolate mofetil), and for critically ill patients all non-glucocorticoid immunosuppression should be discontinued.

^g As part of a step-wise approach in patients who are not highly symptomatic or cannot tolerate chemotherapy secondary to comorbidity.

Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

^h For concurrent or sequential chemoimmunotherapy, [see Suggested Treatment Regimens \(PTLD-A\)](#).

^j Restage in two to four weeks.

^j Re-escalation of immunosuppressive therapy should be individualized, taking into account the extent of initial RI and the nature of the organ allograft. These decisions should be made in conjunction with the transplant team.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Post-Transplant Lymphoproliferative Disorders



Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

^f Response to RI is variable and patients need to be closely monitored; RI should be coordinated with the transplant team. RI: Reduction in calcineurin inhibition (cyclosporin and tacrolimus), discontinuation of antimetabolic agents (azathioprine and mycophenolate mofetil), and for critically ill patients all non-glucocorticoid immunosuppression should be discontinued. Response to RI is variable and patients need to be closely monitored; RI should be coordinated with the transplant team.

^h For concurrent or sequential chemoimmunotherapy, [see Suggested Treatment Regimens \(PTLD-A\)](#).

ⁱ Restage in two to four weeks.

^k [See Principles of Radiation Therapy \(NHODG-D\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Post-Transplant Lymphoproliferative Disorders

SUGGESTED TREATMENT REGIMENS^a

(in alphabetical order)

An FDA-approved biosimilar is an appropriate substitute for rituximab.

Monomorphic PTLD (B-cell type) and Polymorphic PTLD

Sequential chemoimmunotherapy

- Rituximab 375 mg/m² weekly x 4 weeks (on days 1, 8, 15, and 22)^c

- ▶ Restage with PET/CT scan

- ◊ If PET/CT scan negative, rituximab 375 mg/m² every 3 weeks x 4 cycles

- ◊ If PET/CT scan positive, CHOP-21 every 3 weeks + granulocyte colony-stimulating factor (G-CSF) (on days 50, 72, 94, and 116) x 4 cycles

Concurrent chemoimmunotherapy

- RCHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone)

- For frail patients who cannot tolerate anthracycline, no specific regimen has been identified but options may include:^b

- ▶ RCEPP (rituximab, cyclophosphamide, etoposide, prednisone, procarbazine)

- ▶ RCEOP (rituximab, cyclophosphamide, etoposide, vincristine, prednisone)

- ▶ RCVP (rituximab, cyclophosphamide, vincristine, prednisone)

Monomorphic PTLD (T-cell type)

- Brentuximab vedotin + CHP (cyclophosphamide, doxorubicin, prednisone) for CD30+ cases^d

- CHOP

- CHOEP

- For frail patients who cannot tolerate anthracycline, no specific regimen has been identified but options may include:

- ▶ CVP

- ▶ CEOP

- ▶ CEPP

Primary CNS PTLD (B-cell type)

- High-dose methotrexate^e + rituximab

Consider prophylaxis for tumor lysis syndrome ([See NHODG-B](#))
See monoclonal antibody and viral reactivation ([NHODG-B](#))

^a Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^b There are no published data regarding the use of these regimens; however, they are used at NCCN Member Institutions for the treatment of PTLD.

^c Trappe R, Oertel S, Leblond V, et al. Sequential treatment with rituximab followed by CHOP chemotherapy in adult B-cell post-transplant lymphoproliferative disorder (PTLD): the prospective international multicentre phase 2 PTLD-1 trial. *Lancet Oncol* 2012;13:196-206.

^d Horwitz S, O'Connor OA, Pro B, et al. Brentuximab vedotin with chemotherapy for CD30-positive peripheral T-cell lymphoma (ECHELON-2): a global, double-blind, randomised, phase 3 trial. *Lancet* 2019;393:229-240.

^e Patients with PTLD often have renal insufficiency. High-dose methotrexate should be used with caution. Alternate regimens (cytarabine-based) should be considered.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Castleman's Disease

ADDITIONAL DIAGNOSTIC TESTING^{a,b,c}

ESSENTIAL:

- Adequate immunophenotyping to establish diagnosis
 - IHC panel: kappa/lambda, CD20, CD3, CD5, CD138, HHV8
 - EBER-ISH

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- Molecular analysis to detect: immunoglobulin and *TCR* gene rearrangements
- IHC: Ki-67 index; Ig heavy chains,^d CD10, BCL2, BCL6, cyclin D1, CD21, or CD23, CD38, IRF4/MUM1, PAX5
- Cell surface marker analysis by flow cytometry with peripheral blood and/or biopsy specimen: kappa/lambda, CD19, CD20, CD5, CD23, CD10

WORKUP^e

ESSENTIAL:

- Physical exam: attention to node-bearing areas, including Waldeyer's ring, and to size of liver and spleen
- Performance status
- Assess for criteria for active disease^f
- CBC with differential
- Comprehensive metabolic panel
- LDH, CRP, ESR
- Beta-2-microglobulin, SPEP and urine electrophoresis with immunofixation, serum light chains, quantitative immunoglobulins
- HIV ELISA, HHV8 DNA titer by PCR, hepatitis B testing,^g EBV DNA titer by PCR
- Whole-body PET/CT scan (preferred) or C/A/P CT with contrast of diagnostic quality
- Pregnancy testing in women of child-bearing age (if chemotherapy or RT planned)

USEFUL UNDER CERTAIN CIRCUMSTANCES

- If HHV8 or HIV positive, screening for concurrent Kaposi's sarcoma is strongly recommended
- Bone marrow biopsy + aspirate
- Reticulin fibrosis of bone marrow (particularly in patients with TAFRO syndrome)
- Neck CT with contrast
- Echocardiogram or MUGA scan if anthracycline or anthracenedione-based regimen is indicated
- sIL6, sIL10, VEGF, uric acid, ferritin^h
- Hepatitis C testing
- Discussion of fertility issues and sperm banking

Unicentric
Castleman's
disease

→ [See
CD-2](#)

Multicentric
Castleman's
disease (MCD)

→ [See
CD-3](#)

^a For AIDS-related lymphoma associated with Castleman's disease, [see AIDS-1](#). For DLBCL-associated with Castleman's disease in non-HIV patients, [see BCEL-1](#).

^b There are 2 variants – hyaline vascular (virtually always unicentric, HHV8-) and plasma cell (may be multicentric, often HHV8+, +/- HIV+).

^c Two types of DLBCL are associated with the HHV8+ PC type: plasmablastic (EBV-) and "germinotropic" (EBV+).

^d In plasma cell variant HHV8+, plasmablasts are IgM lambda while normal plasma cells are IgG or A polytypic.

^e [See Subtypes of Idiopathic MCD \(CD-A\)](#). Rule out other diseases that can mimic idiopathic MCD ([see Exclusion Criteria on CD-B 1 of 2](#)). If concurrent polyneuropathy and monoclonal plasma cell disorder, a workup for POEMS syndrome is recommended.

^f [See Criteria for Active Disease \(CD-A\)](#).

^g Hepatitis B testing is indicated because of the risk of reactivation with immunotherapy + chemotherapy. Tests include hepatitis B surface antigen and core antibody for a patient with no risk factors. For patients with risk factors or previous history of hepatitis B, add e-antigen ([See NHODG-B](#)). If positive, check viral load and consider consult with gastroenterologist.

^h Measurement of acute phase reactants maybe helpful in monitoring therapy.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PRIMARY TREATMENT^{i,j}

SECOND-LINE THERAPY^{i,j}



ⁱ Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^j An FDA-approved biosimilar is an appropriate substitute for rituximab.

^k [See Principles of Radiation Therapy \(NHODG-D\).](#)

^l Patients with non-bulky disease may be observed after RT.

^m Encourage biopsy to rule out transformation to DLBCL or concomitant development of other malignancies or opportunistic infections.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Castleman's Disease



^f See [Criteria for Active Disease \(CD-A\)](#).

ⁱ Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^j An FDA-approved biosimilar is an appropriate substitute for rituximab.

^m Encourage biopsy to rule out transformation to DLBCL or concomitant development of other malignancies or opportunistic infections.

ⁿ See [Diagnostic Criteria for Idiopathic MCD \(CD-B\)](#).

^o All HIV+ patients should be on combination antiretroviral therapy (cART).

^p Patients with hyaline vascular histology do not benefit from siltuximab.

^q Response assessment using the imaging modalities performed during workup (C/A/P CT with contrast or PET/CT).

^r Occult Kaposi sarcoma (KS) is prevalent in HIV/HHV8+ MCD and may flare after rituximab or prednisone. Consider baseline imaging and direct visualization to screen for pulmonary ± GI KS as well as concurrent KS-directed therapy (ie, addition of liposomal doxorubicin).

^s Rituximab ± prednisone may repeat without limit if progression ≥6 months after completion of rituximab.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PRIMARY TREATMENT^{i,j}

RELAPSED DISEASE



ⁱ Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^j An FDA-approved biosimilar is an appropriate substitute for rituximab.

^m Encourage biopsy to rule out transformation to DLBCL or concomitant development of other malignancies or opportunistic infections.

^q Response assessment using the imaging modalities performed during workup (C/A/P CT with contrast or PET/CT).

^r Occult Kaposi sarcoma (KS) is prevalent in HIV/HHV8+ MCD and may flare after rituximab or prednisone. Consider baseline imaging and direct visualization to screen for pulmonary ± GI KS as well as concurrent KS-directed therapy (ie, addition of liposomal doxorubicin).

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Castleman's Disease

REFRACTORY OR PROGRESSIVE DISEASE^{i,j}



ⁱ Rituximab and hyaluronidase human injection for subcutaneous use may be substituted for rituximab after patients have received the first full dose of rituximab by intravenous infusion. This substitution cannot be made for rituximab used in combination with ibritumomab tiuxetan.

^j An FDA-approved biosimilar is an appropriate substitute for rituximab.

^m Encourage biopsy to rule out transformation to DLBCL or concomitant development of other malignancies or opportunistic infections.

^q Response assessment using the imaging modalities performed during workup (C/A/P CT with contrast or PET/CT).

^t Single-agent therapy is preferred for asymptomatic patients with no organ failure; combination therapy is preferred for patients with fulminant disease and organ failure.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



CRITERIA FOR ACTIVE DISEASE^a

- **Fever**
- **Increased serum C-reactive protein level >20 mg/L in the absence of any other etiology**
- **At least three of the following other MCD-related symptoms:**
 - **Peripheral lymphadenopathy**
 - **Enlarged spleen**
 - **Edema**
 - **Pleural effusion**
 - **Ascites**
 - **Cough**
 - **Nasal obstruction**
 - **Xerostomia**
 - **Rash**
 - **Central neurologic symptoms**
 - **Jaundice**
 - **Autoimmune hemolytic anemia**

^a Gérard L, Bérezné A, Galicier L, et al. Prospective study of rituximab in chemotherapy-dependent human immunodeficiency virus-associated multicentric Castleman's disease: ANRS 117 Castlemab Trial. J Clin Oncol 2007;25:3350-3356.

SUBTYPES OF IDIOPATHIC MCD^a

- **Idiopathic MCD (iMCD-TAFRO)**
 - **Marked inflammatory syndrome**
 - **Thrombocytopenia, anasarca, fever/elevated C-reactive protein (CRP), renal dysfunction/reticulिन myelofibrosis, organomegaly**
 - **Megakaryocytic hyperplasia, hypervascular or mixed histopathology**
 - **Normal immunoglobulin levels**
- **Idiopathic MCD (iMCD-NOS)**
 - **Less intense inflammatory syndrome**
 - **Normal/elevated platelet counts**
 - **Plasmacytic or mixed histopathology**
 - **Polyclonal hypergammaglobulinemia**

^a Iwaki N, Fajgenbaum DC, Nabel CS, et al. Clinicopathologic analysis of TAFRO syndrome demonstrates a distinct subtype of HHV-8-negative multicentric Castleman disease. Am J Hematol 2016 ;91:220-226.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

Castleman's Disease

Consensus Diagnostic Criteria for Idiopathic MCD^a

I. Major Criteria (need both):	1. Histopathologic lymph node features consistent with the iMCD spectrum. Features along the iMCD spectrum include (need grade 2–3 for either regressive GCs or plasmacytosis at minimum): <ul style="list-style-type: none"> - Regressed/atrophic/atretic germinal centers, often with expanded mantle zones composed of concentric rings of lymphocytes in an “onion skinning” appearance - FDC prominence - Vascularity, often with prominent endothelium in the interfollicular space and vessels penetrating into the GCs with a “lollipop” appearance - Sheetlike, polytypic plasmacytosis in the interfollicular space - Hyperplastic GCs 2. Enlarged lymph nodes (≥ 1 cm in short-axis diameter) in ≥ 2 lymph node stations	
II. Minor Criteria (need at least 2 of 11 criteria with at least 1 laboratory criterion)	Laboratory* <ol style="list-style-type: none"> 1. Elevated CRP (>10 mg/L) or ESR (>15 mm/h)[†] 2. Anemia (hemoglobin <12.5 g/dL for males, hemoglobin <11.5 g/dL for females) 3. Thrombocytopenia (platelet count <150 μmL) or thrombocytosis (platelet count >400 μmL) 4. Hypoalbuminemia (albumin <3.5 g/dL) 5. Renal dysfunction (eGFR <60 mL/min/1.73m²) or proteinuria (total protein 150 mg/24 h or 10 mg/100 mL) 6. Polyclonal hypergammaglobulinemia (total γ globulin or immunoglobulin G >1700 mg/dL) 	Clinical <ol style="list-style-type: none"> 1. Constitutional symptoms: night sweats, fever ($>38^{\circ}\text{C}$), weight loss, or fatigue (≥ 2 CTCAE lymphoma score for B-symptoms) 2. Large spleen and/or liver 3. Fluid accumulation: edema, anasarca, ascites, or pleural effusion 4. Eruptive cherry hemangiomatosis or violaceous papules 5. Lymphocytic interstitial pneumonitis
III. Exclusion Criteria (must rule out each of these diseases that can mimic iMCD)	<u>Infection-related disorders</u> <ol style="list-style-type: none"> 1. HHV-8 (infection can be documented by blood PCR; diagnosis of HHV-8–associated MCD requires positive LANA-1 staining by IHC, which excludes iMCD) 2. Clinical EBV-lymphoproliferative disorders such as infectious mononucleosis or chronic active EBV (detectable EBV viral load not necessarily exclusionary) 3. Inflammation and adenopathy caused by other uncontrolled infections (eg, acute or uncontrolled CMV, toxoplasmosis, HIV, active tuberculosis) 	<u>Autoimmune/autoinflammatory diseases</u> (requires full clinical criteria; detection of autoimmune antibodies alone is not exclusionary) <ol style="list-style-type: none"> 1. Systemic lupus erythematosus 2. Rheumatoid arthritis 3. Adult-onset Still disease 4. Juvenile idiopathic arthritis 5. Autoimmune lymphoproliferative syndrome <u>Malignant/lymphoproliferative disorders</u> (these disorders must be diagnosed before or at the same time as iMCD to be exclusionary): <ol style="list-style-type: none"> 1. Lymphoma (Hodgkin and non-Hodgkin) 2. Multiple myeloma 3. Primary lymph node plasmacytoma 4. FDC sarcoma 5. POEMS syndrome[‡]

[Continued](#)

^a Fajgenbaum D, et al. International, evidence-based consensus diagnostic criteria for HHV-8-negative/idiopathic multicentric Castleman disease. Blood 2017;129:1646-1657.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



Consensus Diagnostic Criteria for Idiopathic MCD (continued)^a

- Select additional features supportive of, but not required for diagnosis
 - ▶ Elevated IL-6, sIL-2R, VEGF, IgA, IgE, LDH, and/or B2M
 - ▶ Reticulin fibrosis of bone marrow (particularly in patients with TAFRO syndrome)
 - ▶ Diagnosis of disorders that have been associated with iMCD: paraneoplastic pemphigus, bronchiolitis obliterans organizing pneumonia, autoimmune cytopenias, polyneuropathy (without diagnosing POEMS[‡]), glomerular nephropathy, inflammatory myofibroblastic tumor

^a Fajgenbaum D, et al. International, evidence-based consensus diagnostic criteria for HHV-8-negative/idiopathic multicentric Castleman disease. Blood 2017;129:1646-1657.

Footnotes for [CD-B 1 of 2](#)

* Laboratory cutoff thresholds are provided as guidance. Since some laboratories have slightly different ranges, the upper and lower ranges from a particular laboratory should be used to determine if a patient meets a particular laboratory Minor Criterion.

† Evaluation of CRP is mandatory and tracking CRP levels is highly recommended, but ESR will be accepted if CRP is not available.

‡ POEMS is considered to be a disease “associated” with CD. Because the monoclonal plasma cells are believed to drive the cytokine storm, we do not consider it iMCD, but rather “POEMS-associated MCD.”

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



USE OF IMMUNOPHENOTYPING/GENETIC TESTING IN DIFFERENTIAL DIAGNOSIS OF MATURE B-CELL AND NK/T-CELL NEOPLASMS^a (TO BE USED IN CONJUNCTION WITH CLINICAL AND MORPHOLOGIC CORRELATION)

General Principles

- Morphology ± clinical features drive both the choice and the interpretation of special studies.
- Differential diagnosis is based on morphology ± clinical setting.
- Begin with a broad panel appropriate to morphologic diagnosis, limiting panel of antibodies based on the differential diagnosis.
 - Avoid “shotgun” panels of unnecessary antibodies unless a clinically urgent situation warrants.
- Add antigens in additional panels, based on initial results.
- Follow with genetic studies as needed.
- Return to clinical picture if immunophenotype + morphology are not specific.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

NHODG-A
1 OF 11



USE OF IMMUNOPHENOTYPING/GENETIC TESTING IN DIFFERENTIAL DIAGNOSIS OF MATURE B-CELL AND NK/T-CELL NEOPLASMS^a (TO BE USED IN CONJUNCTION WITH CLINICAL AND MORPHOLOGIC CORRELATION)

B-cell antigens positive^{b,c} (CD19, CD20, CD79a, PAX5)

- **Morphology**
 - **Cytology**
 - ◊ Small cells
 - ◊ Medium-sized cells
 - ◊ Large cells
 - **Pattern**
 - ◊ Diffuse
 - ◊ Nodular, follicular, mantle, marginal
 - ◊ Sinuses
- **Clinical**
 - Age (child, adult)
 - Location
 - ◊ Nodal
 - ◊ Extranodal, specific site
- **Immunophenotype**
 - Naïve B cells: CD5, CD23
 - GCB cells: CD10, BCL6
 - FDC: CD21, CD23
 - Post-GCB cells: IRF4/MUM1, CD138
 - Immunoglobulin heavy and light chains (surface, cytoplasmic, class switch, light chain type)
 - Oncogene products: BCL2, cyclin D1, MYC, BCL6, ALK
 - Viruses: EBV, HHV8
 - Other: CD43, Ki-67
- **Genetic testing**
 - BCL2, BCL6, CCND1, MYC, ALK, MYD88, BRAF, IG rearrangement

T- or NK/T-cell antigens positive^{b,c}

(CD2, CD3, CD5, CD7) [and B-cell antigens negative]

- **Morphology**
 - Anaplastic vs. non-anaplastic
 - Epidermotropic
- **Clinical**
 - Age (child, adult)
 - Location
 - ◊ Cutaneous
 - ◊ Extranodal noncutaneous (specific site)
 - ◊ Nodal
- **Immunophenotype**
 - CD30, ALK*, CD56, βF1, cytotoxic granule proteins
 - CD4, CD8, CD5, CD7, TCRαβ, TCRγδ, CD1a, TdT
 - Follicular T cells: CD10, BCL6, CD57, PD1/CD279
 - Viruses: EBV, HTLV1 (clonal)
- **Genetic testing**
 - ALK, TCR, HTLV1

*Always do ALK if CD30+

[See Initial Morphologic, Clinical, and Immunophenotypic Analysis \(NHODG-A 3 of 11\)](#)

^aThese are meant to be general guidelines. Interpretation of results should be based on individual circumstances and may vary. Not all tests will be required in every case.

^bSome lymphoid neoplasms may lack pan leukocyte (CD45), pan-B, and pan-T antigens. Selection of additional antibodies should be based on the differential diagnosis generated by morphologic and clinical features (eg, plasma cell myeloma, ALK+ DLBCL, plasmablastic lymphoma, anaplastic large cell lymphoma, NK-cell lymphomas).

^cUsually 1 Pan-B (CD20) and 1 Pan-T (CD3) markers are done unless a terminally differentiated B-cell or a specific PTCL is suspected.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



USE OF IMMUNOPHENOTYPING/GENETIC TESTING IN DIFFERENTIAL DIAGNOSIS OF MATURE B-CELL AND NK/T-CELL NEOPLASMS^a (TO BE USED IN CONJUNCTION WITH CLINICAL AND MORPHOLOGIC CORRELATION)

INITIAL MORPHOLOGIC, CLINICAL, AND IMMUNOPHENOTYPIC ANALYSIS



^aThese are meant to be general guidelines. Interpretation of results should be based on individual circumstances and may vary. Not all tests will be required in every case.

^dInitial panel will often include additional markers based on morphologic differential diagnosis and clinical features.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



USE OF IMMUNOPHENOTYPING/GENETIC TESTING IN DIFFERENTIAL DIAGNOSIS OF MATURE B-CELL AND NK/T-CELL NEOPLASMS^a (TO BE USED IN CONJUNCTION WITH CLINICAL AND MORPHOLOGIC CORRELATION)

B-CELL NEOPLASMS



Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 2.2020

B-Cell Lymphomas

USE OF IMMUNOPHENOTYPING/GENETIC TESTING IN DIFFERENTIAL DIAGNOSIS OF MATURE B-CELL AND NK/T-CELL NEOPLASMS^a (TO BE USED IN CONJUNCTION WITH CLINICAL AND MORPHOLOGIC CORRELATION)

B-CELL NEOPLASMS



^aThese are meant to be general guidelines. Interpretation of results should be based on individual circumstances and may vary. Not all tests will be required in every case.

^kStarry sky pattern is typically present in BL and frequently in U-DLBCL/BL.

^lKi-67 is a prognostic factor in some lymphomas (eg, mantle cell and is typically >90% in Burkitt lymphoma). It is not useful in predicting the presence of MYC rearrangement or in classification.

^mRare MCL may be cyclin D1-.

ⁿRare BL may lack detectable MYC rearrangement. Correlation with morphology and clinical features is essential.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 2.2020

B-Cell Lymphomas

USE OF IMMUNOPHENOTYPING/GENETIC TESTING IN DIFFERENTIAL DIAGNOSIS OF MATURE B-CELL AND NK/T-CELL NEOPLASMS^a (TO BE USED IN CONJUNCTION WITH CLINICAL AND MORPHOLOGIC CORRELATION)

B-CELL NEOPLASMS



Large cells:

- Diffuse large B-cell lymphoma (DLBCL), NOS
 - Germinal center B-cell type
 - Non-germinal center B-cell type
 - T-cell/histiocyte-rich large B-cell lymphoma (THRLBCL)
 - Primary DLBCL of the CNS
 - Primary cutaneous DLBCL, leg type
 - EBV-positive DLBCL, NOS (EBV + DLBCL)
- DLBCL associated with chronic inflammation
- Lymphomatoid granulomatosis
- Primary mediastinal (thymic) large B-cell lymphoma (PMBL)
- Intravascular large B-cell lymphoma
- ALK-positive large B-cell lymphoma
- Plasmablastic lymphoma
- HHV8+ Large B-cell lymphoma NOS
- Large B-cell Lymphoma (LBCL) with IRF4 rearrangement
- Primary effusion lymphoma
- B-cell lymphoma, unclassifiable, intermediate between DLBCL (U-DLBCL) and classical Hodgkin lymphoma (CHL)
- Mantle cell lymphoma (MCL), pleomorphic variant

^aThese are meant to be general guidelines. Interpretation of results should be based on individual circumstances and may vary. Not all tests will be required in every case.

^lKi-67 is a prognostic factor in some lymphomas (eg, mantle cell and is typically >90% in Burkitt lymphoma). It is not useful in predicting the presence of MYC rearrangement or in classification.

^oCD5 is included to identify pleomorphic MCL; if CD5 is positive, cyclin D1 staining is done to confirm or exclude MCL.

[Continued](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 2.2020

B-Cell Lymphomas

USE OF IMMUNOPHENOTYPING/GENETIC TESTING IN DIFFERENTIAL DIAGNOSIS OF MATURE B-CELL AND NK/T-CELL NEOPLASMS^a (TO BE USED IN CONJUNCTION WITH CLINICAL AND MORPHOLOGIC CORRELATION)

LARGE CELLS (continued)



^aThese are meant to be general guidelines. Interpretation of results should be based on individual circumstances and may vary. Not all tests will be required in every case.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



USE OF IMMUNOPHENOTYPING/GENETIC TESTING IN DIFFERENTIAL DIAGNOSIS OF MATURE B-CELL AND NK/T-CELL NEOPLASMS^a (TO BE USED IN CONJUNCTION WITH CLINICAL AND MORPHOLOGIC CORRELATION)

B-CELL NEOPLASMS



^aThese are meant to be general guidelines. Interpretation of results should be based on individual circumstances and may vary. Not all tests will be required in every case.


^PThese are assessed both in follicles (if present) and in intrafollicular/diffuse areas. CD10+ BCL6+ germinal centers are present in PCMZL, while both follicular and interfollicular/diffuse areas (tumor cells) are positive for BCL6+/- CD10 in PCFCL.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

USE OF IMMUNOPHENOTYPING/GENETIC TESTING IN DIFFERENTIAL DIAGNOSIS OF MATURE B-CELL AND NK/T-CELL NEOPLASMS^a
(TO BE USED IN CONJUNCTION WITH CLINICAL AND MORPHOLOGIC CORRELATION)

T-CELL NEOPLASMS



Anaplastic morphology

- **Anaplastic large cell lymphoma (ALCL), ALK positive**
- **Anaplastic large cell lymphoma (ALCL), ALK negative**
- **Adult T-cell leukemia/lymphoma (ATLL), anaplastic large cell type**
- **Enteropathy-associated T-cell lymphoma (EATL)**
- **Primary cutaneous CD30-positive T-cell lymphoproliferative disorders**
 - **Lymphomatoid papulosis (LyP)**
 - **Primary cutaneous anaplastic large cell lymphoma (PC-ALCL)**

^aThese are meant to be general guidelines. Interpretation of results should be based on individual circumstances and may vary. Not all tests will be required in every case.

⁹Rare T-cell lymphomas may be CD20+ or PAX5+. Assessment of other Pan-T and -B markers is essential. The expression of multiple markers of 1 lineage and only 1 of the other lineages supports lineage assignment. PCR analysis may be required to determine lineage in such cases.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 2.2020

B-Cell Lymphomas

USE OF IMMUNOPHENOTYPING/GENETIC TESTING IN DIFFERENTIAL DIAGNOSIS OF MATURE B-CELL AND NK/T-CELL NEOPLASMS^a (TO BE USED IN CONJUNCTION WITH CLINICAL AND MORPHOLOGIC CORRELATION)

T-CELL NEOPLASMS



^aThese are meant to be general guidelines. Interpretation of results should be based on individual circumstances and may vary. Not all tests will be required in every case.

^rA minority of MF cases can be CD30+, CD4-, CD8+/-, and TIA1+.

^sAECTCL has distinctive morphology and clinical presentation.

Note: All recommendations are category 2A unless otherwise indicated.



Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

USE OF IMMUNOPHENOTYPING/GENETIC TESTING IN DIFFERENTIAL DIAGNOSIS OF MATURE B-CELL AND NK/T-CELL NEOPLASMS^a (TO BE USED IN CONJUNCTION WITH CLINICAL AND MORPHOLOGIC CORRELATION)



^aThese are meant to be general guidelines. Interpretation of results should be based on individual circumstances and may vary. Not all tests will be required in every case.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



SUPPORTIVE CARE FOR B-CELL LYMPHOMAS

Tumor Lysis Syndrome (TLS)

- **Laboratory hallmarks of TLS:**

- ▶ High potassium
- ▶ High uric acid
- ▶ High phosphorous
- ▶ Low calcium

- **Symptoms of TLS:**

- ▶ Nausea and vomiting, shortness of breath, irregular heartbeat, clouding of urine, lethargy, and/or joint discomfort.

- **TLS features:**

- ▶ Consider TLS prophylaxis for patients with the following risk factors:
 - ◊ Histologies of Burkitt lymphoma and lymphoblastic lymphoma; occasionally DLBCL
 - ◊ Spontaneous TLS
 - ◊ Elevated WBC
 - ◊ Bone marrow involvement
 - ◊ Pre-existing elevated uric acid
 - ◊ Ineffectiveness/intolerance of allopurinol
 - ◊ Renal disease or renal involvement by tumor

- **Treatment of TLS:**

- ▶ TLS is best managed if anticipated and treatment is started prior to chemotherapy.
- ▶ Centerpiece of treatment includes:
 - ◊ Vigorous hydration
 - ◊ Management of hyperuricemia
 - ◊ Frequent monitoring of electrolytes and aggressive correction (essential)
- ▶ First-line and at retreatment for hyperuricemia
 - ◊ Allopurinol or febuxostat if intolerant to allopurinol beginning 2–3 days prior to chemotherapy and continued for 10–14 days or Rasburicase (doses of 3 to 6 mg are usually effective.^a One dose of rasburicase is frequently adequate. Re-dosing should be individualized) is indicated for patients with any of the following risk factors:
 - Urgent need to initiate therapy in a high-bulk patient
 - Situations where adequate hydration may be difficult or impossible
 - Acute renal failure
- ▶ If TLS is untreated, its progression may cause acute kidney failure, cardiac arrhythmias, seizures, loss of muscle control, and death.

^aThere are data to support that fixed-dose rasburicase is very effective in adult patients.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

NHODG-B
1 OF 4



SUPPORTIVE CARE FOR B-CELL LYMPHOMAS

For other immunosuppressive situations, [see NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections](#).

Monoclonal Antibody Therapy and Viral Reactivation

Anti-CD20 Antibody Therapy

Hepatitis B virus (HBV):

- Hepatitis B surface antigen (HBsAg) and Hepatitis B core antibody (HBcAb) testing for all patients receiving anti-CD20 antibody therapy
 - ▶ Quantitative hepatitis B viral load by PCR and surface antibody only if one of the screening tests is positive
- Note: Patients receiving IV immunoglobulin (IVIG) may be HBcAb-positive as a consequence of IVIG therapy.
- Prophylactic antiviral therapy with entecavir is recommended for any patient who is HBsAg-positive and receiving anti-lymphoma therapy. If there is active disease (PCR+), it is considered treatment/management and not prophylactic therapy. In cases of HBcAb positivity, prophylactic antiviral therapy is preferred; however, if there is a concurrent high-level hepatitis B surface antibody, these patients may be monitored with serial hepatitis B viral load.
 - ▶ Entecavir is preferred based on Huang YH, et al. J Clin Oncol 2013;31:2765-2772; Huang H, et al. JAMA 2014;312:2521-2530.
 - ▶ Avoid lamivudine due to risks of resistance development.
 - ▶ Other antivirals including adefovir, telbivudine, and tenofovir are proven active treatments and are acceptable alternatives.
 - ▶ Monitor hepatitis B viral load with PCR monthly through treatment and every 3 months thereafter.
 - ◊ If viral load is consistently undetectable, treatment is considered prophylactic.
 - ◊ If viral load fails to drop or previously undetectable PCR becomes positive, consult hepatologist and discontinue anti-CD20 antibody therapy.
 - ▶ Maintain prophylaxis up to 12 mo after oncologic treatment ends
 - ◊ Consult with hepatologist for duration of therapy in patient with active HBV.

Hepatitis C virus (HCV):

- New evidence from large epidemiology studies, molecular biology research, and clinical observation supports an association of HCV and B-cell NHL. Recently approved direct-acting antiviral agents (DAA) for chronic carriers of HCV with genotype 1 demonstrated a high rate of sustained viral responses.
 - ▶ Low-grade B-cell NHL
 - ◊ According to the American Association for the Study of Liver Diseases, combined therapy with DAA should be considered in asymptomatic patients with HCV genotype 1 since this therapy can result in regression of lymphoma.
 - ▶ Aggressive B-cell NHL
 - ◊ Patients should be initially treated with chemoimmunotherapy regimens according to NCCN Guidelines for NHL.
 - ◊ Liver functional tests and serum HCV RNA levels should be closely monitored during and after chemoimmunotherapy for development of hepatotoxicity.
 - ◊ Antiviral therapy should be considered in patients in complete remission after completion of lymphoma therapy.

Anti-CD20 Antibody Therapy and Brentuximab Vedotin

Progressive multifocal leukoencephalopathy (PML):

- Caused by the JC virus and is usually fatal.
 - ▶ Diagnosis made by PCR of CSF and in some cases brain biopsy.
- No known effective treatment.
- Clinical indications may include changes in behavior such as confusion, dizziness or loss of balance, difficulty talking or walking, and vision problems.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

NHODG-B
2 OF 4



SUPPORTIVE CARE FOR B-CELL LYMPHOMAS^b

Rare Complications of Monoclonal Antibody Therapy

- Rare complications such as mucocutaneous reactions including paraneoplastic pemphigus, Stevens-Johnson syndrome, lichenoid dermatitis, vesiculobullous dermatitis, and toxic epidermal necrolysis can occur. Expert consultation with dermatology is recommended.
- Re-challenge with the same monoclonal antibody is not recommended in patients experiencing rare complications to chosen anti-CD20 monoclonal antibody (rituximab, obinutuzumab, or ofatumumab). An alternative anti-CD20 monoclonal antibody (obinutuzumab or ofatumumab) could be used for patients with intolerance to rituximab, regardless of histology. It is unclear that the use of alternative anti-CD20 monoclonal antibody poses the same risk of recurrence.

Rituximab Rapid Infusion

- If no infusion reactions were experienced with prior cycle of rituximab, a rapid infusion over 90 minutes can be used.

Rituximab-Related Neutropenia

- Usually delayed in onset weeks to months after last exposure
- Occurs in up to 10% of patients
- Can be severe, but usually not presenting with infections
- Can be initially observed for spontaneous recovery, but if prolonged a short course of G-CSF is indicated
- Intravenous immunoglobulin (IVIG) has been anecdotally successful in patients refractory to G-CSF

Renal Dysfunction Associated with Methotrexate

- Consider use of glucarpidase if significant renal dysfunction and methotrexate levels are >10 microM beyond 42 to 48 hours. Leucovorin remains a component in the treatment of methotrexate toxicity and should be continued for at least 2 days following glucarpidase administration. However, be aware that leucovorin is a substrate for glucarpidase, and therefore should not be administered within two hours prior to or following glucarpidase.

Immunizations

- [See NCCN Guidelines for Survivorship - General Principles of Immunizations](#)

Prophylaxis for Pneumocystis Jiroveci Pneumonia (PJP) and Varicella-Zoster Virus (VZV)

- [See NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections](#)

Hypogammaglobulinemia

- Patients receiving anti-CD20 monoclonal antibody and anti-CD19-directed CAR T-cell therapy may experience hypogammaglobulinemia. Patients with recurrent infections may benefit from IVIG replacement.

^bSupportive care measures specific to rituximab include both rituximab as well as rituximab biosimilars.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

NHODG-B
3 OF 4

**SUPPORTIVE CARE FOR B-CELL LYMPHOMAS****Bone Health: Recommendations for Patients Who Have Received Steroid-Containing Regimens^{a,b,c,d}**
(in addition to standard recommendations for screening)**• Evaluation**

- ▶ **Vitamin D, 25-OH level**
- ▶ **Post-treatment bone mineral density (BMD) evaluation (1 year following therapy)**
 - ◊ **Greatest risk in women with chemotherapy-induced premature menopause**
 - If osteopenic (T score between -1.1 and -2.4):
 - ◊ **Use Fracture Risk Assessment Tool (FRAX) to determine if drug therapy is necessary (<https://www.sheffield.ac.uk/FRAX/>)**
 - **20% risk for any major osteoporotic fracture or 3% risk for hip fracture are the thresholds where drug therapy is recommended**
 - If T-score -2 to -2.4 (at any site) or ongoing glucocorticoid exposure repeat BMD every 1–2 years, as long as risk factors persist.^e
 - If T-score -1.5 to -1.9 (at any site) with no risk factors, repeat BMD in 5 years^e

• Therapy

- ▶ **If vitamin D 25-OH is deficient, then replete**
 - ◊ **In lymphoma patients with current elevations in 1,25-dihydroxyvitamin D, deficient 25(OH)D levels should not be aggressively replaced.**
- ▶ **Calcium intake from food (plus supplements if necessary) should be commensurate with National Academy of Medicine recommendations except in cases of lymphoma-induced hypercalciuria/hypercalcemia due to excessive 1,25-dihydroxyvitamin D production.**
 - ◊ **In patients receiving corticosteroid-containing chemotherapy regimens, adequate calcium intake is of paramount importance since corticosteroids block calcium absorption and increase fracture risk.^f**
- ▶ **Patients with osteoporotic bone mineral density, with a history of hip or vertebral fractures, or with asymptomatic vertebral compression deformity (as seen on CT scan or other imaging) should be started on therapy as per National Osteoporosis Foundation (NOF) guidelines; referral to an endocrinologist with expertise in bone health is recommended.**
 - ◊ **In appropriate women with premature menopause, hormone replacement therapy (HRT) up until the expected time of natural menopause, or raloxifene could be considered.**
 - ◊ **Bisphosphonates should be used as first-line pharmacologic treatment for osteoporosis.**
 - ◊ **In patients who cannot tolerate or whose symptoms do not improve with bisphosphonate therapy, denosumab is an effective alternative medication to prevent osteoporotic fractures.**
 - **Teriparatide is contraindicated in patients with a history of radiotherapy; also, theoretical concerns in patients with a recent history of cancer exist.**

^aCrandall CJ, Newberry SJ, Diamant A, et al. Comparative effectiveness of pharmacologic treatments to prevent fractures: an updated systematic review. *Ann Intern Med* 2014;161:711-723.

^bMacLean C, Newberry S, Maglione M, et al. Systematic review: comparative effectiveness of treatments to prevent fractures in men and women with low bone density or osteoporosis. *Ann Intern Med* 2008;148:197-213.

^cCummings SR, San Martin J, McClung MR, et al. Denosumab for prevention of fractures in postmenopausal women with osteoporosis. *N Engl J Med* 2009;361:756-765. [published correction appears in *N Engl J Med*. 2009; 361(19):1914].

^dPaccou L, Merlusca I, Henry-Desailly A, et al. Alterations in bone mineral density and bone turnover markers in newly diagnosed adults with lymphoma receiving chemotherapy: a 1-year prospective pilot study. *Ann Oncol* 2014; 25:481-486.

^ehttps://www.uptodate.com/contents/screening-for-osteoporosis?source=see_link.

^fVan Staa TP, Leufkens HG, Abenhaim L, et al. Use of oral corticosteroids and risk of fractures. *J Bone Miner Res* 2005;20:1487-1494.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

LUGANO RESPONSE CRITERIA FOR NON-HODGKIN LYMPHOMA

PET should be done with contrast-enhanced diagnostic CT and can be done simultaneously or at separate procedures.

Response	Site	PET-CT (Metabolic response)	CT (Radiologic response) ^d
Complete response	Lymph nodes and extralymphatic sites	Score 1, 2, 3 ^a with or without a residual mass on 5-point scale (5-PS) ^{b,c}	All of the following: Target nodes/nodal masses must regress to ≤1.5 cm in longest transverse diameter of a lesion (LDi) No extralymphatic sites of disease
	Non-measured lesion	Not applicable	Absent
	Organ enlargement	Not applicable	Regress to normal
	New lesions	None	None
	Bone marrow	No evidence of FDG-avid disease in marrow	Normal by morphology; if indeterminate and flow cytometry IHC negative
Partial response	Lymph nodes and extralymphatic sites	Score 4 or 5 ^b with reduced uptake compared with baseline. No new progressive lesions. At interim these findings suggest responding disease. At end of treatment these findings may indicate residual disease.	All of the following: ≥50% decrease in SPD of up to 6 target measurable nodes and extranodal sites When a lesion is too small to measure on CT, assign 5 mm x 5 mm as the default value. When no longer visible, 0x0 mm For a node >5 mm x 5 mm, but smaller than normal, use actual measurement for calculation
	Non-measured lesion	Not applicable	Absent/normal, regressed, but no increase
	Organ enlargement	Not applicable	Spleen must have regressed by >50% in length beyond normal
	New lesions	None	None
	Bone marrow	Residual uptake higher than uptake in normal marrow but reduced compared with baseline (diffuse uptake compatible with reactive changes from chemotherapy allowed). If there are persistent focal changes in the marrow in the content of a nodal response, consider further evaluation with biopsy, or an interval scan.	Not applicable

Reprinted with permission. © 2014 American Society of Clinical Oncology. All rights reserved. Cheson B, Fisher R, Barrington S, et al. Recommendations for initial evaluation, staging and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. J Clin Oncol 2014;32:3059-3068.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

NHODG-C
1 OF 3



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

LUGANO RESPONSE CRITERIA FOR NON-HODGKIN LYMPHOMA

PET should be done with contrast-enhanced diagnostic CT and can be done simultaneously or at separate procedures.

Response	Site	PET-CT (Metabolic response)	CT (Radiologic response) ^d
No response or stable disease	Target nodes/nodal masses, extranodal lesions	Score 4 or 5 ^b with no significant change in FDG uptake from baseline at interim or end of treatment. No new or progressive lesions.	<50% decrease from baseline in SPD of up to 6 dominant, measureable nodes and extranodal sites; no criteria for progressive disease are met
	Non-measured lesion	Not applicable	No increase consistent with progression
	Organ enlargement	Not applicable	No increase consistent with progression
	New lesions	None	None
	Bone marrow	No change from baseline	Not applicable
Progressive disease	Individual target nodes/nodal masses, extranodal lesions	Score 4 or 5 ^b with an increase in intensity of uptake from baseline and/or New FDG-avid foci consistent with lymphoma at interim or end-of-treatment assessment ^e	Requires at least one of the following PPD progression: An individual node/lesion must be abnormal with: LDi >1.5 cm and Increase by ≥50% from PPD nadir and An increase in LDi or SDi from nadir 0.5 cm for lesions ≤2 cm 1.0 cm for lesions >2 cm In the setting of splenomegaly, the splenic length must increase by >50% of the extent of its prior increase beyond baseline. If no prior splenomegaly, must increase by at least 2 cm from baseline New or recurrent splenomegaly
	Non-measured lesion	None	New or clear progression of preexisting nonmeasured lesions
	New lesions	New FDG-avid foci consistent with lymphoma rather than another etiology (eg, infection, inflammation). If uncertain regarding etiology of new lesions, biopsy or interval scan may be considered ^e	Regrowth of previously resolved lesions A new node >1.5 cm in any axis A new extranodal site >1.0 cm in any axis; if <1 cm in any axis, its presence must be unequivocal and must be attributable to lymphoma Assessable disease of any size unequivocally attributable to lymphoma
	Bone marrow	New or recurrent FDG-avid foci	New or recurrent involvement

Reprinted with permission. © 2014 American Society of Clinical Oncology. All rights reserved. Cheson B, Fisher R, Barrington S, et al. Recommendations for initial evaluation, staging and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. J Clin Oncol 2014;32:3059-3068.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

NHODG-C
2 OF 3



LUGANO RESPONSE CRITERIA FOR NON-HODGKIN LYMPHOMA

Footnotes

^aScore 3 in many patients indicates a good prognosis with standard treatment, especially if at the time of an interim scan. However, in trials involving PET where de-escalation is investigated, it may be preferable to consider score 3 as an inadequate response (to avoid under-treatment).

^bSee PET Five-Point Scale (5-PS).

^cIt is recognized that in Waldeyer's ring or extranodal sites with high physiological uptake or with activation within spleen or marrow, e.g. with chemotherapy or myeloid colony stimulating factors, uptake may be greater than normal mediastinum and/or liver. In this circumstance, CMR may be inferred if uptake at sites of initial involvement is no greater than surrounding normal tissue even if the tissue has high physiological uptake.

^dFDG-avid lymphomas should have response assessed by PET-CT. Diseases that can typically be followed with CT alone include CLL/SLL and marginal zone lymphomas.

^eFalse-positive PET scans may be observed related to infectious or inflammatory conditions. Biopsy of affected sites remains the gold standard for confirming new or persistent disease at end of therapy.

PET Five-Point Scale (5-PS)

- 1 No uptake above background
- 2 Uptake \leq mediastinum
- 3 Uptake $>$ mediastinum but \leq liver
- 4 Uptake moderately $>$ liver
- 5 Uptake markedly higher than liver and/or new lesions
- X New areas of uptake unlikely to be related to lymphoma

SPD – Sum of the product of the perpendicular diameters for multiple lesions

LDi – Longest transverse diameter of a lesion

SDi – Shortest axis perpendicular to the LDi

PPD – Cross product of the LDi and perpendicular diameter

Measured dominant lesions – Up to 6 of the largest dominant nodes, nodal masses and extranodal lesions selected to be clearly measurable in 2 diameters. Nodes should preferably be from disparate regions of the body, and should include, where applicable, mediastinal and retroperitoneal areas. Non-nodal lesions include those in solid organs, e.g., liver, spleen, kidneys, lungs, etc, gastrointestinal involvement, cutaneous lesions of those noted on palpation.

Non-measured lesions – Any disease not selected as measured, dominant disease and truly assessable disease should be considered not measured. These sites include any nodes, nodal masses, and extranodal sites not selected as dominant, measurable or which do not meet the requirements for measurability, but are still considered abnormal. As well as truly assessable disease which is any site of suspected disease that would be difficult to follow quantitatively with measurement, including pleural effusions, ascites, bone lesions, leptomeningeal disease, abdominal masses and other lesions that cannot be confirmed and followed by imaging.

Reprinted with permission. © 2014 American Society of Clinical Oncology. All rights reserved. Cheson B, Fisher R, Barrington S, et al. Recommendations for initial evaluation, staging and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. J Clin Oncol 2014;32:3059-3068.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF RADIATION THERAPY

General Principles^a

- Treatment with photons, electrons, or protons is appropriate; selection depends upon clinical scenario.
- Advanced radiation therapy technologies such as IMRT/VMAT, proton therapy, breath-hold or respiratory gating, and/or image-guided therapy may offer significant and clinically relevant advantages in specific instances to spare organs at risk (OAR) such as the heart (including coronary arteries and valves), lungs, kidneys, liver, spinal cord, esophagus, bone marrow, breasts, stomach, muscle/soft tissue, and salivary glands and decrease the risk for late, normal tissue toxicity while still achieving the primary goal of local tumor control.
- Reducing dose to normal tissues reduces the risk of late complications. Achieving highly conformal dose distributions is especially important for patients who are being treated with curative intent or who have long life expectancies following therapy.
- For mediastinal and abdominal lymphoma, respiratory motion management such as gating or breath-hold techniques may be advantageous. Breath-hold techniques have been shown to decrease incidental dose to the heart and lungs in many disease presentations. Similarly, for abdominal lymphomas, reduction in radiation exposures to liver and kidneys may be achieved by motion management techniques.
- Since the advantages of these techniques include tightly conformal doses and steep gradients next to normal tissues, target definition and treatment delivery verification require careful monitoring to avoid the risk of tumor geographic miss and subsequent decrease in tumor control. Image guidance may be required to provide this assurance.
- Randomized studies to test these concepts are unlikely to be done since these techniques are designed to decrease late effects, which take 10+ years to evolve. In light of that, the modalities and techniques which best reduce the doses to the OAR in a clinically meaningful way without compromising target coverage should be considered.

^aSee references on [NHODG-D 4 of 4](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

NHODG-D
1 OF 4



PRINCIPLES OF RADIATION THERAPY^a

Volumes

- **ISRT for nodal disease (see Illidge T, et al. Int J Radiat Oncol Biol Phys 2014;89:49-58)**
 - ▶ **ISRT is the recommended approach for volume definition and treatment planning for NHL. Planning for ISRT requires CT-based treatment planning and incorporates volume determinations including GTV, CTV, and PTV. Incorporating other imaging tests such as PET and MRI often enhances treatment volume determination.**
 - ▶ **The pre-chemotherapy or pre-biopsy gross tumor volume (GTV) provides the basis for determining the clinical target volume (CTV). Concerns for questionable subclinical disease and uncertainties in original imaging accuracy or localization may lead to expansion of the CTV and are determined individually using clinical judgment. Further, adjacent uninvolved organs (eg, lungs, bone, muscle, kidney) are excluded from the CTV when disease regresses following chemotherapy.**
 - ▶ **For early-stage indolent NHL treated with RT alone, larger treatment volumes should be considered to encompass potential microscopic disease in adjacent lymph nodes or the immediate vicinity. For example, the CTV definition for treating follicular lymphoma with radiation therapy alone will be greater than that employed for DLBCL with similar disease distribution, as the latter is treated with combined modality therapy.**
 - ▶ **Motion of the target caused by respiration as determined by 4D-CT or fluoroscopy (internal target volume, ITV) should also influence the final CTV.**
 - ▶ **The planning treatment volume (PTV) is an additional expansion of the CTV that accounts only for setup variations (see ICRU definitions).**
 - ▶ **OARs should be outlined for dose-volume analysis and optimizing treatment planning decisions.**
 - ▶ **The treatment plan can be designed with conventional, 3-D conformal, IMRT/VMAT, or proton therapy techniques using clinical treatment planning considerations of target coverage and normal tissue avoidance.**
- **ISRT for extranodal disease (see Yahalom J, et al. Int J Radiat Oncol Biol Phys 2015;92:11-31)**
 - ▶ **Similar principles as for ISRT nodal sites (see above).**
 - ▶ **For most organs and particularly for indolent disease, the CTV consists of the entire affected organ (eg, stomach, salivary gland, thyroid). For other organs, including orbit, breast, lung, bone, localized skin, and in some cases when RT is consolidation after chemotherapy, partial organ RT may be appropriate.**
 - ▶ **For most NHL subtypes, uninvolved lymph nodes should not be targeted.**

^aSee references on [NHODG-D 4 of 4](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**NHODG-D
2 OF 4**



PRINCIPLES OF RADIATION THERAPY^a

General Dose Guidelines:

- **Definitive treatment (1.5–2.0 Gy daily fractions)**

- ▶ **Follicular lymphoma: 24–30 Gy**

- ▶ **Marginal zone lymphoma:**

- ◊ **24–30 Gy**

- ◊ **Gastric MALT lymphoma most commonly treated to 30 Gy in 20 fractions (1.5 Gy/fx) to minimize acute GI toxicity; however, 24 Gy is appropriate as well**

- ▶ **Early-stage mantle cell lymphoma: 24–36 Gy**

- ▶ **DLBCL/HGBCL/PMBL/Grey zone lymphoma**

- ◊ **Consolidation after chemotherapy**

- **CR - 30–36 Gy**

- **PR - 36–50 Gy**

- **Refractory - 40–55 Gy**

- ◊ **RT as primary treatment: 40–55 Gy**

- ◊ **In combination with stem cell transplantation: 20–36 Gy, depending on sites of disease and prior RT exposure (Ng A, et al. Int J Radiat Oncol Biol Phys 2018;100:652-669)**

- **Palliative RT**

- ▶ **FL/MZL/MCL/SLL: 2 Gy X 2 fractions or 4 Gy X 1 fractions (which may be repeated as needed); doses up to 30 Gy may be appropriate in select circumstances**

- ▶ **DLBCL/HGBCL/PMBL/Grey zone lymphoma and Burkitt lymphoma: (higher doses/fraction typically appropriate)**

- ◊ **20–30 Gy in 5–10 fractions. Standard hypofractionated palliative treatment schedules such as 20 Gy in 5 and 30 Gy in 10 are appropriate depending upon clinical scenario.**

- ◊ **AIDS-related B-cell lymphomas and PTLT: Treated based on underlying histologic subtype and treatment intent (curative vs. palliative)**

^aSee references on [NHODG-D 4 of 4](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF RADIATION THERAPY REFERENCES

- ¹Andolino DL, Hoene T, Xiao L, Buchsbaum J, Chang AL. Dosimetric comparison of involved-field three-dimensional conformal photon radiotherapy and breast-sparing proton therapy for the treatment of Hodgkin's lymphoma in female pediatric patients. *Int J Radiat Oncol Biol Phys* 2011;81:e667-671.
- ²Aznar MC, Maraldo MV, Schut DA, et al. Minimizing late effects for patients with mediastinal Hodgkin lymphoma: deep inspiration breath-hold, IMRT, or both? *Int J Radiat Oncol Biol Phys* 2015;92:169-174.
- ³Besson N, Pernin V, Zefkili S, Kirova YM. Evolution of radiation techniques in the treatment of mediastinal lymphoma: from 3D conformal radiotherapy (3DCRT) to intensity-modulated RT (IMRT) using helical tomotherapy (HT): a single-centre experience and review of the literature. *Br J Radiol* 2016;89:20150409.
- ⁴Boda-Heggemann J, Knopf AC, Simeonova-Chergou A, et al. Deep inspiration breath hold-based radiation therapy: a clinical review. *Int J Radiat Oncol Biol Phys* 2016;94:478-492.
- ⁵Claude L, Malet C, Pommier P, Thiesse P, Chabaud S, Carrie C. Active breathing control for Hodgkin's disease in childhood and adolescence: feasibility, advantages, and limits. *Int J Radiat Oncol Biol Phys* 2007;67:1470-1475.
- ⁶Filippi AR, Ragona R, Piva C, et al. Optimized volumetric modulated arc therapy versus 3D-CRT for early stage mediastinal Hodgkin lymphoma without axillary involvement: a comparison of second cancers and heart disease risk. *Int J Radiat Oncol Biol Phys* 2015;92:161-168.
- ⁷Ho CK, Flampouri S, Hoppe BS. Proton therapy in the management of lymphoma. *Cancer J* 2014;20:387-392.
- ⁸Hoppe BS, Flampouri S, Zaiden R, et al. Involved-node proton therapy in combined modality therapy for Hodgkin lymphoma: results of a phase 2 study. *Int J Radiat Oncol Biol Phys* 2014;89:1053-1059.
- ⁹Hoppe BS, Hill-Kayser CE, Tseng YD, et al. Consolidative proton therapy after chemotherapy for patients with Hodgkin lymphoma. *Ann Oncol* 2017;28:2179-2184.
- ¹⁰Horn S, Fournier-Bidoz N, Pernin V, et al. Comparison of passive-beam proton therapy, helical tomotherapy and 3D conformal radiation therapy in Hodgkin's lymphoma female patients receiving involved-field or involved site radiation therapy. *Cancer Radiother* 2016;20:98-103.
- ¹¹Nanda R, Flampouri S, Mendenhall NP, et al. Pulmonary toxicity following proton therapy for thoracic lymphoma. *Int J Radiat Oncol Biol Phys* 2017;99:494-497.
- ¹²Petersen PM, Aznar MC, Berthelsen AK, et al. Prospective phase II trial of image-guided radiotherapy in Hodgkin lymphoma: benefit of deep inspiration breath-hold. *Acta Oncol* 2015;54:60-66.
- ¹³Pinnix CC, Smith GL, Milgrom S, et al. Predictors of radiation pneumonitis in patients receiving intensity modulated radiation therapy for Hodgkin and non-Hodgkin lymphoma. *Int J Radiat Oncol Biol Phys* 2015;92:175-182.
- ¹⁴Tseng YD, Cutter DJ, Plastaras JP, et al. Evidence-based review on the use of proton therapy in lymphoma from the Particle Therapy Cooperative Group (PTCOG) Lymphoma Subcommittee. *Int J Radiat Oncol Biol Phys* 2017;99:825-842.
- ¹⁵Voong KR, McSpadden K, Pinnix CC, et al. Dosimetric advantages of a "butterfly" technique for intensity-modulated radiation therapy for young female patients with mediastinal Hodgkin's lymphoma. *Radiat Oncol* 2014;9:94.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



SPECIAL CONSIDERATIONS FOR THE USE OF SMALL-MOLECULE INHIBITORS¹

Acalabrutinib

• Dosage

- ▶ The recommended dose of acalabrutinib is 100 mg PO approximately every 12 hours

• Toxicity

- ▶ Grade ≥3 bleeding events were observed in 2% of patients on acalabrutinib. The mechanism is not well understood. Acalabrutinib may increase the risk of hemorrhage in patients receiving anti-platelet or anticoagulant therapies. The phase 2 ACE-LY-004 study excluded patients on concomitant warfarin or equivalent vitamin K antagonists. Patients should be monitored for signs of bleeding. Consider the benefit-risk of withholding acalabrutinib for 3–7 days pre-and post-surgery depending on the type of surgery and risk of bleeding.
- ▶ Atrial fibrillation and flutter of any grade was reported in 3% of patients and atrial fibrillation grade 3 was reported in 1% of patients. Monitor for atrial fibrillation and flutter and manage as appropriate.

Copanlisib

• Dosage

- ▶ The recommended dose of copanlisib is 60 mg administered as a 1-hour IV infusion on Days 1, 8, and 15 of a 28-day treatment cycle on an intermittent schedule (3 weeks on and 1 week off).

• Toxicity

- ▶ Infection: Monitor patients for signs and symptoms of infection prior to and during treatment. Copanlisib should be withheld for grade ≥ infection until resolution.
- ▶ PJP: Consider PJP prophylaxis for patients at risk before initiating copanlisib. Copanlisib should be withheld in patients with suspected PJP infection of any grade. If confirmed, treat infection until resolution, then resume copanlisib at previous dose with concomitant PJP prophylaxis.
- ▶ Neutropenia: Monitor blood counts at least weekly during treatment with copanlisib. Interrupt copanlisib, reduce dose, or discontinue copanlisib depending on the severity and persistence of neutropenia.
- ▶ Hyperglycemia: Patients with diabetes mellitus should only be treated with copanlisib following adequate glucose control and should be monitored closely. Interrupt copanlisib, reduce dose, or discontinue copanlisib depending on the severity and persistence of hyperglycemia.
- ▶ Hypertension: Optimal blood pressure control should be achieved before starting each copanlisib infusion. Monitor blood pressure pre- and post-infusion. Interrupt copanlisib, reduce dose, or discontinue copanlisib depending on the severity and persistence of hypertension.

¹Please refer to package insert for full prescribing information and monitoring for adverse reactions, available at www.fda.gov.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

NHODG-E
1 OF 4

**SPECIAL CONSIDERATIONS FOR THE USE OF SMALL-MOLECULE INHIBITORS¹****Duvelisib****• Dosage**

- ▶ The recommended dose of duvelisib is 25 mg PO twice daily, per prescribing recommendations.

• Lymphocytosis

- ▶ Upon initiation of duvelisib, transient increase in absolute lymphocyte count is expected in most patients, which does not signify disease progression. This onset of isolated lymphocytosis occurs during the first few weeks of duvelisib therapy and may persist for several weeks on treatment.

• Toxicity

- ▶ **Hepatotoxicity:** Monitor hepatic function prior to and during treatment. Interrupt if ALT/AST >5 x ULN (upper limit of normal) and when resolved resume at the same dose (25 mg twice daily) for first occurrence or at a reduced dose (15 mg twice daily) for subsequent occurrence. Discontinue duvelisib if ALT/AST > 20 x ULN.
- ▶ **Diarrhea or colitis:** Monitor for the development of severe diarrhea or colitis. Initiate supportive therapy with antidiarrheal agents as appropriate. In case of severe diarrhea or colitis, interrupt duvelisib until resolution and then resume at a reduced dose (15 mg twice daily) or discontinue duvelisib. Severe diarrhea and colitis can be managed with enteric acting steroids (eg, budesonide) or systemic steroids.
- ▶ **Pneumonitis without suspected infectious cause:** Interrupt duvelisib and treat with systemic steroid therapy for grade 2. If pneumonitis recovers to grade 0 or 1, duvelisib may be resumed at reduced dose (15 mg twice daily). Discontinue duvelisib if non-infectious pneumonitis recurs or patient does not respond to steroid therapy or for severe (grade 3) or life-threatening pneumonitis.
- ▶ **Cutaneous reactions:** Monitor closely and initiate supportive care with emollients, antihistamines (for pruritus), or topical steroids. In case of severe cutaneous reactions, interrupt duvelisib until resolution and initiate supportive care with emollients, antihistamines (for pruritus), or topical steroids. Resume at a reduced dose (15 mg twice daily). If severe cutaneous reaction does not improve, worsens, or recurs, discontinue duvelisib.
- ▶ **Infections:** PJP prophylaxis with sulfamethoxazole/trimethoprim or equivalent.
- ▶ **CMV reactivation:** Consider prophylactic antivirals to prevent CMV infection including CMV reactivation.

Ibrutinib**• Dosage**

- ▶ **MCL, DLBCL:** The recommended dose of ibrutinib is 560 mg PO daily, continuous and should be continued until time of progression.

• Lymphocytosis

- ▶ **MCL:** Upon initiation of ibrutinib, transient increase in absolute lymphocyte counts occurred in 33% of patients. The onset of isolated lymphocytosis occurs during the first few weeks of ibrutinib therapy and resolves by a median of 8 weeks.

• Toxicity

- ▶ **Grade >2 bleeding events** were observed in 6% of patients on ibrutinib; the mechanism is not well understood. Consider the benefit-risk of ibrutinib in patients requiring anti-platelet or anticoagulant therapies. Clinical trials excluded patients on concurrent warfarin. Ibrutinib should be held 3 days before and after a minor surgical procedure and 7 days before and after a major surgical procedure. Ibrutinib should not be given concomitantly with warfarin.
- ▶ **New-onset atrial fibrillation** was reported in 6%–9%, associated with ibrutinib administration.
 - ◇ Consider non-warfarin anticoagulation.
 - ◇ Monitor carefully.
 - ◇ Consider switching to alternate therapy.
 - ◇ Patients with recurrent atrial fibrillation that is not medically controllable should be changed to an alternative agent.
- ▶ **Hypertension** associated with ibrutinib has been uncommonly reported as a basis for discontinuation and should be managed with anti-hypertensives as appropriate. Ibrutinib should only be discontinued for uncontrollable hypertension.

¹Please refer to package insert for full prescribing information and monitoring for adverse reactions, available at www.fda.gov.**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.[Continued](#)**NHODG-E**
2 OF 4



SPECIAL CONSIDERATIONS FOR THE USE OF SMALL-MOLECULE INHIBITORS¹

Idelalisib

- **Dosage**
 - The recommended dose of idelalisib is 150 mg PO twice daily, per prescribing recommendations.
- **Toxicity**
 - **Hepatotoxicity:** Monitor hepatic function prior to and during treatment. Interrupt (if ALT/AST > 5 x ULN) and when resolved may resume at a reduced dose (100 mg PO twice daily).
 - **Diarrhea or colitis:** Monitor for the development of severe diarrhea or colitis. Interrupt until resolution and then reduce or discontinue idelalisib. Severe diarrhea and colitis can be managed with systemic or nonabsorbable steroids.
 - **Pneumonitis:** Monitor for pulmonary symptoms and bilateral interstitial infiltrates. Discontinue idelalisib.
 - **Intestinal perforation:** Discontinue idelalisib if intestinal perforation is suspected.
- **Infections:** PJP prophylaxis with sulfamethoxazole/trimethoprim or equivalent.
- **CMV:** Monitor per institutional guidelines or consult with infectious disease specialist.

Venetoclax

- **Dosage**
 - The recommended single-agent dose for venetoclax is 400 mg (higher dose for insufficient response).
 - Initiate venetoclax at 20 mg for one week and gradually escalate to target dose of 400 mg PO daily over 5 weeks to reduce the risk of TLS.
 - **TLS:** See Tumor Lysis Syndrome ([NHODG-B 1 of 4](#))
- **Toxicity**
 - Consider the use of neutrophil growth factors for neutropenia according to standard guidelines.

Zanubrutinib

- **Dosage**
 - The recommended dose of zanubrutinib is 160 mg orally twice daily or 320 mg orally once daily, per prescribing recommendations.
- **Toxicity**
 - **Hemorrhage:** Grade 3 or higher bleeding events including intracranial and gastrointestinal hemorrhage, hematuria, and hemothorax have been reported in 2% of patients treated with zanubrutinib monotherapy. Bleeding events of any grade, including purpura and petechiae, occurred in 50% of patients treated with zanubrutinib monotherapy. Consider the benefit-risk of withholding zanubrutinib for 3–7 days pre- and post-surgery depending upon the type of surgery and the risk of bleeding.
 - **Cytopenias:** Grade 3 or 4 cytopenias, including neutropenia (27%), thrombocytopenia (10%), and anemia (8%) based on laboratory measurements, were reported in patients treated with zanubrutinib monotherapy.
 - **Cardiac arrhythmias:** Atrial fibrillation and atrial flutter have occurred in 2% of patients treated with zanubrutinib monotherapy. Patients with cardiac risk factors, hypertension, and acute infections may be at increased risk. Grade 3 or higher events were reported in 0.6% of patients treated with zanubrutinib monotherapy.
- **Infections:** Grade 3 or higher infections occurred in 23% of patients treated with zanubrutinib monotherapy. Consider prophylaxis for herpes simplex virus, PJP, and other infections according to standard of care in patients who are at increased risk for infections.

[Continued](#)

¹Please refer to package insert for full prescribing information and monitoring for adverse reactions, available at www.fda.gov.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



SPECIAL CONSIDERATIONS FOR THE USE OF SMALL-MOLECULE INHIBITORS¹

Co-administration with CYP3A inhibitors and inducers

• Acalabrutinib

- Avoid concomitant use of strong CYP3A inhibitors or inducers.
- For strong CYP3A inhibitors used short-term, interrupt acalabrutinib during the duration of inhibitor use.
- For concomitant use with a moderate CYP3A inhibitor, reduce acalabrutinib dose to 100 mg once daily.
- If concomitant use with a strong CYP3A inducer cannot be avoided, increase acalabrutinib dose to 200 mg twice daily.

• Copanlisib

- Avoid concomitant use of strong CYP3A inhibitors or inducers.
- If concomitant use with strong CYP3A inhibitors cannot be avoided, reduce the copanlisib dose to 45 mg.

• Duvelisib

- Avoid concomitant use of strong CYP3A inducers.
- Patients taking concomitant strong CYP3A4 inhibitors should be monitored more closely for signs of duvelisib toxicity. Reduce dose to 15 mg twice daily when co-administered with strong CYP3A4 inhibitors.
- Monitor for signs of toxicities when coadministering with sensitive CYP3A substrates.

• Ibrutinib

- Avoid concomitant use of strong and moderate CYP3A inhibitors. Consider alternative agents with less CYP3A inhibition.
 - ◊ For strong CYP3A inhibitors used short-term (eg, antifungals and antibiotics for 7 days or less; eg, ketoconazole, itraconazole, voriconazole, posaconazole, clarithromycin, telithromycin), consider interrupting ibrutinib during the duration of inhibitor use. Avoid strong CYP3A inhibitors that are needed chronically.
 - ◊ If a moderate CYP3A inhibitor must be used, reduce ibrutinib dose to 140 mg.
 - ◊ Patients taking concomitant strong or moderate CYP3A4 inhibitors should be monitored more closely for signs of ibrutinib toxicity
- Avoid concomitant use of strong CYP3A inducers (eg, carbamazepine, rifampin, phenytoin, St. John's Wort). Consider alternative agents with less CYP3A induction.

• Idelalisib

- Avoid concomitant use of strong CYP3A inhibitors or inducers.
- Patients taking concomitant strong CYP3A4 inhibitors should be monitored more closely for signs of idelalisib toxicity.

• Venetoclax

- Avoid concomitant use of strong CYP3A inhibitors or inducers.

• Zanubrutinib

- Modify dose with strong CYP3A inhibitors. Interrupt dose as recommended for adverse reactions.
- Modify dose with moderate CYP3A inhibitors. Modify dose as recommended for adverse reactions.
- Avoid concomitant use with moderate or strong CYP3A inducers.

¹Please refer to package insert for full prescribing information and monitoring for adverse reactions, available at www.fda.gov.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



Classification

Table 1

WHO Classification of the Mature B-Cell, T-Cell, and NK-Cell Neoplasms (2017)

Mature B-Cell Neoplasms

- Chronic lymphocytic leukemia/small lymphocytic lymphoma
- Monoclonal B-cell lymphocytosis
- B-cell prolymphocytic leukemia
- Splenic marginal zone lymphoma
- Hairy cell leukemia
- *Splenic lymphoma/leukemia, unclassifiable**
 - ▶ *Splenic diffuse red pulp small B-cell lymphoma**
 - ▶ *Hairy cell leukemia-variant**
- Lymphoplasmacytic lymphoma
 - ▶ Waldenström macroglobulinemia
- Monoclonal gammopathy of undetermined significance (MGUS), IgM
- Mu heavy chain disease
- Gamma heavy chain disease
- Alpha heavy chain disease
- Monoclonal gammopathy of undetermined significance (MGUS), IgG/A
- Plasma cell myeloma
- Solitary plasmacytoma of bone
- Extramedullary plasmacytoma
- Monoclonal immunoglobulin deposition diseases
- Extranodal marginal zone lymphoma of mucosa-associated lymphoid tissue (MALT lymphoma)
- Nodal marginal zone lymphoma
 - ▶ *Pediatric nodal marginal zone lymphoma**
- Follicular lymphoma
 - ▶ In situ follicular neoplasia
 - ▶ Duodenal-type follicular lymphoma
- Pediatric-type follicular lymphoma
- *Large B-cell lymphoma with IRF4 rearrangement*
- Primary cutaneous follicle center lymphoma
- Mantle cell lymphoma
 - ▶ In situ mantle cell neoplasia
- Diffuse large B-cell lymphoma (DLBCL), NOS
 - ▶ Germinal center B-cell type
 - ▶ Activated B-cell type
- T-cell/histiocyte-rich large B-cell lymphoma
- Primary DLBCL of the central nervous system (CNS)
- Primary cutaneous DLBCL, leg type
- EBV-positive DLBCL, NOS
- *EBV-positive mucocutaneous ulcer**
- DLBCL associated with chronic inflammation
- Lymphomatoid granulomatosis
- Primary mediastinal (thymic) large B-cell lymphoma
- Intravascular large B-cell lymphoma
- ALK-positive large B-cell lymphoma
- Plasmablastic lymphoma
- Primary effusion lymphoma
- *HHV8-positive DLBCL, NOS**
- Burkitt lymphoma
- *Burkitt-like lymphoma with 11q aberration**
- High-grade B-cell lymphoma, with *MYC* and *BCL2* and/or *BCL6* rearrangements
- High-grade B-cell lymphoma, NOS
- B-cell lymphoma, unclassifiable, with features intermediate between DLBCL and classical Hodgkin lymphoma

*Provisional entities are listed in italics.

[Continued](#)

Swerdlow SH, CE, Harris NL, Jaffe ES, Pileri SA, Stein H, Thiele J, ed. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Revised 4th ed. Lyon: IARC; 2017.



Staging

Lugano Modification of Ann Arbor Staging System* (for primary nodal lymphomas)

<u>Stage</u>	<u>Involvement</u>	<u>Extranodal (E) status</u>
Limited		
Stage I	One node or a group of adjacent nodes	Single extranodal lesions without nodal involvement
Stage II	Two or more nodal groups on the same side of the diaphragm	Stage I or II by nodal extent with limited contiguous extranodal involvement
Stage II bulky**	II as above with “bulky” disease	Not applicable
Advanced		
Stage III	Nodes on both sides of the diaphragm Nodes above the diaphragm with spleen involvement	Not applicable
Stage IV	Additional non-contiguous extralymphatic involvement	Not applicable

*Extent of disease is determined by PET/CT for avid lymphomas, and CT for non-avid histologies.

Note: Tonsils, Waldeyer’s ring, and spleen are considered nodal tissue.

**Whether II bulky is treated as limited or advanced disease may be determined by histology and a number of prognostic factors.

Categorization of A versus B has been removed from the Lugano Modification of Ann Arbor Staging.

Reprinted with permission. © 2014 American Society of Clinical Oncology. All rights reserved. Cheson B, Fisher R, Barrington S, et al. Recommendations for initial evaluation, staging, and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. J Clin Oncol 2014;32:3059-3068.



NCCN Categories of Evidence and Consensus

Category 1	Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
Category 2A	Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
Category 2B	Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.
Category 3	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.

NCCN Categories of Preference

Preferred intervention	Interventions that are based on superior efficacy, safety, and evidence; and, when appropriate, affordability.
Other recommended intervention	Other interventions that may be somewhat less efficacious, more toxic, or based on less mature data; or significantly less affordable for similar outcomes.
Useful in certain circumstances	Other interventions that may be used for selected patient populations (defined with recommendation).

All recommendations are considered appropriate.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

This discussion corresponds to the NCCN Guidelines for B-Cell Lymphomas. Update is in progress. Last updated on 12/18/2019.

Discussion

Table of Contents

Overview	MS-2
Classification	MS-2
Diagnosis	MS-3
Workup	MS-3
Role of PET Scans	MS-4
Principles of Radiation Therapy	MS-5
Biosimilars	MS-6
Supportive Care	MS-7
Follicular Lymphoma	MS-24
Marginal Zone Lymphomas	MS-53
Mantle Cell Lymphoma	MS-76
Diffuse Large B-Cell Lymphoma	MS-96
High-Grade B-Cell Lymphomas with translocations of <i>MYC</i> and <i>BCL2</i> and/or <i>BCL6</i>	MS-113
Burkitt Lymphoma	MS-128
AIDS-Related B-Cell Lymphomas	MS-137
Post-Transplant Lymphoproliferative Disorders	MS-150



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Overview

Non-Hodgkin's lymphomas (NHL) are a heterogeneous group of lymphoproliferative disorders originating in B-lymphocytes, T-lymphocytes, or natural killer (NK) cells (NK/T-cell lymphomas are very rare). In 2019, an estimated 74,200 people will be diagnosed with NHL and there will be approximately 19,970 deaths due to the disease.¹ NHL is the seventh leading site of new cancer cases among men and women, accounting for 4% to 5% of new cancer cases and 3% to 4% of cancer-related deaths.¹ In prospectively collected data from the National Cancer Database, diffuse large B-cell lymphoma (DLBCL; 32%), chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL; 19%), follicular lymphoma (FL; 17%), marginal zone lymphoma (MZL; 8%), mantle cell lymphoma (MCL; 4%), and peripheral T-cell lymphoma not otherwise specified (PTCL-NOS; 2%) were the major subtypes of NHL diagnosed in the United States between 1998 and 2011.²

The incidence of NHL has increased dramatically between 1970 and 1995; the increase has moderated since the mid-1990s. This increase has been attributed partly to the human immunodeficiency virus (HIV) epidemic and the development of acquired immunodeficiency virus (AIDS)-related NHL. However, much of the increase in incidence has been observed in patients in their sixth and seventh decades; a large part of this increase incidence has paralleled a major decrease in mortality from other causes. The median age of individuals with NHL has risen in the last two decades.³ As a result, patients with NHL may also have significant comorbid conditions, which complicate treatment options.

The National Comprehensive Cancer Network (NCCN®) Guidelines (NCCN Guidelines®) were developed as a result of meetings convened by a multidisciplinary panel of NHL experts, with the aim to provide recommendations for diagnostic workup, treatment, and surveillance strategies for the most common subtypes of NHL, in addition to a general

discussion on the classification systems used in NHL and supportive care considerations.

The most common B-cell lymphoma subtypes that are covered in these NCCN Guidelines are listed below:

- ♦ Follicular lymphoma (FL)
- ♦ Marginal zone lymphomas (MZLs)
 - Gastric MALT lymphoma
 - Non gastric MALT lymphoma
 - Nodal MZL
 - Splenic MZL
- ♦ Mantle cell lymphoma (MCL)
- ♦ Diffuse large B-cell lymphoma (DLBCL)
- ♦ Burkitt lymphoma (BL)
- ♦ AIDS-related B-cell lymphomas
- ♦ Post-transplant lymphoproliferative disorders
- ♦ Castleman's disease

Classification

The 2001 World Health Organization (WHO) classification of hematopoietic and lymphoid neoplasms represented the first international consensus on classification of hematologic malignancies. After consideration of cell of origin (B, T, or NK), the classification subdivides lymphomas into those derived from precursor lymphocytes versus those derived from mature lymphocytes. The classification was further refined based on immunophenotype, genetic, and clinical features to aid in defining active treatment for specific subtypes of lymphoma.

Genetic features, detected by cytogenetics or fluorescence in situ hybridization (FISH), are increasingly important in defining specific NHL subtypes. In addition, detection of viruses, particularly Epstein-Barr virus (EBV), HHV8, and HTLV1, is often necessary to establish a specific



diagnosis. The WHO classification was revised in 2008 and 2017 to include new disease entities and better define some of the heterogeneous and ambiguous subtypes, based on the evolving genetic and molecular landscape of various subtypes of NHL.^{4,5}

Revisions to the 2008 and 2017 WHO classifications are discussed under the respective subtypes of B-cell lymphomas.

Diagnosis

An accurate pathologic diagnosis of the subtype is the most important first step in the management of B-cell lymphomas. The basic pathologic evaluation is the same for each subtype, although some further evaluation may be useful in certain circumstances to clarify a particular diagnosis; these are outlined in the pathologic evaluation of the individual Guidelines.

An incisional or excisional lymph node biopsy is recommended to establish the diagnosis of B-cell lymphomas. Fine-needle aspiration (FNA) biopsy alone is not generally suitable for the initial diagnosis of lymphoma.^{6,7} Recent studies have shown that the diagnostic accuracy of FNA improves significantly when it is used in combination with immunohistochemistry (IHC) and flow cytometry.⁸⁻¹⁰ A core needle biopsy is not optimal but can be used under certain circumstances when a clinical situation dictates that this is the only safe means of obtaining diagnostic tissue. In certain circumstances, when a lymph node is not easily accessible for excisional or incisional biopsy, a combination of core needle biopsy (multiple biopsies preferred) and FNA biopsies in conjunction with appropriate ancillary techniques (IHC, flow cytometry, karyotype and fluorescence in situ hybridization [FISH] for major translocations and molecular analysis for immunoglobulin heavy chain variable [*IGHV*] and T-cell receptor [*TCR*] gene rearrangements) may be sufficient for the differential diagnosis. Hematopathology review of all slides (with at least

one paraffin block representative of the tumor) is recommended. Rebiopsy should be done if consult material is nondiagnostic.

IHC is essential for the differentiation of various subtypes of NHL to establish the proper diagnosis. It can be performed by flow cytometry and/or IHC; the choice depends on the antigens as well as the expertise and resources available to the hematopathologist. In some cases flow cytometry and IHC are complementary diagnostic tools.¹¹ Cytogenetic or molecular genetic analysis may be necessary under certain circumstances to identify the specific chromosomal translocations that are characteristic of some NHL subtypes or to establish clonality.

The NCCN Guidelines panel developed a series of algorithms for the use of immunophenotyping in the diagnosis of mature lymphoid neoplasms. These algorithms were developed to provide guidance for surgical pathologists as well as an aid to the clinician in the interpretation of pathology reports, and they should be used in conjunction with clinical and pathologic correlation. See “*Use of Immunophenotyping/Genetic Testing in Differential Diagnosis of Mature B-Cell and NK/T-Cell Neoplasms*” in the algorithm.

Workup

Essential workup procedures include a complete physical exam with particular attention to node-bearing areas and the size of liver and spleen, symptoms present, performance status, and laboratory studies including complete blood count (CBC), serum lactate dehydrogenase (LDH), hepatitis B virus (HBV) testing (see below), comprehensive metabolic panel, and CT chest/abdominal/pelvic with oral and intravenous contrast (unless coexistent renal insufficiency). Multigated acquisition (MUGA) scan or echocardiograms are recommended when anthracyclines and anthracenedione-containing regimens are used.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

HBV reactivation (resulting in liver failure and death) has been reported in patients treated with anti-CD20 monoclonal antibody (MAB)-based regimens.¹² HBV carriers with lymphoid malignancies have a high risk of HBV reactivation and disease, especially those treated with anti-CD20 MAB-based regimens.¹³ The panel has included testing for hepatitis B surface antigen (HBsAg) and hepatitis B core antibody (HBcAb) as part of essential workup prior to initiation of treatment in all patients who will receive anti-CD20 MAB-based regimens. See *Hepatitis B virus Reactivation* in the *Supportive Care* section.

Large population-based or multicenter case-control studies have demonstrated a strong association between seropositivity for hepatitis C virus (HCV) and the development of B-cell lymphomas.¹⁴⁻²¹ The prevalence of HCV seropositivity was consistently increased among patients with DLBCL and MZL.^{14,15,19,20} HCV testing is needed in high-risk patients and in patients with splenic MZL. See *Hepatitis C Virus-associated B-Cell Lymphomas* in the *Supportive Care* section.

Optional workup procedures (depending on specific lymphoma type) include beta-2-microglobulin, CT or PET/CT scans, endoscopic ultrasound, head CT, or brain MRI and lumbar puncture to analyze cerebrospinal fluid (CSF). Discussion of fertility issues and sperm banking should be addressed in the appropriate circumstances.²²

Role of PET Scans

Staging

PET has a high positivity and specificity when used for the staging and restaging of lymphoma.²³ PET is nearly universally positive at diagnosis in DLBCL, FL, and nodal MZL, but less sensitive for extranodal MZL.^{24,25} However, a number of benign conditions including sarcoid, infection, and inflammation can result in false-positive PET scans, complicating the interpretation. Lesions <1 cm are not reliably visualized with PET scans.

Although PET scans may detect additional disease sites at diagnosis, the clinical stage is modified only in 15% to 20% of patients and a change in treatment in only 8% of patients.

PET scans are now virtually always performed as combined PET/CT scans. PET/CT has distinct advantages in both staging and restaging compared to full-dose diagnostic CT or PET alone.^{26,27} In a retrospective study, PET/CT performed with low-dose non-contrast-enhanced CT was found to be more sensitive and specific than contrast-enhanced CT for the evaluation of lymph node and organ involvement in patients with Hodgkin lymphoma or high-grade NHL.²⁶ Preliminary results of a prospective study (47 patients; patients who had undergone prior diagnostic CT were excluded) also showed a good correlation between low-dose non-contrast-enhanced PET/CT and full-dose contrast-enhanced PET/CT for the evaluation of lymph nodes and extranodal disease in lymphomas.²⁷

PET/CT should be done with contrast-enhanced diagnostic CT and is recommended for initial staging and restaging of all FDG-avid lymphomas.^{28,29} PET/CT is particularly important for staging before consideration of radiation therapy (RT) and baseline PET/CT will aid in the interpretation of post-treatment response evaluation based on the 5-point scale (5-PS) as described below.²⁹

Response Assessment

The guidelines for response criteria for lymphoma were first published in 1999 by the International Working Group (IWG).³⁰ These response criteria are based on the reduction in the size of the enlarged lymph node as measured by CT scan and the extent of bone marrow involvement that is determined by bone marrow aspirate and biopsy.³⁰ These guidelines were revised in 2007 by the International Harmonization Project to incorporate IHC, flow cytometry, and PET scans in the definition of response for lymphoma.³¹ In the revised guidelines, the response is categorized as



complete response (CR), partial response (PR), stable disease (SD), and relapsed disease or progressive disease (PD) based on the result of a PET scan. The response category of complete response uncertain (CRu) was essentially eliminated.

In 2014, revised response criteria, known as the Lugano criteria, were introduced for response assessment using PET/CT scans according to the 5-PS.^{28,29} The 5-PS is based on the visual assessment of fluorodeoxyglucose (FDG) uptake in the involved sites relative to that of the mediastinum and the liver.³²⁻³⁴ A score of 1 denotes no abnormal FDG-avidity, while a score of 2 represents uptake less than the mediastinum. A score of 3 denotes uptake greater than the mediastinum but less than the liver, while scores of 4 and 5 denote uptake greater than the liver, and greater than the liver with new sites of disease, respectively. Different clinical trials have considered scores of either 1 to 2 or 1 to 3 to be PET-negative, but a score of 1 to 3 is now widely considered to be PET negative. Scores of 4 to 5 are universally considered PET-positive. A score of 4 on an interim or end-of-treatment restaging scan may be consistent with a PR if the FDG avidity has declined from initial staging, while a score of 5 denotes PD.

However, the application of PET/CT to response assessment is limited to FDG-avid lymphomas and the revised response criteria have thus far only been validated for DLBCL and Hodgkin lymphoma. The application of the revised response criteria to other histologies requires validation and the original IWG guidelines should be used. False-positive PET scans may be observed related to infectious or inflammatory conditions. Biopsy of affected sites remains the gold standard for confirming new or persistent disease at end of therapy.

Principles of Radiation Therapy

RT can be delivered with photons, electrons, or protons, depending upon clinical circumstances.³⁵ Advanced RT techniques emphasize tightly conformal doses and steep gradients next to normal tissues. Therefore, target definition and delineation and treatment delivery verification require careful monitoring to avoid the risk of missing geographic location of the tumor and subsequent decrease in tumor control. Image guidance may be required to facilitate target definition. Preliminary results from single-institution studies have shown that significant dose reduction to organs at risk (OARs; eg, lungs, heart, breasts, kidneys, spinal cord, esophagus, carotid artery, bone marrow, stomach, muscle, soft tissue and salivary glands) can be achieved with advanced RT planning and delivery techniques such as 4D-CT simulation, intensity-modulated RT (IMRT), image-guided RT, respiratory gating, or deep inspiration breath hold.³⁶⁻³⁹ These techniques offer significant and clinically relevant advantages in specific instances to spare OARs and reduce the risk of late complications from normal tissue damage. This is especially important for patients being treated with curative intent or who have long life expectancies following therapy.⁴⁰⁻⁴³ In mediastinal lymphoma, the use of 4D-CT simulation and the adoption of strategies to deal with respiratory motion such as inspiration breath-hold techniques, and image guided RT during treatment delivery is also important.

Randomized prospective studies to test these concepts are unlikely to be done since these techniques are designed to decrease late effects, which usually develop ≥ 10 years after completion of treatment. Therefore, the guidelines recommend that RT delivery techniques that are found to best reduce the doses to the OAR in a clinically meaningful manner without compromising target coverage should be considered.

Involved-site RT (ISRT) is recommended as the appropriate field for NHL as it limits the radiation exposure to adjacent uninvolved organs (such as



lungs, bone, muscle, or kidney) when lymphadenopathy regresses following chemotherapy, thus minimizing the potential long-term complications.^{44,45} ISRT targets the initially involved nodal and extranodal sites detectable at presentation.^{44,45} Larger RT fields should be considered for limited-stage indolent NHL, often treated with RT alone.⁴⁴

Treatment planning for ISRT requires the use of CT-based simulation. The incorporation of additional imaging techniques such as PET and MRI often enhances the treatment planning. The OARs should be outlined for optimizing treatment plan decisions. The treatment plan is designed using conventional, 3D conformal, or IMRT techniques using clinical treatment planning considerations of coverage and dose reductions for OAR.⁴⁴

The principles of ISRT are similar for both nodal and extranodal disease. The gross tumor volume (GTV) defined by radiologic imaging prior to biopsy, chemotherapy, or surgery provides the basis for determining the clinical target volume (CTV).⁴⁶ Possible movement of the target by respiration as determined by 4D-CT or fluoroscopy should also influence the final CTV. The presence of suspected subclinical disease and uncertainties in original imaging accuracy or localization may lead to the expansion of the CTV. The planning treatment volume (PTV) is an additional expansion of the CTV that accounts only for setup variations.

In the case of extranodal disease, particularly for indolent lymphoma, in most cases the whole organ comprises the CTV (eg, stomach, salivary gland, and thyroid). For other organs, including orbit, breast, lung, bone, localized skin, and in some cases when RT is consolidation after chemotherapy, partial organ RT may be appropriate. No radiation is required for uninvolved lymph nodes for most NHL subtypes.

The general dose guidelines for individual subtypes of NHL are outlined in the *Principles of Radiation Therapy* section of the algorithm.

Biosimilars

A biosimilar is a biological product that is highly similar to the FDA approved reference biological product with the exception of minor differences in clinically inactive components and no clinically meaningful differences in safety, purity, or potency.⁴⁷

Pharmacokinetic (drug exposure) and pharmacodynamic (response) studies in the appropriate patient population are essential to demonstrate the efficacy and safety of the biosimilar.⁴⁸ Biosimilars require only one clinical trial to demonstrate equivalent safety and efficacy in the most sensitive indication for the reference biological product. If the mechanism of action, pharmacokinetics, and pharmacodynamics are similar, the biosimilar may be approved for all of the same indications as the reference biological product and can be substituted for the reference biological product.⁴⁸ Extrapolation of clinical and safety data from one indication to other approved indications is a key concept in the development of biosimilars that potentially provides substantial cost savings in oncology care, as biosimilars are typically more affordable than their reference products. Extrapolation should only be considered for indications where the mechanism of action is identical to that studied in the pivotal trial.

In November 2018, the FDA approved rituximab-abbs as the first biosimilar to rituximab for the following indications: 1) relapsed or refractory, low grade or follicular, CD20-positive B-cell NHL as a single agent; 2) previously untreated follicular, CD20-positive, B-cell NHL in combination with first-line chemotherapy and, in patients achieving a CR or PR to a rituximab product in combination with chemotherapy, as single-agent maintenance therapy; and 3) non-progressing (including SD), low-grade, CD20-positive, B-cell NHL as a single agent after first-line cyclophosphamide, vincristine and prednisone (CVP) chemotherapy.



In July 2019, the FDA approved rituximab-pvvr as the second biosimilar to rituximab for all of the same indications as rituximab-abbs. In addition, rituximab-pvvr is also approved for previously untreated DLBCL, CD20-positive NHL in combination with CHOP or other anthracycline-based chemotherapy regimens.

The approval of these two biosimilars is based on a review of evidence that included extensive structural and functional characterization, animal study data, human pharmacokinetic data, clinical immunogenicity data, and other clinical data demonstrating that the efficacy and safety profiles of rituximab-abbs and rituximab-pvvr are comparable to that of the reference biologic product (rituximab).⁴⁹⁻⁵¹

The efficacy, pharmacokinetics, and safety of rituximab-abbs (as a single agent and in combination with CVP chemotherapy) in comparison to rituximab was evaluated in 2 phase III randomized trials in patients with previously untreated low-burden FL (258 patients were randomized in the trial that evaluated rituximab-abbs as a single agent and 140 patients were enrolled in the trial that evaluated rituximab-abbs in combination with CVP).^{49,50} The efficacy and safety profile of rituximab-abbs were comparable to that of the reference biologic product (rituximab), although the incidences of grade 3 neutropenia were higher with rituximab-abbs in both studies.

The efficacy, pharmacokinetics, and safety of rituximab-pvvr as a single agent was evaluated in a phase III randomized trial in patients with previously untreated low-burden FL (394 patients were randomized to rituximab-pvvr or the reference biologic product). The efficacy, safety and immunogenicity, pharmacokinetics and pharmacodynamics of rituximab-pvvr were comparable to that of the reference biologic product (rituximab) up to 26 weeks.⁵¹

Rituximab-abbs is an FDA-approved biosimilar without an extrapolation to all of the indications as that of the reference biological product (rituximab) for NHL. Therefore, initially, the panel consensus was to include rituximab-abbs as an appropriate substitute for rituximab only for patients with indolent lymphomas (FL and MZL) as per the FDA approval. Rituximab-pvvr, however, is approved for all of the same indications as the reference biological product (rituximab), although the safety and efficacy of rituximab-pvvr was established only in patients with FL. Based on this approval, the panel consensus was to include an FDA-approved biosimilar (eg. rituximab-abbs or rituximab-pvvr) as an appropriate substitute for rituximab (for use as a single agent or in combination with chemotherapy) in all subtypes of B-cell lymphomas.

Alternating between the biosimilar and the reference product is acceptable without the intervention of a health care provider only if a biosimilar is designated as interchangeable since such a substitution will not result in higher toxicity or diminished efficacy.⁴⁷ However, alternating between the biosimilar and reference product is not recommended, if the biosimilar is not designated as interchangeable. Rituximab-abbs and rituximab-pvvr have not been approved as interchangeable biological products. Therefore, during a single course of therapy, the patient should remain on the same product that was used to initiate treatment (rituximab or rituximab-abbs or rituximab-pvvr) throughout the course of the treatment.

Supportive Care

Tumor Lysis Syndrome

Tumor lysis syndrome (TLS) is a potentially serious complication of anticancer therapy characterized by metabolic and electrolyte abnormalities caused by the disintegration of malignant cells by anticancer therapy and rapid release of intracellular contents into peripheral blood. It is usually observed within 12 to 72 hours after start of chemotherapy.⁵²



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Laboratory TLS is defined as a 25% increase in the levels of serum uric acid, potassium, or phosphorus or a 25% decrease in calcium levels.⁵³ Clinical TLS refers to laboratory TLS with clinical toxicity that requires intervention. Hyperkalemia, hyperuricemia, hyperphosphatemia, and hypocalcemia are the primary electrolyte abnormalities associated with TLS. Clinical symptoms may include nausea and vomiting, diarrhea, seizures, shortness of breath, renal insufficiency, or cardiac arrhythmias. Untreated TLS can induce profound metabolic changes resulting in cardiac arrhythmias, seizures, loss of muscle control, acute renal failure, and even death. The cornerstone of TLS management is hydration and the management of hyperuricemia. Allopurinol, febuxostat, and rasburicase are highly effective for the management of hyperuricemia.

Allopurinol is a xanthine analog and a competitive inhibitor of xanthine oxidase, thereby blocking the conversion of purine metabolites to uric acid and decreasing the formation of uric acid production.⁵⁴ Since the drug inhibits new uric acid formation rather than reduce existing uric acid, it can take several days for elevated levels of uric acid to normalize after the initiation of allopurinol, which may delay the start of chemoimmunotherapy. Furthermore, allopurinol may lead to the accumulation of xanthine crystals in renal tubules leading to acute obstructive uropathy. Allopurinol will also reduce clearance of 6-mercaptopurine and high-dose methotrexate.

Rasburicase is a recombinant urate oxidase, which catalyzes the oxidation of uric acid to a highly soluble non-toxic metabolite that is readily excreted. It has been shown to be safe and highly effective in the prevention and treatment of chemotherapy-induced hyperuricemia in both children and adults with hematologic malignancies.⁵⁵⁻⁵⁷ In a prospective, multicenter, randomized phase III trial that compared the efficacy and safety of rasburicase and allopurinol in adult patients with hematologic malignancies at high or potential risk for TLS (275 patients were randomized to receive rasburicase alone [0.20 mg/kg/d IV for days

1–5; n = 92], rasburicase combined with allopurinol [rasburicase 0.20 mg/kg/d IV for days 1–3; allopurinol 300 mg/d PO for days 3–5; n = 92], or allopurinol alone [300 mg/d PO for days 1–5; n = 91]), rasburicase was superior to allopurinol in the overall study population as well as in patients at high-risk for TLS.⁵⁷ The incidence of clinical TLS was similar across treatment arms, occurring in 3%, 3%, and 4% of patients, respectively. The incidence of laboratory TLS was 21%, 27%, and 41%, respectively, with significantly lower incidence observed in the rasburicase arm compared with allopurinol ($P = .003$). The median time to control for serum uric acid in hyperuricemic patients was 4 hours for rasburicase, 4 hours for rasburicase combined with allopurinol, and 27 hours for allopurinol. The rate of uric acid response (defined as plasma uric acid levels ≤ 7.5 mg/dL for all measurements from days 3–5 was 87% for rasburicase, 78% for rasburicase combined with allopurinol, and 66% for allopurinol. The response rate with rasburicase was superior to allopurinol in the overall study population (87% vs. 66%, as above; $P = .001$) as well as in patients with high-risk TLS (89% vs. 68%; $P = .001$) and in patients with baseline hyperuricemia (90% vs. 53%; $P = .015$). Potential hypersensitivity to study regimen was reported in 4% of patients in the rasburicase arm and 1% in the combination arm; no anaphylaxis or grade 4 hypersensitivity reactions were reported in this trial.⁵⁷ However, rasburicase can induce anaphylactic reactions. Other adverse reactions include methemoglobinemia and severe hemolysis in patients with glucose-6-phosphate dehydrogenase (G6PD) deficiency.

There are data to suggest that single fixed dose (6 mg or 3 mg) or single weight-based dose of rasburicase (0.05–0.15 mg/kg) are effective in adult patients with hyperuricemia or high-risk factors for TLS.⁵⁸⁻⁶³ In the phase II randomized trial that compared the efficacy of rasburicase administered as a single dose (0.15 mg/kg, followed by additional days of dosing as needed) versus rasburicase (0.15 mg/kg/d) given for 5 days in 80 adult patients at high risk or potential risk for TLS, nearly all treated



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

patients (99%) showed normalization of uric acid levels within 4 hours after the first dose of rasburicase; levels of uric acid were undetectable (<0.7 mg/dL) in 84% of patients.⁶³ The median pretreatment uric acid level was 8.5 mg/dL for high-risk patients (n = 40) and 5.6 mg/dL for potential risk patients (n = 40). In the single-dose rasburicase arm, 85% of patients had sustained uric acid response compared with 98% of patients in the 5-day rasburicase arm. Among high-risk patients within the single-dose arm, 6 patients received a second dose of rasburicase to achieve uric acid response.

In a randomized trial that compared the efficacy and safety of febuxostat and allopurinol in 346 adult patients with hematologic malignancies at intermediate or high risk for TLS, one fixed dose of febuxostat achieved a significantly superior serum uric acid control in comparison to allopurinol with comparable renal function preservation and safety profile.⁶⁴

TLS is best managed if anticipated and when treatment is started prior to chemoimmunotherapy. Histologies of Burkitt lymphoma, lymphoblastic lymphoma and occasionally DLBCL, bone marrow involvement, bulky tumors that are chemosensitive, rapidly proliferative or aggressive hematologic malignancies, an elevated leukocyte count or pretreatment LDH, pre-existing elevated uric acid, renal disease, or renal involvement of tumor are considered as risk factors for developing TLS.⁶⁵ TLS prophylaxis should be considered for patients with any of these risk factors. Frequent monitoring of electrolytes and aggressive correction is essential. The NCCN Guidelines recommend allopurinol or febuxostat (if intolerant to allopurinol) or rasburicase as first-line and at retreatment of hyperuricemia. Allopurinol should be started 2 to 3 days prior to the initiation of chemotherapy and continued for 10 to 14 days. Rasburicase is recommended for patients with any of the following risk factors: bulky disease requiring immediate therapy; patients in whom adequate hydration is not possible; or acute renal failure. A single dose is

adequate in most cases; repeat dosing should be given on an individual basis.

Hepatitis B Virus Reactivation

Testing for HBsAg and HBcAb can determine the HBV status of an individual. Because of the widespread use of the hepatitis B vaccine, HBsAb positivity is of limited value; however, in rare cases, HBsAb levels can help to guide therapy. Patients with malignancies who are positive for either HBsAg or HBcAb are at risk for HBV reactivation with cytotoxic chemotherapy; approximately 20% to 50% of patients with HBsAg positivity and 3% to 45% with HBcAb positivity develop HBV reactivation.⁶⁶⁻⁷⁵ False-negative HBsAg results may occur in chronic liver disease; therefore, patients with a history of hepatitis in need of chemotherapy should be assessed by viral load measurement.⁷⁶ HBsAb positivity is generally equated with protective immunity, although reactivated HBV disease may occur in the setting of significant immunosuppression in HBcAb-positive individuals.^{73,77} In patients with B-cell lymphomas treated with rituximab-containing regimens, HBV reactivation was observed in patients with HBcAb positivity (with or without HBsAb positivity), even among those who were HBsAg negative prior to initiation of treatment.^{68,74,75} A meta-analysis and evaluation of the FDA safety reports concerning HBV reactivation in patients with lymphoproliferative disorders reported that HBcAb positivity was correlated with increased incidence of rituximab-associated HBV reactivation.⁶⁷ Vaccination against HBV should be strongly considered in HBV-naïve patients (ie, negative for HBsAg, HBsAb, and HBcAb).^{73,78}

Recommended strategies for the management of HBV reactivation in patients with hematologic malignancies undergoing immunosuppressive therapy include upfront antiviral prophylaxis or pre-emptive therapy. Antiviral prophylaxis involves treating patients who are HBsAg-positive or HBcAb-positive with antiviral therapy, regardless of viral load or presence



of clinical manifestations of HBV reactivation. Pre-emptive therapy involves close surveillance with a highly sensitive quantitative assay for HBV, combined with antiviral therapy given at the time of serological evidence of HBV reactivation based upon a rising HBV DNA load.⁷³

Lamivudine has been shown to reduce the risks for HBV reactivation in HBsAg-positive patients with hematologic malignancies treated with immunosuppressive cytotoxic agents.^{13,79-81} A small randomized study in 30 HBsAg-positive patients with lymphoma showed that antiviral prophylaxis with lamivudine was superior to deferred pre-emptive therapy.⁷⁹ HBV reactivation occurred in 53% of patients in the deferred therapy arm compared with none in the prophylaxis arm. In a meta-analysis of clinical trials evaluating the benefit of lamivudine prophylaxis in HBsAg-positive lymphoma patients treated with immunosuppressive regimens, prophylaxis resulted in significant reductions in HBV reactivation and a trend for reduced HBV-related deaths compared with no prophylaxis.⁸¹

Entecavir has been shown to be more effective than lamivudine in preventing rituximab-associated HBV reactivation.^{82,83} In one randomized controlled trial, entecavir prophylaxis (before initiation of chemotherapy to 3 months after completion of chemotherapy) was more effective in preventing HBV reactivation than the control (initiation of entecavir therapy at the time of HBV reactivation and HBsAg reversed seroconversion after chemotherapy).⁸² The cumulative HBV reactivation rates at months 6, 12, and 18 after chemotherapy were 8%, 11%, and 26%, respectively, in the control group, and 0%, 0%, and 4% in the entecavir prophylaxis ($P = .019$). In another prospective study that compared the efficacy of antiviral prophylaxis with entecavir ($n = 61$) and lamivudine ($n = 60$) in HBsAg-positive patients with newly diagnosed DLBCL treated with R-CHOP chemoimmunotherapy, entecavir was associated with significantly lower rates of HBV reactivation (7% vs. 30%, $P = .001$),

HBV-related hepatitis (0% vs. 13%, $P = .003$), and disruption of chemotherapy (2% vs. 18%, $P = .002$) than lamivudine.⁸³

The panel recommends HBsAg and HBcAb testing for all patients planned for treatment with anti-CD20 MAB-containing regimens. In individuals who test positive for HBsAg and/or HBcAb, baseline quantitative polymerase chain reaction (PCR) for HBV DNA should be obtained to determine viral load. However, a negative baseline PCR does not preclude the possibility of reactivation. Patients receiving intravenous immunoglobulin (IVIG) may be HBcAb positive as a consequence of IVIG therapy, although HBV viral load monitoring is recommended.⁸⁴

Entecavir prophylaxis is recommended for patients who are HBsAg positive and receiving anti-lymphoma therapy.^{82,83} Entecavir prophylaxis is also the preferred approach for patients who are HBsAg negative but HBcAb positive; however, if these patients concurrently have high levels of HBsAb, they may be monitored with serial measurements of HBV viral load and treated with pre-emptive antivirals upon increasing viral load. Lamivudine should be avoided due to the risks for the development of resistance.⁸⁵⁻⁸⁷ Other antivirals such as adefovir, telbivudine, and tenofovir have also demonstrated antiviral efficacy in patients with chronic HBV infection and are acceptable alternatives.⁸⁸⁻⁹¹

The optimal choice of prophylactic antiviral therapy will be driven by institutional standards or recommendation from hepatology or an infectious disease consultant. The appropriate duration of prophylaxis remains undefined, but the panel recommended that surveillance and antiviral prophylaxis should be continued for up to 12 months after the completion of oncologic treatment.⁷³ During the treatment period, viral load should be monitored monthly with PCR and then every 3 months after completion of treatment. If viral load is consistently undetectable, prophylaxis with antivirals should be continued. If viral load fails to drop or a previously undetectable PCR becomes positive, consultation with a



hepatologist and discontinuation of anti-CD20 antibody therapy is recommended.

Hepatitis C Virus-associated B-Cell Lymphomas

As noted earlier, large population-based or multicenter case-control studies have demonstrated a strong association between seropositivity for HCV and development of B-cell lymphomas.¹⁴⁻²¹ The prevalence of HCV seropositivity was consistently increased among patients with DLBCL and MZL.^{14,15,19,20} A retrospective study in patients with HCV infection showed that the cumulative incidence of developing malignant lymphomas was significantly higher among patients with persistent HCV infection compared with those who had sustained virologic response (SVR) to interferon-containing therapy (15-year incidence rate 3% vs. 0%; $P = .016$).¹⁷ Based on multivariate analysis, persistent HCV infection remained a significant independent factor associated with development of malignant lymphomas. This study suggested that achievement of SVR with interferon-based therapy may reduce the incidence of malignant lymphoma in patients with HCV infection.¹⁷

Several published reports suggested that treatment with antivirals (typically interferon with or without ribavirin) led to regression of NHLs in HCV-positive patients, which provides additional evidence for the involvement of HCV infection in the pathogenesis of lymphoproliferative diseases.⁹²⁻⁹⁸ In a retrospective study in patients with NHL (N = 343; indolent and aggressive histologies) who achieved a CR after chemotherapy, the subgroup of HCV-positive patients treated with antivirals (interferon and ribavirin; n = 25) had significantly longer disease-free survival compared with HCV-positive patients who did not receive antiviral therapy (n = 44); the probability of relapse-free survival at 5-year follow up was 76% and 55%, respectively.⁹⁷ In addition, none of the patients with an SVR to antivirals (n = 0 of 8) relapsed compared with 29% who did not respond to antivirals (n = 5 of 17).

In a multicenter retrospective study from a large series of HCV-positive patients with indolent NHL, antiviral therapy (interferon or pegylated interferon, with or without ribavirin) resulted in HCV-RNA clearance being achieved in 80% of patients who received first-line antivirals (n = 100) and in 67% of those who received antivirals as second-line therapy after failure of initial treatment (n = 34).⁹⁸ Patients in this analysis did not require immediate treatment for their lymphoma. The ORR for patients treated with antiviral in the first-line setting was 77% (44% CR and 33% PR) and the ORR for patients treated with antiviral in the second-line setting was 85% (56% CR and 29% PR). In the group of patients who received antivirals in first line, hematologic response was significantly associated with achievement of HCV-RNA clearance. Thus, in HCV-positive patients with indolent NHL not requiring immediate anti-tumor therapy with chemoimmunotherapy regimens, initial treatment with interferon (with or without ribavirin) appeared to induce lymphoma regression in a high proportion of patients. In HCV-positive patients with NHL who achieve a remission with anti-tumor therapy, subsequent treatment with antivirals may be associated with lower risk of disease relapse.

The optimal management of NHL in HCV-positive patients remains to be defined. Patients with indolent NHL may benefit from antiviral treatment as initial therapy, as demonstrated in several reports.^{92,94,96,98,99} In patients with aggressive NHL, an earlier analysis of pooled data from GELA clinical studies (prior to the rituximab era) suggested that HCV seropositivity in patients with DLBCL was associated with significantly decreased survival outcomes, due, in part, to severe hepatotoxicity among those with HCV infection.¹⁰⁰ Subsequent studies in the rituximab era showed that HCV seropositivity was not predictive of outcomes in terms of PFS or OS in patients with DLBCL.^{101,102} However, the incidence of hepatotoxicity with chemoimmunotherapy was higher among HCV-positive patients, confirming the observation made from the GELA studies.



The treatment of chronic HCV infection has improved with the advent of newer antiviral agents, especially those that target carriers of HCV genotype 1. Direct-acting antiviral agents (DAA) administered in combination with standard antivirals (pegylated interferon and ribavirin) have shown significantly higher rates of SVR compared with standard therapy alone in chronic carriers of HCV genotype 1.¹⁰³⁻¹⁰⁶ Telaprevir and boceprevir are DAA that were recently approved by the FDA for the treatment (in combination with pegylated interferon and ribavirin) of patients with HCV genotype 1 infection. The updated guidelines for the management of HCV infection from the American Association for the Study of Liver Diseases (AASLD) recommended that DAA be incorporated into standard antiviral therapy for patients infected with HCV genotype 1.¹⁰⁷

The panel recommends initial antiviral therapy in asymptomatic patients with HCV-positive low-grade B-cell NHL. For those with HCV genotype 1, triple antiviral therapy with inclusion of DAAs should be considered as per AASLD guidelines. Patients with HCV-positive aggressive B-cell NHL should initially be treated with appropriate chemoimmunotherapy regimens according to the NCCN Guidelines for NHL. Liver function and serum HCV RNA levels should be closely monitored for the development of hepatotoxicity during and after chemoimmunotherapy. Antiviral therapy should then be considered in patients who achieve a CR after completion of chemoimmunotherapy.

Progressive Multifocal Leukoencephalopathy

Progressive multifocal leukoencephalopathy (PML) is a rare but serious and usually fatal CNS infection caused by reactivation of the latent JC polyomavirus. PML generally occurs in severely immunocompromised individuals, as in the case of patients with AIDS and in patients with low CD4+ T-cells prior to or during anti-lymphoma treatment.¹⁰⁸⁻¹¹⁰ PML has been reported in patients with NHL receiving treatment with rituximab

(usually in combination with chemotherapy) or brentuximab vedotin.^{108,111} In a report of 57 patients from the Research on Adverse Drug Events and Reports project, 52 patients with lymphoproliferative disorders developed PML after treatment with rituximab and other treatments, which included hematopoietic stem cell transplantation or chemotherapy with purine analogs or alkylating agents.¹⁰⁸ Median time from last rituximab dose to PML diagnosis was 6 months. Median time to death after PML diagnosis was 2 months, with a case fatality rate of 90%.

PML is clinically suspected based on neurologic signs and symptoms that may include confusion, motor weakness or poor motor coordination, visual changes, and/or speech changes.¹⁰⁸ PML is usually diagnosed with PCR of CSF or, in some cases, by analysis of brain biopsy material. There is no effective treatment for PML. Patients should be carefully monitored for the development of any neurologic symptoms. There is currently no consensus on pretreatment evaluations that can be undertaken to predict for the subsequent development of PML.

Management of anti-CD20 Monoclonal Antibody Therapy Intolerance

Rare complications such as mucocutaneous reactions including paraneoplastic pemphigus, Stevens-Johnson syndrome, lichenoid dermatitis, vesiculobullous dermatitis, and toxic epidermal necrolysis can occur in patients treated with anti-CD20 MAB. Expert consultation with a dermatologist is recommended.

Re-challenge with the same anti-CD20 MAB is not recommended in patients experiencing the aforementioned complications to chosen anti-CD20 MAB (rituximab, obinutuzumab, or ofatumumab). There are data supporting the use of ofatumumab in patients experiencing intolerance to rituximab.^{112,113} An alternative anti-CD20 MAB (obinutuzumab or ofatumumab) could be used for patients with intolerance



to rituximab, regardless of histology. However, it is unclear if such a substitution poses the same risk of recurrence.

Rituximab and hyaluronidase human injection for subcutaneous use is approved by the FDA for the treatment of patients with previously untreated and relapsed/refractory FL and previously untreated DLBCL, only for patients who have received at least one full dose of intravenous rituximab. The FDA approval was based on the results of two large phase III randomized trials that confirmed the non-inferiority for subcutaneous rituximab (1400 mg SC dose) compared with intravenous rituximab (375 mg/m²) when used in combination with chemotherapy (CHOP or CVP) in patients with previously untreated FL (SABRINA study)¹¹⁴ or DLBCL (MabEase study).¹¹⁵

The guidelines recommend that rituximab and hyaluronidase human injection for subcutaneous use may be substituted for intravenous rituximab after patients have received the first full dose of rituximab by intravenous infusion. Switching to subcutaneous rituximab is not recommended until a full intravenous dose of rituximab is successfully administered without experiencing severe adverse reactions. This substitution with subcutaneous rituximab is also not recommended for intravenous rituximab used in combination with ibritumomab tiuxetan.

Management of Bone Health in Patients Receiving Steroid-Containing Regimens

Steroid-containing systemic therapy regimens have been associated with increased risk of fractures and treatment-induced bone loss in patients with NHL.¹¹⁶⁻¹¹⁸ The risk of treatment-induced bone loss is higher among young women with chemotherapy-induced premature menopause and older patients receiving chemotherapy.^{119,120} In addition, patients with newly diagnosed NHL are also at risk of low bone mineral density (BMD), which may worsen during treatment with steroid-based systemic therapy.¹²¹

Evaluation of vitamin D levels and post-treatment BMD evaluation using a fracture risk assessment tool is recommended for patients receiving steroid-based systemic therapy.^{122,123} Referral to an endocrinologist with expertise on bone health and initiation of treatment as per National Osteoporosis Foundation guidelines is recommended for patients with osteoporotic BMD, with a history of hip or vertebral fractures, or with asymptomatic vertebral compression deformity (as seen on imaging studies).¹²⁴

Adequate calcium intake is of paramount importance since corticosteroids block calcium absorption and increase the risk of fracture. Raloxifene or hormone replacement therapy up until the expected time of natural menopause could be considered in appropriate women with premature menopause. The use of bisphosphonates (pamidronate and zoledronic acid) has been shown to effectively stabilize BMD, prevent bone loss, and reduce the risk of new vertebral fractures in patients with NHL.^{125,126} Bisphosphonates should be used as first-line pharmacologic treatment for osteoporosis. Denosumab is an effective alternative option to prevent osteoporotic fractures in patients who cannot tolerate or whose symptoms do not improve with bisphosphonate therapy.¹²⁷



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. CA Cancer J Clin 2019;69:7-34. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30620402>.
2. Al-Hamadani M, Habermann TM, Cerhan JR, et al. Non-Hodgkin lymphoma subtype distribution, geodemographic patterns, and survival in the US: A longitudinal analysis of the National Cancer Data Base from 1998 to 2011. Am J Hematol 2015;90:790-795. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26096944>.
3. Groves FD, Linet MS, Travis LB, Devesa SS. Cancer surveillance series: non-Hodgkin's lymphoma incidence by histologic subtype in the United States from 1978 through 1995. J Natl Cancer Inst 2000;92:1240-1251. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10922409>.
4. Swerdlow SH, Campo E, Harris NL, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues, Fourth Edition, Lyon, France: IARC; ; 2008.
5. Swerdlow SH, Harris NL, Jaffe ES, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Revised 4th ed. Lyon, France: IARC; 2017.
6. Hehn ST, Grogan TM, Miller TP. Utility of fine-needle aspiration as a diagnostic technique in lymphoma. J Clin Oncol 2004;22:3046-3052. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15284254>.
7. Meda BA, Buss DH, Woodruff RD, et al. Diagnosis and subclassification of primary and recurrent lymphoma. The usefulness and limitations of combined fine-needle aspiration cytomorphology and flow cytometry. Am J Clin Pathol 2000;113:688-699. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10800402>.
8. Dong HY, Harris NL, Preffer FI, Pitman MB. Fine-needle aspiration biopsy in the diagnosis and classification of primary and recurrent lymphoma: a retrospective analysis of the utility of cytomorphology and flow cytometry. Mod Pathol 2001;14:472-481. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11353059>.
9. Jeffers MD, Milton J, Herriot R, McKean M. Fine needle aspiration cytology in the investigation on non-Hodgkin's lymphoma. J Clin Pathol 1998;51:189-196. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9659258>.
10. Zeppa P, Marino G, Troncone G, et al. Fine-needle cytology and flow cytometry immunophenotyping and subclassification of non-Hodgkin lymphoma: a critical review of 307 cases with technical suggestions. Cancer 2004;102:55-65. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14968418>.
11. Dunphy CH. Applications of flow cytometry and immunohistochemistry to diagnostic hematopathology. Arch Pathol Lab Med 2004;128:1004-1022. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15335254>.
12. Kusumoto S, Arcaini L, Hong X, et al. Risk of HBV reactivation in patients with B-cell lymphomas receiving obinutuzumab or rituximab immunochemotherapy. Blood 2019;133:137-146. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30341058>.
13. Yeo W, Chan PKS, Ho WM, et al. Lamivudine for the prevention of hepatitis B virus reactivation in hepatitis B s-antigen seropositive cancer patients undergoing cytotoxic chemotherapy. J Clin Oncol 2004;22:927-934. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14990649>.
14. Engels EA, Chatterjee N, Cerhan JR, et al. Hepatitis C virus infection and non-Hodgkin lymphoma: results of the NCI-SEER multi-center case-control study. Int J Cancer 2004;111:76-80. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15185346>.
15. Nieters A, Kallinowski B, Brennan P, et al. Hepatitis C and risk of lymphoma: results of the European multicenter case-control study EPILYMPH. Gastroenterology 2006;131:1879-1886. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17087949>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

16. Arcaini L, Burcheri S, Rossi A, et al. Prevalence of HCV infection in nongastric marginal zone B-cell lymphoma of MALT. *Ann Oncol* 2007;18:346-350. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17071937>.

17. Kawamura Y, Ikeda K, Arase Y, et al. Viral elimination reduces incidence of malignant lymphoma in patients with hepatitis C. *Am J Med* 2007;120:1034-1041. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18060923>.

18. Schollkopf C, Smedby KE, Hjalgrim H, et al. Hepatitis C infection and risk of malignant lymphoma. *Int J Cancer* 2008;122:1885-1890. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18271005>.

19. de Sanjose S, Benavente Y, Vajdic CM, et al. Hepatitis C and non-Hodgkin lymphoma among 4784 cases and 6269 controls from the International Lymphoma Epidemiology Consortium. *Clin Gastroenterol Hepatol* 2008;6:451-458. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18387498>.

20. Spinelli JJ, Lai AS, Krajden M, et al. Hepatitis C virus and risk of non-Hodgkin lymphoma in British Columbia, Canada. *Int J Cancer* 2008;122:630-633. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17935132>.

21. Arcaini L, Merli M, Volpetti S, et al. Indolent B-cell lymphomas associated with HCV infection: clinical and virological features and role of antiviral therapy. *Clin Dev Immunol* 2012;2012:638185. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22956970>.

22. Howell SJ, Shalet SM. Fertility preservation and management of gonadal failure associated with lymphoma therapy. *Curr Oncol Rep* 2002;4:443-452. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12162920>.

23. Isasi CR, Lu P, Blaufox MD. A metaanalysis of 18F-2-deoxy-2-fluoro-D-glucose positron emission tomography in the staging and restaging of patients with lymphoma. *Cancer*

2005;104:1066-1074. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16047335>.

24. Trotman J, Fournier M, Lamy T, et al. Positron emission tomography-computed tomography (PET-CT) after induction therapy is highly predictive of patient outcome in follicular lymphoma: analysis of PET-CT in a subset of PRIMA trial participants. *J Clin Oncol* 2011;29:3194-3200. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21747087>.

25. Hoffmann M, Kletter K, Becherer A, et al. 18F-fluorodeoxyglucose positron emission tomography (18F-FDG-PET) for staging and follow-up of marginal zone B-cell lymphoma. *Oncology* 2003;64:336-340. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12759529>.

26. Schaefer NG, Hany TF, Taverna C, et al. Non-Hodgkin lymphoma and Hodgkin disease: coregistered FDG PET and CT at staging and restaging--do we need contrast-enhanced CT? *Radiology* 2004;232:823-829. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15273335>.

27. Rodriguez-Vigil B, Gomez-Leon N, Pinilla I, et al. PET/CT in lymphoma: prospective study of enhanced full-dose PET/CT versus unenhanced low-dose PET/CT. *J Nucl Med* 2006;47:1643-1648. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17015900>.

28. Barrington SF, Mikhael NG, Kostakoglu L, et al. Role of imaging in the staging and response assessment of lymphoma: consensus of the International Conference on Malignant Lymphomas Imaging Working Group. *J Clin Oncol* 2014;32:3048-3058. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25113771>.

29. Cheson BD, Fisher RI, Barrington SF, et al. Recommendations for initial evaluation, staging, and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. *J Clin Oncol* 2014;32:3059-3068. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25113753>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

30. Cheson BD, Horning SJ, Coiffier B, et al. Report of an international workshop to standardize response criteria for non-Hodgkin's lymphomas. NCI Sponsored International Working Group. J Clin Oncol 1999;17:1244. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10561185>.

31. Cheson BD, Pfistner B, Juweid ME, et al. Revised response criteria for malignant lymphoma. J Clin Oncol 2007;25:579-586. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17242396>.

32. Barrington SF, Qian W, Somer EJ, et al. Concordance between four European centres of PET reporting criteria designed for use in multicentre trials in Hodgkin lymphoma. European Journal of Nuclear Medicine and Molecular Imaging 2010;37:1824-1833. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20505930>.

33. Meignan M, Gallamini A, Haioun C, Polliack A. Report on the Second International Workshop on interim positron emission tomography in lymphoma held in Menton, France, 8-9 April 2010. Leuk Lymphoma 2010;51:2171-2180. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21077737>.

34. Meignan M, Gallamini A, Itti E, et al. Report on the Third International Workshop on Interim Positron Emission Tomography in Lymphoma held in Menton, France, 26-27 September 2011 and Menton 2011 consensus. Leuk Lymphoma 2012;53:1876-1881. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22432519>.

35. Tseng YD, Cutter DJ, Plastaras JP, et al. Evidence-based Review on the Use of Proton Therapy in Lymphoma From the Particle Therapy Cooperative Group (PTCOG) Lymphoma Subcommittee. Int J Radiat Oncol Biol Phys 2017;99:825-842. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28943076>.

36. Jang JW, Brown JG, Mauch PM, Ng AK. Four-dimensional versus 3-dimensional computed tomographic planning for gastric mucosa associated lymphoid tissue lymphoma. Pract Radiat Oncol 2013;3:124-129. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24674315>.

37. Xu LM, Li YX, Fang H, et al. Dosimetric evaluation and treatment outcome of intensity modulated radiation therapy after doxorubicin-based chemotherapy for primary mediastinal large B-cell lymphoma. Int J Radiat Oncol Biol Phys 2013;85:1289-1295. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23245279>.

38. Besson N, Pernin V, Zefkili S, Kirova YM. Evolution of radiation techniques in the treatment of mediastinal lymphoma: from 3D conformal radiotherapy (3DCRT) to intensity-modulated RT (IMRT) using helical tomotherapy (HT): a single-centre experience and review of the literature. Br J Radiol 2016;89:20150409. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26744079>.

39. Boda-Heggemann J, Knopf AC, Simeonova-Chergou A, et al. Deep Inspiration Breath Hold-Based Radiation Therapy: A Clinical Review. Int J Radiat Oncol Biol Phys 2016;94:478-492. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26867877>.

40. Girinsky T, Pichenot C, Beaudre A, et al. Is intensity-modulated radiotherapy better than conventional radiation treatment and three-dimensional conformal radiotherapy for mediastinal masses in patients with Hodgkin's disease, and is there a role for beam orientation optimization and dose constraints assigned to virtual volumes? Int J Radiat Oncol Biol Phys 2006;64:218-226. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16169675>.

41. Nieder C, Schill S, Kneschaurek P, Molls M. Influence of different treatment techniques on radiation dose to the LAD coronary artery. Radiat Oncol 2007;2:20. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17547777>.

42. Filippi AR, Ragona R, Fusella M, et al. Changes in breast cancer risk associated with different volumes, doses, and techniques in female Hodgkin lymphoma patients treated with supra-diaphragmatic radiation therapy. Pract Radiat Oncol 2013;3:216-222. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24674367>.

43. Charpentier AM, Conrad T, Sykes J, et al. Active breathing control for patients receiving mediastinal radiation therapy for lymphoma: Impact on



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

normal tissue dose. *Pract Radiat Oncol* 2014;4:174-180. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24766684>.

44. Illidge T, Specht L, Yahalom J, et al. Modern radiation therapy for nodal non-Hodgkin lymphoma-target definition and dose guidelines from the International Lymphoma Radiation Oncology Group. *Int J Radiat Oncol Biol Phys* 2014;89:49-58. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24725689>.

45. Yahalom J, Illidge T, Specht L, et al. Modern radiation therapy for extranodal lymphomas: field and dose guidelines from the International Lymphoma Radiation Oncology Group. *Int J Radiat Oncol Biol Phys* 2015;92:11-31. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25863750>.

46. Hoskin PJ, Diez P, Williams M, et al. Recommendations for the use of radiotherapy in nodal lymphoma. *Clin Oncol (R Coll Radiol)* 2013;25:49-58. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22889569>.

47. U.S. Food and Drug Administration: Biosimilar and Interchangeable Products. Available at: <https://www.fda.gov/Drugs/DevelopmentApprovalProcess/HowDrugsareDevelopedandApproved/ApprovalApplications/TherapeuticBiologicApplications/Biosimilars/ucm580419.htm>.

48. U.S. Food and Drug Administration: Biosimilar Development, Review, and Approval. Available at: <https://www.fda.gov/Drugs/DevelopmentApprovalProcess/HowDrugsareDevelopedandApproved/ApprovalApplications/TherapeuticBiologicApplications/Biosimilars/ucm580429.htm>.

49. Kim WS, Buske C, Ogura M, et al. Efficacy, pharmacokinetics, and safety of the biosimilar CT-P10 compared with rituximab in patients with previously untreated advanced-stage follicular lymphoma: a randomised, double-blind, parallel-group, non-inferiority phase 3 trial. *Lancet Haematol* 2017;4:e362-e373. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28712940>.

50. Ogura M, Sancho JM, Cho SG, et al. Efficacy, pharmacokinetics, and safety of the biosimilar CT-P10 in comparison with rituximab in patients with previously untreated low-tumour-burden follicular lymphoma: a randomised, double-blind, parallel-group, phase 3 trial. *Lancet Haematol* 2018;5:e543-e553. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30389036>.

51. Sharman J, Liberati AM, Santucci Silva R, et al. A randomized, double-blind efficacy and safety study of PF-05280586 (a potential rituximab biosimilar) compared with rituximab reference product (MabThera) in subjects with previously untreated CD20-positive, low tumor burden follicular lymphoma (LTB-FL) [abstract]. *Blood* 2018;132:Abstract 394. Available at: http://www.bloodjournal.org/content/132/Suppl_1/394.abstract.

52. Coiffier B, Altman A, Pui C, et al. Guidelines for the management of pediatric and adult tumor lysis syndrome: an evidence-based review. *J Clin Oncol* 2008;26:2767-2778. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18509186>.

53. Cairo MS, Bishop M. Tumour lysis syndrome: new therapeutic strategies and classification. *Br J Haematol* 2004;127:3-11. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15384972>.

54. Krakoff IH, Meyer RL. Prevention of hyperuricemia in leukemia and lymphoma: use of allopurinol, a xanthine oxidase inhibitor. *JAMA* 1965;193:1-6. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14297704>.

55. Bosly A, Sonet A, Pinkerton CR, et al. Rasburicase (recombinant urate oxidase) for the management of hyperuricemia in patients with cancer: report of an international compassionate use study. *Cancer* 2003;98:1048-1054. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12942574>.

56. Coiffier B, Mounier N, Bologna S, et al. Efficacy and safety of rasburicase (recombinant urate oxidase) for the prevention and treatment of hyperuricemia during induction chemotherapy of aggressive non-Hodgkin's lymphoma: results of the GRAAL1 (Groupe d'Etude des



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Lymphomes de l'Adulte Trial on Rasburicase Activity in Adult Lymphoma) study. J Clin Oncol 2003;21:4402-4406. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14581437>.

57. Cortes J, Moore JO, Maziarz RT, et al. Control of plasma uric acid in adults at risk for tumor lysis syndrome: efficacy and safety of rasburicase alone and rasburicase followed by allopurinol compared with allopurinol alone—results of a multicenter phase III study. J Clin Oncol 2010;28:4207-4213. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20713865>.

58. McDonnell AM, Lenz KL, Frei-Lahr DA, et al. Single-dose rasburicase 6 mg in the management of tumor lysis syndrome in adults. Pharmacotherapy 2006;26:806-812. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16716134>.

59. Campara M, Shord SS, Haaf CM. Single-dose rasburicase for tumour lysis syndrome in adults: weight-based approach. J Clin Pharm Ther 2009;34:207-213. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19250141>.

60. Trifilio SM, Pi J, Zook J, et al. Effectiveness of a single 3-mg rasburicase dose for the management of hyperuricemia in patients with hematological malignancies. Bone Marrow Transplant 2011;46:800-805. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20818444>.

61. Vines AN, Shanholtz CB, Thompson JL. Fixed-dose rasburicase 6 mg for hyperuricemia and tumor lysis syndrome in high-risk cancer patients. Ann Pharmacother 2010;44:1529-1537. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20841516>.

62. Knoebel RW, Lo M, Crank CW. Evaluation of a low, weight-based dose of rasburicase in adult patients for the treatment or prophylaxis of tumor lysis syndrome. J Oncol Pharm Pract 2011;17:147-154. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20332174>.

63. Vadhan-Raj S, Fayad LE, Fanale MA, et al. A randomized trial of a single-dose rasburicase versus five-daily doses in patients at risk for tumor

lysis syndrome. Ann Oncol 2012;23:1640-1645. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22015451>.

64. Spina M, Nagy Z, Ribera JM, et al. FLORENCE: a randomized, double-blind, phase III pivotal study of febuxostat versus allopurinol for the prevention of tumor lysis syndrome (TLS) in patients with hematologic malignancies at intermediate to high TLS risk. Ann Oncol 2015;26:2155-2161. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26216382>.

65. Cairo MS, Coiffier B, Reiter A, et al. Recommendations for the evaluation of risk and prophylaxis of tumour lysis syndrome (TLS) in adults and children with malignant diseases: an expert TLS panel consensus. Br J Haematol 2010;149:578-586. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20331465>.

66. Borentain P, Colson P, Coso D, et al. Clinical and virological factors associated with hepatitis B virus reactivation in HBsAg-negative and anti-HBc antibodies-positive patients undergoing chemotherapy and/or autologous stem cell transplantation for cancer. J Viral Hepat 2010;17:807-815. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20002298>.

67. Evens AM, Jovanovic BD, Su YC, et al. Rituximab-associated hepatitis B virus (HBV) reactivation in lymphoproliferative diseases: meta-analysis and examination of FDA safety reports. Ann Oncol 2011;22:1170-1180. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21115603>.

68. Fukushima N, Mizuta T, Tanaka M, et al. Retrospective and prospective studies of hepatitis B virus reactivation in malignant lymphoma with occult HBV carrier. Ann Oncol 2009;20:2013-2017. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19561036>.

69. Ji D, Cao J, Hong X, et al. Low incidence of hepatitis B virus reactivation during chemotherapy among diffuse large B-cell lymphoma patients who are HBsAg-negative/ HBcAb-positive: a multicenter retrospective study. Eur J Haematol 2010;85:243-250. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20491883>.



70. Koo YX, Tan DS, Tan IB, et al. Hepatitis B virus reactivation and role of antiviral prophylaxis in lymphoma patients with past hepatitis B virus infection who are receiving chemioimmunotherapy. *Cancer* 2010;116:115-121. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19899164>.

71. Lalazar G, Rund D, Shouval D. Screening, prevention and treatment of viral hepatitis B reactivation in patients with haematological malignancies. *Br J Haematol* 2007;136:699-712. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17338776>.

72. Lau GK, Leung YH, Fong DY, et al. High hepatitis B virus (HBV) DNA viral load as the most important risk factor for HBV reactivation in patients positive for HBV surface antigen undergoing autologous hematopoietic cell transplantation. *Blood* 2002;99:2324-2330. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11895763>.

73. Liang R. How I treat and monitor viral hepatitis B infection in patients receiving intensive immunosuppressive therapies or undergoing hematopoietic stem cell transplantation. *Blood* 2009;113:3147-3153. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19144986>.

74. Matsue K, Kimura S, Takanashi Y, et al. Reactivation of hepatitis B virus after rituximab-containing treatment in patients with CD20-positive B-cell lymphoma. *Cancer* 2010;116:4769-4776. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20597091>.

75. Yeo W, Chan TC, Leung NW, et al. Hepatitis B virus reactivation in lymphoma patients with prior resolved hepatitis B undergoing anticancer therapy with or without rituximab. *J Clin Oncol* 2009;27:605-611. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19075267>.

76. Brechot C, Degos F, Lugassy C, et al. Hepatitis B virus DNA in patients with chronic liver disease and negative tests for hepatitis B surface antigen. *N Engl J Med* 1985;312:270-276. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2981408>.

77. Dhedin N, Douvin C, Kuentz M, et al. Reverse seroconversion of hepatitis B after allogeneic bone marrow transplantation: a retrospective

study of 37 patients with pretransplant anti-HBs and anti-HBc. *Transplantation* 1998;66:616-619. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9753342>.

78. Lok AS, McMahon BJ. Chronic hepatitis B. *Hepatology* 2007;45:507-539. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17256718>.

79. Lau GKK, Yiu HHY, Fong DYT, et al. Early is superior to deferred preemptive lamivudine therapy for hepatitis B patients undergoing chemotherapy. *Gastroenterology* 2003;125:1742-1749. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14724827>.

80. Tsutsumi Y, Tanaka J, Kawamura T, et al. Possible efficacy of lamivudine treatment to prevent hepatitis B virus reactivation due to rituximab therapy in a patient with non-Hodgkin's lymphoma. *Ann Hematol* 2004;83:58-60. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14513286>.

81. Ziakas PD, Karsaliakos P, Mylonakis E. Effect of prophylactic lamivudine for chemotherapy-associated hepatitis B reactivation in lymphoma: a meta-analysis of published clinical trials and a decision tree addressing prolonged prophylaxis and maintenance. *Haematologica* 2009;94:998-1005. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19454492>.

82. Huang YH, Hsiao LT, Hong YC, et al. Randomized controlled trial of entecavir prophylaxis for rituximab-associated hepatitis B virus reactivation in patients with lymphoma and resolved hepatitis B. *J Clin Oncol* 2013;31:2765-2772. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23775967>.

83. Huang H, Li X, Zhu J, et al. Entecavir vs lamivudine for prevention of hepatitis B virus reactivation among patients with untreated diffuse large B-cell lymphoma receiving R-CHOP chemotherapy: a randomized clinical trial. *JAMA* 2014;312:2521-2530. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25514302>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

84. Arnold DM, Crowther MA, Meyer RM, et al. Misleading hepatitis B test results due to intravenous immunoglobulin administration: implications for a clinical trial of rituximab in immune thrombocytopenia. *Transfusion* 2010;50:2577-2581. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20576011>.

85. Law JK, Ali JA, Harrigan PR, et al. Fatal postlymphoma chemotherapy hepatitis B reactivation secondary to the emergence of a YMDD mutant strain with lamivudine resistance in a noncirrhotic patient. *Am J Hematol* 2006;81:969-972. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16937392>.

86. Picardi M, Pane F, Quintarelli C, et al. Hepatitis B virus reactivation after fludarabine-based regimens for indolent non-Hodgkin's lymphomas: high prevalence of acquired viral genomic mutations. *Haematologica* 2003;88:1296-1303. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14607759>.

87. Schnepf N, Sellier P, Bendenoun M, et al. Reactivation of lamivudine-resistant occult hepatitis B in an HIV-infected patient undergoing cytotoxic chemotherapy. *J Clin Virol* 2007;39:48-50. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17368969>.

88. Peters MG, Hann HW, Martin P, et al. Adefovir dipivoxil alone or in combination with lamivudine in patients with lamivudine-resistant chronic hepatitis B. *Gastroenterology* 2004;126:91-101. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14699491>.

89. Chan HL, Heathcote EJ, Marcellin P, et al. Treatment of hepatitis B e antigen positive chronic hepatitis with telbivudine or adefovir: a randomized trial. *Ann Intern Med* 2007;147:745-754. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17909201>.

90. Marcellin P, Heathcote EJ, Buti M, et al. Tenofovir disoproxil fumarate versus adefovir dipivoxil for chronic hepatitis B. *N Engl J Med* 2008;359:2442-2455. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19052126>.

91. Leung N, Peng C-Y, Hann H-W, et al. Early hepatitis B virus DNA reduction in hepatitis B e antigen-positive patients with chronic hepatitis B: A randomized international study of entecavir versus adefovir. *Hepatology* 2009;49:72-79. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19065670>.

92. Hermine O, Lefrere F, Bronowicki JP, et al. Regression of splenic lymphoma with villous lymphocytes after treatment of hepatitis C virus infection. *N Engl J Med* 2002;347:89-94. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12110736>.

93. Kelaidi C, Rollot F, Park S, et al. Response to antiviral treatment in hepatitis C virus-associated marginal zone lymphomas. *Leukemia* 2004;18:1711-1716. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15284859>.

94. Tursi A, Brandimarte G, Torello M. Disappearance of gastric mucosa-associated lymphoid tissue in hepatitis C virus-positive patients after anti-hepatitis C virus therapy. *J Clin Gastroenterol* 2004;38:360-363. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15087696>.

95. Gisbert JP, Garcia-Buey L, Pajares JM, Moreno-Otero R. Systematic review: regression of lymphoproliferative disorders after treatment for hepatitis C infection. *Aliment Pharmacol Ther* 2005;21:653-662. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15771751>.

96. Vallisa D, Bernuzzi P, Arcaini L, et al. Role of anti-hepatitis C virus (HCV) treatment in HCV-related, low-grade, B-cell, non-Hodgkin's lymphoma: a multicenter Italian experience. *J Clin Oncol* 2005;23:468-473. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15659492>.

97. La Mura V, De Renzo A, Perna F, et al. Antiviral therapy after complete response to chemotherapy could be efficacious in HCV-positive non-Hodgkin's lymphoma. *J Hepatol* 2008;49:557-563. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18678434>.

98. Arcaini L, Vallisa D, Rattotti S, et al. Antiviral treatment in patients with indolent B-cell lymphomas associated with HCV infection: a study of the



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Fondazione Italiana Linfomi. Ann Oncol 2014;25:1404-1410. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24799461>.

99. Pellicelli AM, Marignani M, Zoli V, et al. Hepatitis C virus-related B cell subtypes in non Hodgkin's lymphoma. World J Hepatol 2011;3:278-284. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22125661>.

100. Besson C, Canioni D, Lepage E, et al. Characteristics and outcome of diffuse large B-cell lymphoma in hepatitis C virus-positive patients in LNH 93 and LNH 98 Groupe d'Etude des Lymphomes de l'Adulte programs. J Clin Oncol 2006;24:953-960. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16418500>.

101. Ennishi D, Maeda Y, Niitsu N, et al. Hepatic toxicity and prognosis in hepatitis C virus-infected patients with diffuse large B-cell lymphoma treated with rituximab-containing chemotherapy regimens: a Japanese multicenter analysis. Blood 2010;116:5119-5125. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20823454>.

102. Nishikawa H, Tsudo M, Osaki Y. Clinical outcome in diffuse large B-cell lymphoma with hepatitis C virus infection in the rituximab era: a single center experience. Oncol Rep 2012;28:835-840. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22736295>.

103. Bacon BR, Gordon SC, Lawitz E, et al. Boceprevir for previously treated chronic HCV genotype 1 infection. N Engl J Med 2011;364:1207-1217. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21449784>.

104. Jacobson IM, McHutchison JG, Dusheiko G, et al. Telaprevir for previously untreated chronic hepatitis C virus infection. N Engl J Med 2011;364:2405-2416. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21696307>.

105. McHutchison JG, Everson GT, Gordon SC, et al. Telaprevir with peginterferon and ribavirin for chronic HCV genotype 1 infection. N Engl J Med 2009;360:1827-1838. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19403902>.

106. Poordad F, McCone J, Jr., Bacon BR, et al. Boceprevir for untreated chronic HCV genotype 1 infection. N Engl J Med 2011;364:1195-1206. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21449783>.

107. Ghany MG, Nelson DR, Strader DB, et al. An update on treatment of genotype 1 chronic hepatitis C virus infection: 2011 practice guideline by the American Association for the Study of Liver Diseases. Hepatology 2011;54:1433-1444. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21898493>.

108. Carson KR, Evens AM, Richey EA, et al. Progressive multifocal leukoencephalopathy after rituximab therapy in HIV-negative patients: a report of 57 cases from the Research on Adverse Drug Events and Reports project. Blood 2009;113:4834-4840. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19264918>.

109. Hopfinger G, Plessl A, Grisold W, et al. Progressive multifocal leukoencephalopathy after rituximab in a patient with relapsed follicular lymphoma and low IgG levels and a low CD4+ lymphocyte count. Leuk Lymphoma 2008;49:2367-2369. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19052987>.

110. Yokoyama H, Watanabe T, Maruyama D, et al. Progressive multifocal leukoencephalopathy in a patient with B-cell lymphoma during rituximab-containing chemotherapy: case report and review of the literature. Int J Hematol 2008;88:443-447. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18855101>.

111. Carson KR, Newsome SD, Kim EJ, et al. Progressive multifocal leukoencephalopathy associated with brentuximab vedotin therapy: a report of 5 cases from the Southern Network on Adverse Reactions (SONAR) project. Cancer 2014;120:2464-2471. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24771533>.

112. Castillo JJ, Kanan S, Meid K, et al. Rituximab intolerance in patients with Waldenstrom macroglobulinaemia. Br J Haematol 2016;174:645-648. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26523929>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

113. Chen LY, Shah R, Cwynarski K, et al. Ofatumumab is a feasible alternative anti-CD20 therapy in patients intolerant of rituximab. *Br J Haematol* 2019;184:462-465. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29363752>.

114. Davies A, Merli F, Mihaljevic B, et al. Efficacy and safety of subcutaneous rituximab versus intravenous rituximab for first-line treatment of follicular lymphoma (SABRINA): a randomised, open-label, phase 3 trial. *Lancet Haematol* 2017;4:e272-e282. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28476440>.

115. Lugtenburg P, Avivi I, Berenschot H, et al. Efficacy and safety of subcutaneous and intravenous rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisone in first-line diffuse large B-cell lymphoma: the randomized MabEase study. *Haematologica* 2017;102:1913-1922. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28935843>.

116. Pfeilschifter J, Diel IJ. Osteoporosis due to cancer treatment: pathogenesis and management. *J Clin Oncol* 2000;18:1570-1593. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10735906>.

117. Van Staa TP, Leufkens HG, Abenham L, et al. Use of oral corticosteroids and risk of fractures. June, 2000. *J Bone Miner Res* 2005;20:1487-1494; discussion 1486. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16149171>.

118. Svendsen P, Shekhrajka N, Nielsen KL, et al. R-CHOP(-like) treatment of diffuse large B-cell lymphoma significantly reduces CT-assessed vertebral bone density: a single center study of 111 patients. *Leuk Lymphoma* 2017;58:1105-1113. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27736260>.

119. Ratcliffe MA, Lanham SA, Reid DM, Dawson AA. Bone mineral density (BMD) in patients with lymphoma: the effects of chemotherapy, intermittent corticosteroids and premature menopause. *Hematol Oncol* 1992;10:181-187. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1398514>.

120. Cabanillas ME, Lu H, Fang S, Du XL. Elderly patients with non-Hodgkin lymphoma who receive chemotherapy are at higher risk for osteoporosis and fractures. *Leuk Lymphoma* 2007;48:1514-1521. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17701582>.

121. Paccou J, Merlusca L, Henry-Desailly I, et al. Alterations in bone mineral density and bone turnover markers in newly diagnosed adults with lymphoma receiving chemotherapy: a 1-year prospective pilot study. *Ann Oncol* 2014;25:481-486. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24401926>.

122. Shaikh AJ, Memon WA. Bone health in survivors of lymphoma, treated with high dose steroids - is there a need for clearer guidelines on bone care? *Asian Pac J Cancer Prev* 2011;12:1105-1106. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21790261>.

123. Ottanelli S. Prevention and treatment of bone fragility in cancer patient. *Clin Cases Miner Bone Metab* 2015;12:116-129. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26604936>.

124. Cosman F, de Beur SJ, LeBoff MS, et al. Clinician's Guide to Prevention and Treatment of Osteoporosis. *Osteoporos Int* 2014;25:2359-2381. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25182228>.

125. Kim SH, Lim SK, Hahn JS. Effect of pamidronate on new vertebral fractures and bone mineral density in patients with malignant lymphoma receiving chemotherapy. *Am J Med* 2004;116:524-528. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15063813>.

126. Westin JR, Thompson MA, Cataldo VD, et al. Zoledronic acid for prevention of bone loss in patients receiving primary therapy for lymphomas: a prospective, randomized controlled phase III trial. *Clin Lymphoma Myeloma Leuk* 2013;13:99-105. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23276888>.

127. Cummings SR, San Martin J, McClung MR, et al. Denosumab for prevention of fractures in postmenopausal women with osteoporosis. *N*



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Engl J Med 2009;361:756-765. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19671655>.

Discussion
update in
progress



Follicular Lymphoma

Overview

Follicular lymphoma (FL) is the most common subtype of indolent non-Hodgkin's lymphoma (NHL), and accounts for about 22% of all newly diagnosed cases of NHL.¹ About 90% of patients have a t(14;18) translocation, which juxtaposes *BCL2* with the *IGH* locus resulting in the deregulated expression of *BCL2*. Pathologic grading according to the number of centroblasts is considered to be a clinical predictor of outcome. Clinical outcomes for patients with FL1 and FL2 do not differ and this classification has been deemed unreliable. Therefore, in the WHO classification, FL1 and FL2 are grouped under a single grade (FL1-2). The WHO classification mandates stratifying FL3 into either 3A (centrocytes still present) or 3B (sheets of centroblasts).² Thus, FL is divided into three grades (FL1-2, FL3A, and FL3B) based on the number of centroblasts.

FL1-2 should be managed according to the treatment recommendations for FL. However, controversy exists regarding management of FL grade 3. There is no difference in survival outcomes between patients with FL3A and FL3B. However, FL3 with >50% diffuse component has an inferior survival outcome similar to that of diffuse large B-cell lymphoma (DLBCL).³ FL3B with *BCL2* translocation appears to have a clinical course similar to FL1-3A, whereas FL3B with cytogenetic abnormalities of *BCL6* is thought to be genetically more akin to germinal center type and is associated with a more aggressive clinical course.⁴ Since FL3B is rare, the clinical behavior of FL3 in most studies is based mainly on FL3A cases. Some may treat FL3A as FL and others may treat it as DLBCL. FL3B is commonly treated according to the treatment recommendations for DLBCL. Any area of DLBCL in a FL of any grade should be diagnosed and treated as a DLBCL.

FL with predominantly diffuse pattern is characterized by the absence the t(14;18) translocation, presence of 1p36 deletion, frequent inguinal

lymph node involvement, uniform CD23 expression, and low clinical stage.^{5,6} FL with 1p36 deletion is typically grade 1–2 and has a good prognosis. It should be managed according to treatment recommendations for FL.

Large B-cell lymphoma (LBCL) with *IRF4* rearrangement, which occurs most commonly in children and young adults, is considered as a new distinct provisional entity in the 2017 WHO classification.² These lymphomas are characterized by strong expression of IRF4/MUM1, and may have a follicular, follicular and diffuse, or pure diffuse growth pattern resembling FL3B or DLBCL.⁷ Patients typically present with Waldeyer's ring involvement and/or cervical lymph nodes and locally aggressive disease that responds well to chemotherapy with or without radiation therapy (RT). LBCL with *IRF4* rearrangement should be managed according to treatment recommendations for DLBCL.

Pediatric-type FL (PTFL) is considered as a definite entity in the 2017 WHO classification, since it is a biologically and clinically distinct indolent lymphoma and can also occur in adults.² PTFL is generally characterized by lack of *BCL2* rearrangement and t(14,18), which constitute the genetic hallmark of conventional FL cases seen in adults and *BCL6* rearrangement is also typically absent in PTFL.⁸⁻¹¹ *MAP2K1* and *TNFRSF14* mutations are the most frequent genetic alterations found in PTFL.¹²⁻¹⁴ The diagnosis and management of PTFL in adults is discussed on MS-18.

In the 2017 WHO classification, FL in situ (with the presence of FL-like B-cells in the germinal centers of morphologically reactive lymph nodes) has been renamed as in situ follicular neoplasia (ISFN).² It is characterized by the preservation of the lymph node architecture, with the incidental finding of focal strongly positive staining for *BCL2* (restricted to germinal centers) and CD10 in the involved follicles, and the detection of t(14;18) by fluorescence in situ hybridization (FISH).¹⁵⁻¹⁷ ISFN has been



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

reported in patients with prior FL or concurrent FL (at other sites), as well as in individuals with no known history of FL and the prevalence of ISFN in the general population has been reported to be 2%.^{15,16,18} Although uncommon (5%–6%), the development of or progression to overt lymphoma has been reported in patients reported to have ISFN.^{19,20} The significance or potential for malignancy of ISFN in patients without known FL remains unclear. These cases may potentially represent the tissue counterpart of circulating B-cells with t(14;18), or may represent a very early lesion with t(14;18) but without other genetic abnormalities that lead to overt lymphoma.^{19,21} The WHO classification recommends that a diagnosis of FL not be made in such cases, but that the report should suggest evaluation for the presence of FL elsewhere, and possibly close follow-up.

FL of the gastrointestinal tract (primary intestinal FL) is a recently described entity, which is common in the small intestine with the vast majority of cases occurring in the duodenum. The morphology, immunophenotype, and genetic features are similar to those of nodal FL. However, most patients have clinically indolent and localized disease. Survival appears to be excellent even without treatment. In many of the other extranodal sites, the morphology, immunophenotype, and genetic features are similar to those of nodal FL. Patients usually have localized disease and systemic relapses are rare.

Literature Search Criteria and Guidelines Update Methodology

Prior to the update of this version of the NCCN Guidelines® for B-Cell Lymphomas an electronic search of the PubMed database was performed to obtain key literature in FL published since the previous Guidelines update using the following search terms: follicular lymphoma and indolent lymphoma. The PubMed database was chosen as it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.²²

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.

The data from key PubMed articles deemed as relevant to these Guidelines and discussed by the panel have been included in this version of the Discussion section (eg, e-publications ahead of print, meeting abstracts). Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.

The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

Diagnosis

Immunophenotyping using immunohistochemistry (IHC) and/or flow cytometry for cell surface marker analysis is required to establish a diagnosis. FL has a characteristic immunophenotype, which includes CD20+, CD10+, BCL2+, CD23+/-, CD43-, CD5-, CCND1-, and BCL6+. Occasional cases of FL may be CD10- or BCL2-. The diagnosis is easily established on histologic grounds, but immunophenotyping is encouraged to distinguish FL from a nodular mantle cell lymphoma (MCL) or small lymphocytic lymphoma (SLL). Molecular genetic analysis to detect *BCL2* rearrangement; karyotype or FISH to identify t(14;18) and *BCL6*, *1p36*, and *IRF4/MUM1* gene rearrangements; and IHC for Ki-67 and *IRF4/MUM1* for FL grade 3, cyclin D1 may be useful under certain circumstances. Low-grade FL with a high proliferation index ($\geq 30\%$ as determined by Ki-67 immunostaining) has been shown to be associated with an aggressive clinical behavior.^{23,24} There is no evidence, however, that high Ki-67 should guide the selection of therapy.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

The Follicular Lymphoma International Prognostic Index (FLIPI) is a prognostic scoring system that divides patients into three distinct prognostic groups. FLIPI1 is based on age, Ann Arbor stage, and number of nodal sites involved; hemoglobin level; and serum lactate dehydrogenase (LDH) level.²⁵ FLIPI1 was developed in the pre-rituximab era but it has been shown to retain its prognostic significance in the modern chemoimmunotherapy era also.²⁶ FLIPI-2 was developed based on prospective collection of data from patients with newly diagnosed FL treated in the rituximab era and is based on age, hemoglobin level, longest diameter of largest involved lymph node, beta-2 microglobulin level, and bone marrow involvement.²⁷ FLIPI-2 was highly predictive of treatment outcomes, and separated patients into three distinct risk groups with 5-year progression-free survival (PFS) rates of 79%, 51%, and 20% for patients at low-risk, intermediate-risk and high-risk, respectively ($P < .00001$). The FLIPI-2 also defined distinct risk groups among the subgroup of patients treated with rituximab-containing regimens, with 5-year PFS rates of 98%, 88%, and 77% for patients at low-risk, intermediate-risk and high-risk, respectively ($P < .0001$).²⁷ Thus, FLIPI-2 may be useful for assessing prognosis in patients receiving active therapy with rituximab-based regimens. A simpler prognostic index incorporating only the baseline serum beta 2-microglobulin and LDH levels has been developed, which is easier to apply and also appears to be as predictive of outcomes as the FLIPI-1 and FLIPI-2 indices.²⁸

Although these index scores predict for prognosis, they have not yet been established as a means of selecting treatment options.

Workup

The initial workup should include a thorough physical examination with attention to node-bearing areas, and evaluation of performance status and constitutional symptoms. Laboratory assessments should include CBC with differential and a comprehensive metabolic panel, in addition to

measurements of serum LDH levels. Hepatitis B virus (HBV) testing prior to initiation of treatment is recommended for all patients who will receive anti-CD20 monoclonal antibody (MAB) based regimens due to increased risk of HBV reactivation. Measurement of uric acid, serum beta-2 microglobulin (necessary for calculation of FLIPI-2), and hepatitis C testing may be useful for certain cases.

Bone marrow biopsy with aspirate is essential to document clinical stage I–II disease. Adequate trephine biopsy (specimen ≥ 1.6 cm) should be obtained for initial staging evaluation, along with bone marrow aspiration.^{29,30} However, in patients with low burden indolent lymphoma with radiographic clinical stage III disease, an initial staging bone marrow evaluation can be deferred if observation is the initial option as it will not change the clinical recommendations. If radioimmunotherapy is considered, bilateral core biopsy is recommended; in such instances, the pathologist should provide the percent of overall cellular elements and the percent of cellular elements involved in the marrow.

Chest/abdomen/pelvic CT with contrast of diagnostic quality and/or whole-body PET/CT scan are recommended as part of initial diagnostic workup. PET/CT scan is essential if RT is planned for stage I–II disease. CT scan of the neck may also assist in defining the extent of local disease. Multigated acquisition (MUGA) scan or echocardiogram is recommended for patients being considered for treatment regimens containing anthracyclines.

Role of PET/CT in FL

Diagnosis and Workup

PET/CT scans are more accurate than CT scans alone in the detection of disease in patients with indolent lymphomas and several studies have reported high sensitivity (94%–98%) and specificity (88%–100%) for PET/CT scans for indolent lymphomas.³¹⁻³⁴



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

PET scans may also be useful in identifying occult sites of disease and detecting histologic transformation of FL to DLBCL.^{32,35,36} Standard fluorodeoxyglucose (FDG) uptake values (SUVs) on PET have been reported to be higher among transformed than non-transformed cases of indolent lymphomas.³² PET scans do not replace histologic confirmation of the diagnosis; however, if there are sites with discordantly high FDG avidity, these represent the most likely sites of histologic transformation. High SUVs on PET scan should raise the suspicion of histologic transformation to DLBCL, and can be used to direct the optimal site of biopsy for histologic confirmation.³⁶

Post-treatment Response Evaluation

The prognostic utility of post-treatment PET scans (PET negativity was associated with a longer PFS compared to PET positivity) has also been demonstrated in several studies.^{34,37-42}

In a retrospective study in patients with FL treated with RCHOP, PET/CT imaging was found to be more accurate than CT imaging in detecting both nodal and extranodal lesions at staging and in assessing response to treatment.³⁹ Post-treatment PET/CT negativity was associated with more favorable PFS outcomes; median PFS was 48 months among PET/CT-negative cases compared with 17 months for positive cases ($P < .001$).³⁹

An exploratory retrospective analysis of the prognostic value of post-induction PET/CT scans was conducted based on data obtained from the PRIMA trial of patients with FL.⁴⁰ Among patients with a post-induction PET/CT scan ($n = 122$), those with a positive PET/CT scan had a significantly inferior PFS rate compared with those who were PET negative (33% vs. 71% at 42 months; $P < .001$). The median PFS was 21 months and not reached, respectively. Among the patients randomized to observation, the 42-month PFS rate was 29% for PET/CT-positive patients compared with 68% in PET/CT-negative cases; median PFS was 30

months and 52 months, respectively. Among the patients randomized to rituximab maintenance ($n = 47$), PET/CT positivity was associated with inferior (but not statistically significant) PFS outcomes compared with PET/CT negativity (56% vs. 77% at 41 months); median PFS has not yet been reached in either the PET/CT-positive or PET/CT-negative subgroups. PET/CT status was also associated with OS outcomes in this exploratory analysis. Patients who were PET/CT positive after induction therapy had significantly inferior OS compared with PET/CT-negative patients (79% vs. 97% at 42 months; $P = .001$).

The prognostic value of PET imaging in patients with high-tumor-burden FL treated with first-line therapy with 6 cycles of RCHOP ($n = 121$; no rituximab maintenance administered) was also evaluated in a prospective study.⁴¹ PET scans were performed after 4 cycles of RCHOP (interim PET) and at the end of treatment (final PET), and all scans were centrally reviewed. A positive PET was defined as Deauville score 4 or higher. Among patients with an interim PET scan ($n = 111$), 76% had a PET-negative response. Among patients with a final PET ($n = 106$), 78% had a PET-negative response.⁴¹ At the end of treatment, nearly all patients (98%) who achieved a CR based on International Working Group (IWG) response criteria also achieved a PET-negative response. Interim PET was associated with significantly higher 2-year PFS (86% for PET negative vs. 61% for PET-positive; $P = .0046$) but no significant difference in terms of OS. Final PET-negativity was associated with both significantly higher 2-year PFS (87% vs. 51%; $P < .001$) and higher OS (100% vs. 88%; $P = .013$).⁴¹

These studies suggest that post-treatment imaging studies may have a role as a predictive factor for survival outcomes in patients with FL. End-of-treatment PET/CT scan is now considered a standard part of post-treatment response evaluation in patients with indolent lymphomas.



Further prospective studies are warranted to determine whether interim PET scans have a role in guiding post-induction therapeutic interventions.

Surveillance

Little data exist on the potential role of follow-up surveillance imaging for detection of relapse in patients with indolent NHL. In an early retrospective study, patients with stage I to stage III FL with a CR after induction were evaluated with clinical, laboratory, and imaging studies during routine follow up (n = 257).⁴³ Patients underwent CT scans of the abdomen and/or pelvis during follow-up visits. Follow-up was typically performed every 3 to 6 months for the first 5 years of treatment, and then annually thereafter. The median follow-up was 80 months. Relapse was detected in 78 patients, with the majority of relapses (77%) occurring within the first 5 years of treatment.⁴³ Eleven of the relapses were detected with abdominal and/or pelvic CT scans alone. Thus, in this analysis, 4% of patients with an initial CR had recurrence determined by routine surveillance with CT scans.⁴³

The role of surveillance PET scans in patients with lymphomas (Hodgkin lymphoma and NHL) with a CR after induction was also evaluated in a prospective study.⁴⁴ PET scans were performed every 6 months for the first 2 years after completion of induction, then annually thereafter. In the cohort of patients with indolent NHL (n = 78), follow-up PET scans detected true relapses in 10% of patients at 6 months, 12% at 12 months, 9% at 18 months, 9% at 24 months, 8% at 36 months, and 6% at 48 months. Among 13 patients who were PET-positive without a corresponding abnormality on CT scan, relapse was documented by biopsy in 8 patients. Of the 47 patients with PET-positive relapses, 38 patients were detected on CT and 30 patients were detected clinically at the same time as the PET. It is unclear whether this earlier detection of relapse in a proportion of patients translates to improved outcomes.

In the absence of evidence demonstrating improved survival outcomes with early PET detection of relapse, PET scans are not recommended for routine surveillance in patients who have achieved a CR after treatment.

Stage I–II FL

RT is an effective treatment option for patients with stage I–II disease resulting in long-term disease control rates of >90% with the 10-year PFS and OS rates ranging from 40% to 59% and 58% to 86%, respectively.^{45–49} The 15-year PFS outcomes were influenced by disease stage (66% for stage I vs. 26% for stage II disease) and maximal tumor size (49% for tumors <3 cm vs. 29% for ≥3 cm). The OS rate was not significantly different between extended-field RT compared with involved-field RT (49% vs. 40%, respectively) and the reduction in radiation field (RT of involved nodes only) did not impact PFS or OS outcomes.^{48,49} A more recent multicenter retrospective study conducted by the International Lymphoma Radiation Oncology Group also established RT alone (≥24 Gy) as a potentially curative treatment option for patients with untreated stage I to II FL (512 patients staged by PET/CT; 410 patients had stage I disease).⁵⁰ After a median follow-up of 52 months, the 5-year freedom-from-progression (FFP) and OS rates were 69% and 96% for the entire study population. The 5-year FFP rate was 74% for patients with stage I FL compared to 49% for those with stage II FL ($P < .0001$).

The addition of systemic therapy (rituximab, chemotherapy, or chemoimmunotherapy) to involved-field RT (IFRT) has been shown to improve failure-free survival (FFS) and PFS but did not impact OS in patients with early-stage disease.^{51–58}

Long-term outcomes from a study of RT in patients with early-stage grade 1–2 FL (with or without chemotherapy) reported a median OS of 19 years and a 15-year OS rate of 62%.⁵³ In this study, the majority of patients (74%) had stage I disease and 24% had received chemotherapy with RT,



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

which may have resulted in the higher OS rate reported compared with the other studies. In a small prospective randomized study of RT alone compared with RT with CHOP in patients with stage I low- or intermediate-grade NHL (n = 44), the addition of adjuvant CHOP to RT did not improve relapse-free survival (RFS) or OS in the subgroup of patients with early-stage low-grade NHL.⁵¹ The results of a multicenter observation study (n = 94) showed that the addition of rituximab to involved-site RT (ISRT) significantly prolongs PFS in patients with stage I–II disease.⁵⁷ The 10-year PFS rates were 65% and 51%, respectively ($P < .05$), for patients treated with rituximab + RT and RT alone. However, the OS rates were not significantly different between the treatment groups. In the phase III randomized trial of the Trans-Tasman Radiation Oncology Group (TROG 99.03) that evaluated IFRT vs. IFRT followed by CVP (cyclophosphamide, vincristine, and prednisone) plus rituximab (RCVP) in 150 patients with early-stage FL, at a median follow-up of 10 years, IFRT followed by RCVP was associated with superior PFS compared to IFRT alone.⁵⁸ The 10-year PFS rates were 59% and 41%, respectively. However, the 10-year OS rate was not significantly different between the treatment arms (95% and 87%, respectively).

In retrospective studies that have evaluated the outcomes of patients with stage I–II FL with various treatment approaches (observation, RT, rituximab monotherapy, chemoimmunotherapy, and combined modality with RT), no differences in OS outcomes were observed between the various management approaches.^{59,60} Carefully selected patients (requirement of large abdominal radiation field, advanced age, concern for xerostomia, or patient refusal) with stage I–II disease who did not receive immediate treatment had comparable outcomes to those who were treated with RT.⁶¹ Observation may be appropriate in circumstances where potential toxicity of ISRT outweighs potential clinical benefit.

Treatment of Stage I (<7 cm) or Contiguous Stage II (<7 cm) FL

ISRT (24–30 Gy) is recommended for patients with non-bulky disease. No further treatment is necessary for patients achieving CR or PR to ISRT. Clinical follow-up with a complete history and physical exam and laboratory assessment is recommended every 3 to 6 months for the first 5 years, and then annually (or as clinically indicated) thereafter. Surveillance imaging with CT scans can be performed no more than every 6 months up to the first 2 years following completion of treatment, and then no more than annually thereafter.

Patients with disease not responding to ISRT should be managed as described for stage III or IV disease.

Treatment of Stage I (≥7 cm) or Contiguous Stage II (≥7 cm), or Non-Contiguous Stage II FL

Anti-CD20 monoclonal antibody (MAB) with or without chemotherapy with or without ISRT is recommended for patients with bulky disease (stage I or contiguous stage II) or non-contiguous stage II disease. The addition of ISRT is included with a category 2B recommendation. Clinical follow-up with a complete history and physical exam, laboratory assessment, and surveillance imaging (as described above) are recommended for patients with CR or PR after completion of treatment.

Patients with non-responsive disease should be managed as described below for stage III or IV disease.

Treatment of Stage III–IV FL

Several prospective randomized trials conducted in the pre-rituximab era have failed to demonstrate a survival benefit with immediate treatment versus watchful waiting in patients with advanced stage, low-tumor-burden or asymptomatic FL.^{62–64} Results from more recent studies have shown that although immediate treatment with rituximab monotherapy results in



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

significantly longer median time to initiation of new therapy compared with watchful waiting, it does not improve OS.⁶⁵⁻⁶⁷

In an analysis of the data from the F2-study registry of the International Follicular Lymphoma Prognostic Factor Project, outcomes in a cohort of patients with asymptomatic, advanced stage, low-tumor-burden FL who were initially managed by a “watch and wait” approach (n = 107) were compared with the outcomes of patients who had low-tumor-burden, asymptomatic FL, but were initially treated with rituximab-containing regimens (n = 242).⁶⁵ The endpoint for the comparison was freedom from treatment failure (FFTF), which was defined as the time from diagnosis to one of the following events: progression during treatment, initiation of second-line therapy, relapse, or death from any cause. In the “watch and wait” cohort, initiation of first-line therapy was not considered an event for FFTF. The 4-year FFTF rate was 79% in the “watch and wait” cohort compared to 69% in the cohort of patients initially treated with rituximab-containing regimens; the difference was not significant after adjusting for differences in baseline disease factors between the cohorts. In addition, the 5-year OS rates were similar (87% vs. 88%, respectively).

The role of immediate treatment with rituximab (with or without additional rituximab maintenance) versus watchful waiting in patients with advanced stage, asymptomatic low-tumor-burden FL was evaluated in a randomized phase III intergroup trial (187 patients were assigned to watchful waiting; 192 patients were assigned to maintenance rituximab; and 84 patients were assigned to the induction therapy with rituximab, though this arm was closed early).⁶⁶ The primary endpoint of this trial was time to initiation of new treatment from randomization. Long-term follow-up data showed that there was a significant difference in the percentage of patients not needing new treatment at 3 years between the watchful waiting group and rituximab maintenance group (46% vs. 88%; $P < .0001$) or rituximab induction group (46% vs. 78%; $P < .0001$),

suggesting that rituximab monotherapy should be considered for patients with asymptomatic, advanced-stage, low-tumor-burden FL. However, no differences in OS were observed between the study arms. After a median follow-up of 50 months for the 3-arm study, the 3-year OS rates were 94%, 97%, and 96%, respectively, for the watchful waiting group, rituximab maintenance group, and induction rituximab group. The endpoint chosen for this trial, however, is also rather controversial considering that one arm of the trial involved initiation of early therapy; a more justifiable endpoint for this study could have been “time to initiation of second therapy.”

A recent report from the National LymphoCare Study that compared the outcomes of patients with stage II–IV FL who were managed with watchful waiting (n = 386), rituximab monotherapy (n = 296), or rituximab plus chemotherapy (n = 1072) as initial management strategy also confirmed that there was no significant difference in OS between the 3 different management strategies.⁶⁷ With a median follow-up of 8 years, the estimated 8-year OS rates were 74%, 67%, and 72%, respectively, for the watchful waiting group, rituximab monotherapy group, and rituximab plus chemotherapy group.

Collectively, findings from clinical studies suggest that immediate initial therapy with rituximab in patients not meeting the modified GELF criteria does not improve OS and outside the context of clinical trials, observation is still the standard practice for patients with advanced stage low-tumor-burden FL.⁶⁵⁻⁶⁷ Treatment should only be initiated when a patient presents with indications for treatment (based on the modified GELF criteria). The modified criteria used to determine treatment initiation include: symptoms attributable to FL (not limited to B symptoms); threatened end-organ function; cytopenia secondary to lymphoma; bulky disease (single mass >7 cm or 3 or more masses >3 cm), splenomegaly; and steady progression over at least 6 months. Suggested treatment regimens for first-line therapy are discussed below. Patients achieving CR



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

or PR to first-line therapy can either be observed or treated with optional consolidation or extended therapy. Options are discussed under “*First-line Consolidation Therapy* or *Extended Dosing*.”

The NCCN B-Cell Lymphomas Panel stratified all the regimens into 3 categories (based on the evidence, efficacy, toxicity, preexisting comorbidities, and in some cases access to certain agents): preferred regimens, other recommended regimens, and useful under certain circumstances.

First-line Therapy: Preferred Regimens

Chemoimmunotherapy with anti-CD20 Monoclonal Antibody (obinutuzumab or rituximab)

Bendamustine, CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) or CVP (cyclophosphamide, vincristine, and prednisone) with obinutuzumab or rituximab and lenalidomide + rituximab are included as preferred regimens, based on the results from phase III randomized trials (discussed below).

The multicenter randomized phase III study (StiL NHL1) that compared bendamustine and rituximab (BR) and RCHOP as first-line treatment for patients with indolent and MCLs showed that BR was superior to RCHOP in terms of PFS in all histologic subtypes.^{68,69} At a median follow-up of 45 months, the median PFS was 69 months and 31 months ($P < .0001$), respectively, for BR and RCHOP.⁶⁸ The ORR was similar between treatment arms (93% with BR; 91% with RCHOP), although the CR rate was significantly higher in the BR arm (40% vs. 30%; $P = .021$). BR was associated with less frequent grade 3 or 4 neutropenia (29% vs. 69%) or infections (any grade; 37% vs. 50%), whereas erythema (16% vs. 9%) and allergic skin reactions (15% vs. 6%) were more common with BR compared with RCHOP. The incidence of secondary malignancies was similar (8% with BR and 9% with RCHOP). However, the OS outcomes were not significantly different between treatment arms, even after

long-term follow-up; the estimated 10-year survival rates were 71% and 66%, respectively, for BR and RCHOP.⁶⁹

Another randomized phase III study (BRIGHT) demonstrated that BR was noninferior to RCHOP or RCVP (in terms of PFS) as first-line treatment of patients with indolent or MCL (224 patients were randomized to receive BR and 223 patients were randomized to receive RCHOP or RCVP).⁷⁰ At a median follow up of 5 years, the 5-year PFS rate for the overall study population was 66% and 56% ($P = .0025$), for BR and RCHOP/RCVP, respectively. The 5-year OS rate was not statistically different between the treatment groups and the incidence of opportunistic infections and secondary malignancies were slightly higher in patients treated with BR.⁷⁰

The phase III randomized trial (GALLIUM trial) compared the efficacy and safety of obinutuzumab compared to rituximab when used in combination with chemotherapy (bendamustine, CHOP, or CVP) in patients with previously untreated advanced stage FL (1202 patients were randomized [1:1] to receive obinutuzumab or rituximab in combination with chemotherapy).⁷¹ Patients with disease responding to induction chemoimmunotherapy received maintenance treatment with the same antibody for up to 2 years. However, this trial was not designed to compare the chemotherapy regimens. After a median follow-up of 34 months, a planned interim analysis showed that obinutuzumab-based chemoimmunotherapy was associated with significantly longer PFS and lower risk of progression and relapse than rituximab-based chemoimmunotherapy. The estimated 3-year PFS rates were 80% and 73%, respectively. However, response rates at the end of induction treatment were not significantly different between the two groups (88% for the obinutuzumab-based chemoimmunotherapy and 87% for rituximab-based chemoimmunotherapy) and OS was similar in both groups. In addition, grade 3 to 5 adverse events were higher with obinutuzumab than with rituximab (infections, 20% vs. 16%; neutropenia,



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

46% vs. 39%; and infusion-related reactions, 12% vs. 7%) and bendamustine was associated with higher rates of grade 3 to 5 infections and secondary cancers. Non-relapse-related fatal adverse events were also more common with bendamustine (6% in the obinutuzumab group and 4% in the rituximab group) than with CHOP or CVP (2% for both regimens when used in combination with obinutuzumab or rituximab).

The panel acknowledged that chemoimmunotherapy with anti-CD20 monoclonal antibody (obinutuzumab or rituximab) is an appropriate first-line therapy for patients with advanced stage FL requiring treatment. However, in the absence of data from randomized trials showing significant OS benefit for one chemoimmunotherapy regimen over another, the panel concluded that the available data are not strong enough to designate obinutuzumab-based chemoimmunotherapy as superior to rituximab-based chemoimmunotherapy. The panel consensus was to list all of the chemoimmunotherapy regimens as preferred regimens with category 2A recommendation.

The selection of a chemoimmunotherapy regimen should be highly individualized according to the patient's age, extent of disease, presence of comorbid conditions, and the goals of therapy. When choosing an initial therapy, care should be given to avoid excessively myelotoxic regimens in patients who may subsequently be candidates for HDT/ASCR. Chemoimmunotherapy regimens may be associated with risks for HBV reactivation, which can lead to hepatitis and hepatic failure. Therefore, prior to initiation of therapy, HBV testing (including HBsAg and HBcAb testing) should be performed for all patients; viral load should be monitored routinely for patients with positive test results. In addition, the use of empiric antiviral therapy or upfront prophylaxis should be incorporated into the treatment plan. In the GALLIUM study, there was an increased risk of mortality from opportunistic infections and secondary malignancies in patients receiving bendamustine, and the rates of severe infections were also higher with bendamustine than CHOP or CVP during

the maintenance and follow-up phases.⁷¹ The panel recommends that prophylaxis for pneumocystis jirovecii pneumonia (PJP) and varicella zoster virus (VZV) should be administered for patients receiving bendamustine.

Lenalidomide + Rituximab

In phase II studies, lenalidomide + rituximab has demonstrated activity in patients with previously untreated FL resulting in an ORR of 95% to 98% and 2-year PFS rates of 86% to 89%.^{72,73}

The results from the multicenter, international, randomized phase III study (RELEVANCE) showed that the efficacy of lenalidomide + rituximab was similar to that of chemotherapy + rituximab in patients with previously untreated advanced stage FL.⁷⁴ In this study, 1030 patients were randomized to receive lenalidomide + rituximab (n = 513) or chemotherapy + rituximab (investigator's choice of one the 3 regimens: RCHOP, RCVP, or BR; n = 517) followed by maintenance therapy with rituximab. The ORR (84% for lenalidomide + rituximab and 89% for chemotherapy + rituximab), CR rate at 120 weeks (48% and 53%, respectively; $P = .13$), and the interim 3-year PFS rate (77% and 78%, respectively) were similar in the two treatment groups. Lenalidomide + rituximab was associated with lower rates of grade 3 or 4 neutropenia (32% vs. 50%) and febrile neutropenia of any grade (2% vs. 7%) than chemotherapy + rituximab. Grade 3 or 4 cutaneous reactions were higher with lenalidomide + rituximab (7% vs. 1%).

Although the RELEVANCE trial (which was designed as a superiority trial) did not show that lenalidomide + rituximab was superior to chemotherapy + rituximab in terms of efficacy, this trial confirmed lenalidomide + rituximab as an effective alternative to chemoimmunotherapy for patients with previously untreated FL.⁷⁴ Based on the results of this study, the panel consensus was to include lenalidomide + rituximab as an option for preferred regimen with a category 2A recommendation.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

First-line Therapy: Other Recommended Regimens

Rituximab

Rituximab has been shown to induce high complete molecular response rate in patients with low burden FL and also has demonstrated single-agent activity in patients with previously untreated advanced stage FL.⁷⁵⁻⁷⁷ Rituximab should be considered as a first-line treatment for patients with low burden disease.

First-line Consolidation or Extended Therapy

Several studies have reported that prolonged administration of rituximab (or rituximab maintenance) significantly improved event-free survival (EFS) in chemotherapy-naïve patients responding to initial rituximab induction, although this benefit did not translate to OS advantage.⁷⁸⁻⁸¹

The phase III randomized trial (E4402 study; RESORT) compared rituximab maintenance versus rituximab retreatment in patients with previously untreated low burden FL responding to rituximab induction therapy.⁸² In this study, 289 patients were randomized to receive maintenance with rituximab or rituximab retreatment. Patients receiving rituximab retreatment were eligible for re-treatment at each disease progression until treatment failure. The primary endpoint of this trial was TTF. At a median follow-up of 5 years, the estimated TTF was similar for patients receiving rituximab maintenance and rituximab re-treatment (4 years for both treatment arms; $P = .54$). The 3-year freedom from cytotoxic therapy was 95% and 84%, respectively, for those receiving rituximab maintenance and rituximab re-treatment ($P = .03$). These results suggest that rituximab retreatment provides comparable disease control to rituximab maintenance in patients with low burden FL.

The phase III randomized PRIMA trial evaluated the role of rituximab maintenance in patients with FL responding to first-line chemoimmunotherapy.⁸³ In this study, patients with FL responding to first-line chemoimmunotherapy (RCVP, RCHOP, or RFCM) were

randomized to observation only or rituximab maintenance for 2 years ($n = 1018$). After a median follow-up of 36 months, the 3-year PFS rate was 75% in the rituximab maintenance arm and 58% in the observation arm ($P = .0001$). Two years after randomization, 71% of patients in the rituximab maintenance arm were in CR/CRu compared with 52% in the observation group. However, no significant difference was observed in OS between the two groups. Based on multivariate analysis, induction therapy with RCHOP or RFCM was one of the independent factors associated with improved PFS, suggesting that RCVP induction was not as beneficial in this study. Long-term follow-up data confirmed that rituximab maintenance is associated with significant PFS benefit over observation.⁸⁴ At 10 years, the estimated PFS rate was 51% in the rituximab maintenance arm and 35% in the observation arm. The OS estimates were identical (80%) in both treatment arms. The benefit of rituximab maintenance for PFS was significant irrespective of quality of response to chemoimmunotherapy.

The GALLIUM trial (discussed above) demonstrated that obinutuzumab-based chemoimmunotherapy followed by obinutuzumab maintenance resulted in significantly longer PFS than rituximab-based chemoimmunotherapy followed by rituximab maintenance.⁷¹ But the OS was not significantly different between the two treatment approaches.

Patients achieving CR or PR to first-line therapy can either be observed or treated with optional consolidation or extended therapy:

- Rituximab maintenance (one dose every 8 weeks up to 2 years) is included as an option with a category 1 recommendation for patients with high tumor burden treated with RCVP and RCHOP based on the PRIMA study.^{83,84} There are not enough data to support the use of rituximab maintenance in patients achieving CR after first-line therapy with BR. In a multicenter real-world analysis that evaluated maintenance rituximab versus observation following first-line treatment with BR, rituximab maintenance was associated



with a significant improvement in PFS only in patients achieving PR but not in patients achieving CR.⁸⁵

- Obinutuzumab maintenance is included with a category 2A recommendation based on the results of the GALLIUM trial.⁷¹
- Consolidation with rituximab should be considered if initially treated with single-agent rituximab.
- Ibritumomab tiuxetan is included with a category 2B recommendation based on the results of the First-line Indolent Trial (FIT).⁸⁶ Since this trial included only a small number of patients (14%) who received rituximab in combination with chemotherapy as induction therapy, the effects of consolidation with radioimmunotherapy (RIT) following rituximab-containing regimens cannot be fully evaluated.

Clinical follow-up with a complete physical exam and laboratory assessment (every 3–6 months for the first 5 years, and then annually [or as clinically indicated]) is recommended. Surveillance imaging with CT scans can be performed no more than every 6 months up to the first 2 years following completion of treatment, and then no more than annually (or as clinically indicated) thereafter.

Second-line and Subsequent Therapy

Frequently, patients with disease relapse or progression of disease (POD) after first-line therapy will benefit from a second period of observation. Considerations and indications for treatment of relapsed/refractory or progressive disease include, among other factors, the modified GELF criteria, which include: symptoms attributable to FL (not limited to B symptoms); threatened end-organ function; significant cytopenia secondary to lymphoma; bulky disease; splenomegaly; and steady progression over at least 6 months.

Progressive disease should be histologically documented to exclude transformation, especially in the presence of raising LDH levels, disproportional growth in one area, development of extranodal disease, or development of new constitutional symptoms. Areas of high SUV, especially in values in excess of 13, should raise suspicion for the presence of transformation.³⁶ However, a positive PET/CT scan does not replace a biopsy; rather, the results of the PET/CT scan should be used to direct the optimal site of biopsy for histologic confirmation to enhance the diagnostic yield from the biopsy.

POD ≤24 months of diagnosis and failure to achieve EFS at 12 months (EFS12) after initial treatment with chemoimmunotherapy have been identified as prognostic indicators of poor survival.⁸⁷⁻⁸⁹ In the National LymphoCare Study, the 5-year OS rate was 50% for patients with POD ≤2 years after first-line therapy with RCHOP compared with 90% for those with POD >2 years.⁸⁷ In a population-based analysis of relative survival of patients with FL compared to age-and-sex-matched controls (from U.S. and French data sets), the group of patients achieving EFS12 following initial management had similar OS outcomes compared to age-and-sex-matched general populations, whereas patients who failed to achieve EFS12 had lower subsequent OS.⁸⁸

Alternate non-cross-resistant chemoimmunotherapy regimens or the combination of lenalidomide and rituximab are options for second-line therapy for patients with relapsed/refractory disease. Duration of response to first-line therapy is an important factor in the selection of second-line therapy. Rituximab monotherapy may be appropriate for patients with late relapse as well, particularly if disease burden is low. Patients with POD ≤2 years after first-line therapy should be considered for treatment with lenalidomide-based regimens, novel approaches including clinical trials or referral for the consideration of HDT/ASCR.⁹⁰⁻⁹²



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Preferred Regimens

Chemoimmunotherapy with Anti-CD20 Monoclonal Antibody

The BR regimen has also demonstrated efficacy with acceptable toxicity in patients with relapsed or refractory indolent lymphomas.⁹³⁻⁹⁵ In a randomized phase III study of 230 patients with relapsed or refractory indolent lymphoma or MCL (114 patients assigned to BR and 105 patients assigned to fludarabine plus rituximab [FR]), BR was more effective than FR in terms of PFS.⁹⁵ At a median follow-up of 96 months, the median PFS was 34 months and 12 months, respectively, for BR and FR.

Obinutuzumab-based chemoimmunotherapy has also been evaluated in patients with relapsed or refractory FL.⁹⁶⁻⁹⁸

The safety and efficacy of CHOP plus obinutuzumab in patients with relapsed or refractory FL was demonstrated in a randomized study (56 patients were randomized to obinutuzumab in combination with CHOP or FC [fludarabine and cyclophosphamide]).⁹⁶ The ORRs were 96% and 93%, respectively, for patients treated with CHOP plus obinutuzumab and FC plus obinutuzumab. The corresponding CR rates were 39% and 50%, respectively. In the CHOP plus obinutuzumab group, 25% of patients with rituximab-refractory disease achieved CR and in the FC plus obinutuzumab, 30% achieved CR. All of the patients with rituximab-refractory disease achieved at least PR. FC plus obinutuzumab was associated with more adverse events than CHOP plus obinutuzumab.

The phase III randomized trial (GADOLIN) compared bendamustine + obinutuzumab versus bendamustine monotherapy in patients with rituximab-refractory indolent lymphoma (413 patients; 335 patients had FL; 164 patients were randomized to bendamustine + obinutuzumab; 171 patients were randomized to bendamustine monotherapy).^{97,98} Patients without POD in the bendamustine + obinutuzumab group received obinutuzumab maintenance. After a median follow-up of 32 months, the median PFS was significantly longer with bendamustine + obinutuzumab

than with bendamustine monotherapy (25 months vs. 14 months; $P < .001$) in patients with FL.¹⁰⁰ The median OS was not reached in the bendamustine + obinutuzumab group compared to 54 months for the bendamustine group. The most frequent grade ≥ 3 adverse events were neutropenia (33% for bendamustine plus obinutuzumab vs. 26% for bendamustine monotherapy), thrombocytopenia (11% vs. 16%), anemia (8% vs. 10%), and infusion-related reactions (11% vs. 6%).

Given the concerns for increased incidences of opportunistic infections and secondary malignancies in patients treated with BR as first-line therapy,^{71,99} BR is not recommended as an option for second-line therapy if it was previously used as first-line therapy.

Rituximab

Rituximab has also demonstrated single-agent activity in patients with relapsed or refractory disease resulting in a response rate of 48%.¹⁰⁰

Lenalidomide with or without Rituximab

In a phase II trial of patients with relapsed/refractory indolent NHL ($n = 43$), single-agent lenalidomide induced an ORR of 23% (7% CR).¹⁰¹ The ORR was 27% among the subgroup of 22 patients with FL. The median duration of response was longer than 16 months, and has not been reached. Median PFS for all patients was 4 months.

In the randomized phase II trial that evaluated the activity of lenalidomide monotherapy compared with lenalidomide plus rituximab (CALGB 50401 study) in patients with recurrent FL ($n = 91$), the ORR was significantly higher for patients receiving lenalidomide + rituximab than for those receiving lenalidomide monotherapy (76% vs. 53%; $P = .029$).¹⁰² With a median follow-up of 2.5 years, the median time to progression was 2 years and 1 year, respectively, for lenalidomide + rituximab and lenalidomide monotherapy. The incidences of grade ≥ 3 adverse events were similar in both treatment arms (53% for lenalidomide plus rituximab



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

vs. 58% for lenalidomide monotherapy). However, lenalidomide alone was associated with more treatment failures, with 22% of patients discontinuing treatment due to adverse events. The most common grade 3 or 4 toxicities included neutropenia (20% vs. 16%), fatigue (13% vs. 9%), and thrombosis (4% vs. 16%).

In the multicenter, double-blind, randomized phase III study (AUGMENT; n = 358; 295 patients had FL), lenalidomide + rituximab was superior to rituximab monotherapy in terms of PFS in all histologic subtypes of previously treated indolent lymphomas except MZL.¹⁰³ At median follow-up of 28 months, the median PFS was 39 months for lenalidomide + rituximab compared to 14 months for rituximab monotherapy ($P < .0001$) in the subgroup of patients with previously treated FL. The ORRs were 80% (35% CR) and 55% (20% CR), respectively ($P < .0001$). The estimated 2-year OS rates were 95% and 86%, respectively for lenalidomide + rituximab and rituximab monotherapy. Infections (63% vs. 49%), cutaneous reactions (32% vs. 12%), constipation (26% vs. 14%), thrombocytopenia (15% vs. 4%), and tumor flare reaction (11% vs. 1%) were more common with lenalidomide + rituximab with neutropenia (50% vs. 13%) and leukopenia (7% vs. 2%) being the most common grade 3 or 4 toxicities.

Other Recommended Regimens

Ibritumomab Tiuxetan

In a randomized phase III study of 143 patients with relapsed or refractory low-grade, follicular, or transformed lymphoma, ⁹⁰Y-ibritumomab tiuxetan produced a statistically and clinically significant higher ORR (80% vs. 56%) and CR rate (30% vs. 16%) compared with rituximab alone.¹⁰⁴ At a median follow-up of 44 months, median TTP (15 vs. 10 months) and duration of response (17 vs. 11 months) were longer for patients treated with ⁹⁰Y-ibritumomab compared with rituximab.¹⁰⁵

Phosphatidylinositol 3-kinase (PI3K) Inhibitors

Idelalisib, copanlisib, and duvelisib have shown promising results in the treatment of relapsed/refractory disease after 2 prior therapies.¹⁰⁶⁻¹¹⁰

Idelalisib and copanlisib have also been shown to be effective for high-risk patients with disease relapse ≤ 2 years after first-line therapy.^{109,111}

Based on the recent FDA approvals, idelalisib, copanlisib, and duvelisib are included as options for relapsed/refractory disease after 2 prior therapies. Evidence from clinical trials supporting this recommendation are discussed below.

Idelalisib

In a phase II multicenter, single-arm study of 125 patients with rituximab- and alkylating agent-refractory indolent NHL (72 patients had FL), idelalisib resulted in tumor reductions in 90% of the patients, with an ORR of 57% (6% CR).¹⁰⁶ The median duration of response and the median PFS were 13 months and 11 months, respectively. In a post hoc subgroup analysis of 72 patients with FL, the ORR was 56% (14% CR; 42% PR) and the overall disease control rate was 87%.¹⁰⁷ The PFS rate at 12 months and the estimated OS rate at 24 months were 43% and 70%, respectively. Grade ≥ 3 or greater toxicities included neutropenia (27%), thrombocytopenia (6%), elevated alanine transaminase (ALT) (13%), diarrhea (13%), and pneumonia (7%). In a post hoc analysis, the median PFS was 8 months for patients with POD within 12 months compared to 14 months for those with POD between 12 months and 24 months.¹¹¹ The ORR, however, was not significantly different between the 2 groups (71% and 45%, respectively, for patients with POD within 12 months and those with POD between 12 months and 24 months; $P = .18$).

Copanlisib

In a phase II study (CHRONOS-1) of 142 patients with relapsed/refractory indolent NHL after ≥ 2 prior lines of therapy, copanlisib resulted in an ORR



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

of 59% in the subgroup of 104 patients with FL (14% CR; 44% PR) and the median duration of response was 12 months. The median PFS and OS were 11 months and not reached, respectively.¹⁰⁸ The 2-year follow-up data also confirmed these results. The ORR was 61% (20% CR; 39% PR) and the median PFS and OS were 13 months and 43 months, respectively.¹⁰⁹ Among the 98 patients with FL and POD within 24 months, the ORR was 60% (22% CR; 38% PR), which was not significantly different from ORR of 59% for patients with POD after 24 months; however, the CR rate was higher in the POD <24 months group (22%) compared with 18% in the POD >24 months group.¹⁰⁹ Grade ≥3 toxicities included neutropenia (24%), thrombocytopenia (7%), hyperglycemia (41%), hypertension (24%), pneumonia (15%), and diarrhea (5%).

Duvelisib

In the phase II study (DYNAMO) evaluating the safety and efficacy of duvelisib in 129 patients with indolent NHL refractory to both rituximab and chemotherapy or radioimmunotherapy (RIT), duvelisib resulted in an ORR of 47% (42% for the subgroup of patients with FL).¹¹⁰ The estimated median duration of response and median PFS were 10 months and 9.5 months, respectively. Grade ≥3 toxicities included neutropenia (25%), anemia (15%), thrombocytopenia (12%), and diarrhea (15%).

Second-line Consolidation or Extended Dosing

Two large randomized trials have demonstrated a PFS advantage with rituximab maintenance over observation following second-line therapy.^{112,113}

In a prospective phase III randomized study by the GLSG, rituximab maintenance after second-line treatment with RFCM significantly prolonged duration of response in the subgroup of patients with recurring or refractory FL (n = 81); median PFS with rituximab maintenance was not reached compared with 26 months in the observation arm ($P = .035$).¹¹²

In a phase III randomized Intergroup trial (EORTC 20981) in patients with relapsed or resistant FL (n = 334) responding to CHOP or RCHOP induction therapy, rituximab maintenance significantly improved median PFS (4 years vs. 1 year; $P < .001$) compared with observation alone.¹¹³ This PFS benefit was observed regardless of the induction therapy employed (CHOP or RCHOP). With a median follow-up of 6 years, the 5-year OS rate was not significantly different between study arms (74% vs. 64%, respectively).¹¹³

In the GADOLIN study (discussed above), obinutuzumab maintenance following second-line therapy with bendamustine plus obinutuzumab improved PFS in patients with rituximab-refractory FL.^{97,98}

In a randomized study that evaluated the benefit of rituximab maintenance versus rituximab retreatment at progression in patients with indolent lymphomas previously treated with chemotherapy (n = 114), rituximab maintenance significantly improved PFS compared with rituximab retreatment (31 months vs. 7 months; $P = .007$).¹¹⁴ However, the duration of benefit was similar in both treatment groups (31 months vs. 27 months). Therefore, either approach (maintenance or rituximab retreatment at progression) could be beneficial for this patient population.

Patients achieving CR or PR to second-line or subsequent therapy can either be observed or treated with optional consolidation or extended therapy.

- Rituximab maintenance (one dose every 12 weeks up to 2 years) is included with a category 1 recommendation.^{112,113} However, the panel recognizes that the efficacy of rituximab maintenance in the second-line setting would likely be impacted by a patient's response to first-line maintenance with rituximab. The clinical benefit of rituximab maintenance in the second-line setting is likely



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

very minimal in patients with POD during or within 6 months of first-line maintenance with rituximab.

- Obinutuzumab maintenance (1 g every 8 weeks for a total of 12 doses) is preferred for patients with rituximab refractory disease, which includes POD on or within 6 months of prior rituximab therapy.^{97,98}

HDT/ASCR is an appropriate consolidative therapy for patients with second or third remission. HDT/ASCR as consolidation therapy has been shown to prolong OS and PFS in patients with relapsed or refractory disease.¹¹⁵⁻¹¹⁹ Allogeneic hematopoietic cell transplant (HCT) results in lower relapse rates than HDT/ASCR, but it is associated with high transplant-related mortality (TRM) rate.¹¹⁹⁻¹²¹ Allogeneic HCT may also be considered for highly selected patients.

Clinical follow-up with a complete physical exam and laboratory assessment (every 3 to 6 months for the first 5 years, and then annually [or as clinically indicated]) is recommended. Surveillance imaging with CT scans can be performed no more than every 6 months up to the first 2 years following completion of treatment, and then no more than annually (or as clinically indicated) thereafter.

Suggested Treatment Regimens for Elderly or Infirm Patients

As discussed earlier, rituximab monotherapy has demonstrated single-agent activity in patients with previously untreated low burden FL and advanced stage FL as well as in patients with relapsed/refractory disease.^{75,76,100} In a retrospective study of 75 patients with newly diagnosed FL, chlorambucil + rituximab resulted in an ORR of 97% (75% CR). The 5-year EFS and OS rates were 71% and 98%, respectively.¹²² Single-agent cyclophosphamide was associated with similar OS and CR rates compared with cyclophosphamide-based combination chemotherapy.¹²³

In an international phase II trial of 59 patients with stage III or IV FL (age >50 years; median age 66 years), ⁹⁰Y-ibritumomab tiuxetan when used as a first-line therapy resulted in an ORR of 87% (41% CR) at 6 months after therapy.¹²⁴ After a median follow-up of 31 months, the median PFS was 26 months and median OS had not been reached. The most common toxicities included grade 3 or 4 thrombocytopenia (48%; grade 4 in 7%) and neutropenia (32%; grade 4 in 17%). No grade 3 or 4 non-hematologic toxicities were reported. Grade 2 infections occurred in 20% and grade 2 GI toxicities in 10% of patients.

Rituximab monotherapy is the preferred treatment option for untreated as well as relapsed/refractory disease in elderly or infirm patients who are not able to tolerate any of the chemoimmunotherapy regimens recommended for first-line therapy. Ibritumomab tiuxetan and alkylating agent-based chemotherapy (cyclophosphamide or chlorambucil) with or without rituximab are included as alternative options. Ibritumomab tiuxetan is included with a category 2B recommendation. Selection of patients requires adequate marrow cellularity (>15%), <25% of bone marrow involvement, and platelet count >100,000. Referral to a tertiary care center is highly recommended for patients treated with prior ASCR.

Histologically Transformed Follicular Lymphoma

Histologic transformation of FL to DLBCL occurs in approximately 15% of patients with an estimated annual rate of 2% to 3% and is generally associated with a poor clinical outcome.¹²⁵⁻¹²⁷ In a pooled analysis of French and U.S. cohorts of patients with newly diagnosed FL, histologically transformed FL (TFL) after diagnosis was the leading cause death (77 of 140 deaths).¹²⁸ Recent studies that have evaluated the outcomes of patients with TFL in the rituximab era have reported improved survival in a subset of patients (eg, histologic transformation after early-stage FL and patients with no previous exposure to chemotherapy or rituximab prior to histologic transformation).^{127,129,130}



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Risk Factors

Advanced stage FL, high-risk FLIPI and IPI scores at diagnosis, elevated LDH, and B symptoms at diagnosis have been reported as risk factors for histologic transformation to DLBCL.^{125-127,129,131,132}

Early initiation of treatment at diagnosis relative to observation has been suggested to decrease the risk of transformation in some studies.^{127,129,131} However, in the randomized phase III intergroup trial that evaluated the role of immediate treatment with rituximab versus observation in patients with advanced stage, asymptomatic, low tumor burden FL, the trial also addressed whether early intensive rituximab therapy would change the risk of histologic transformation. No difference in time to transformation or incidence of histologic transformation was detected between the 3 groups, after a median follow-up of almost 4 years.⁶⁶

Treatment Options

There are no data from randomized studies to support the optimal treatment for patients with TFL, since clinical trials have often excluded this group of patients. Results from retrospective cohort studies suggest that rituximab monotherapy or in combination with anthracycline-based chemotherapy is associated with improved OS.^{127,129,132} RIT has also been shown to induce high response rates and durable remissions in patients with TFL.^{105,133}

In a series that reported the outcomes of 60 patients with biopsy-proven TFL, RCHOP was the most common treatment for TFL (n = 35; 59%) associated with a median OS of 50 months.¹²⁷ At a median follow-up of 60 months, the 5-year OS rate was 66% for patients with TFL treated with RCHOP, which was similar to the outcome of patients with de novo DLBCL treated with RCHOP in the validation cohort (5-year OS rate of 73%). In the National LymphoCare Study that included 147 pathologically confirmed and 232 clinically suspected cases of TFL, the majority of patients were treated with rituximab-based therapy (26% of patients

received rituximab monotherapy and 35% received chemotherapy with rituximab).¹²⁹ At a median follow-up of 7 years, the median PFS and OS were 12 months and 60 months, respectively, for patients with biopsy-proven histologic transformation to DLBCL. In another study that evaluated the outcomes of patients with FL who had histologic transformation after response to first-line chemoimmunotherapy in the PRIMA trial, the majority of patients with TFL received chemoimmunotherapy regimens recommended for DLBCL with a 5-year OS rate of 48% to 50%.¹³²

HDT/ASCR as consolidation therapy has been evaluated only in retrospective studies¹³⁴⁻¹⁴¹ with some series reporting survival benefit for patients who did proceed to transplant.^{137,139,141} Allogeneic HCT has been shown to benefit selected patients with disease relapse following HDT/ASCR, but is also associated with significant TRM.^{137,138,140,142} However, it should be noted that the efficacy of HDT/ASCR or allogeneic HCT in patients with TFL has not been confirmed in prospective controlled studies.

Histologic Transformation After Minimal Or No Prior Therapy

Anthracycline-based chemoimmunotherapy (with regimens recommended for first-line therapy for DLBCL, unless contraindicated) with or without ISRT is recommended for TFL without double-hit cytogenetics.^{4,5} Histologic transformation to high-grade B-cell lymphomas (HGBL) with translocations of *MYC* with either *BCL2* or *BCL6*, also called double- or triple-hit lymphoma, occurs in about 24% of cases and has been associated with an inferior survival compared to TFL without double-hit cytogenetics (2-year survival rates of 50% and 73%, respectively).¹⁴³ TFL with double-hit cytogenetics should be managed with more aggressive chemoimmunotherapy regimens as recommended for HGBL with translocations of *MYC* and *BCL2* and/or *BCL6*.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Consolidation therapy with HDT/ASCR with or without ISRT (if not previously given) or observation are included as treatment options for patients achieving CR or PR to initial treatment.^{137,139,141} If transformation is coexisting with extensive FL, rituximab maintenance should be considered for patients achieving CR. Additional systemic therapy \pm ISRT to induce CR should be considered for patients achieving PR, if proceeding to transplant. Repeat biopsy should be strongly considered prior to additional therapy for patients with PET-positive PR because PET positivity may represent post-treatment inflammation. Allogeneic HCT should be considered only in selected patients.

RIT with ibritumomab tiuxetan or ISRT (for localized residual disease and/or residual FDG-avid disease not previously irradiated) are included as alternative treatment options for patients achieving PR.^{105,133} RIT with ibritumomab tiuxetan or best supportive care are recommended for patients with non-responsive or progressive disease.^{105,133} Based on the FDA approval, chimeric antigen receptor (CAR) T-cell therapy (axicabtagene ciloleucel or tisagenlecleucel) is included as an option for patients who have received ≥ 2 prior chemoimmunotherapy regimens for indolent or transformed disease.¹⁴⁴⁻¹⁴⁶

Histologic Transformation After Multiple Lines of Prior Therapies

Enrollment in an appropriate clinical trial is the preferred option. In the absence of a suitable clinical trial, treatment options include chemoimmunotherapy (with regimens recommended for second-line therapy for DLBCL) with or without ISRT, RIT with ibritumomab tiuxetan, ISRT, or best supportive care.^{132,105,133} Based on the FDA approval, CAR T-cell therapy (axicabtagene ciloleucel or tisagenlecleucel) is included as an option for patients who have received ≥ 2 prior chemoimmunotherapy regimens for indolent or transformed disease.¹⁴⁴⁻¹⁴⁶

Consolidation therapy with HDT/ASCR with or without ISRT (if not previously given) or observation are included as treatment options for

patients achieving CR.^{137,139,141} Allogeneic HCT should be considered only in selected patients.

For patients achieving PR to initial therapy of TFL, treatment options include second-line regimens for DLBCL, allogeneic HCT with or without ISRT (only in the context of a clinical trial), CAR T-cell therapy (axicabtagene ciloleucel or tisagenlecleucel for patients who have received ≥ 2 prior chemoimmunotherapy regimens for indolent or transformed disease) if not previously given, or ISRT for localized residual disease and/or residual FDG-avid disease not previously irradiated. However, it should be noted that data on the efficacy of transplant in patients who have received CAR T-cell therapy are not available. HDT/ASCR is not recommended after CAR T-cell therapy. Allogeneic HCT could be considered but remains investigational.

Patients with non-responsive or progressive disease can be treated with any of the treatment options (not received previously), if they are candidates for additional therapy.

Pediatric-type Follicular Lymphoma in Adults

PTFL is generally characterized by lack of *BCL2* rearrangement and t(14,18), and *BCL6* rearrangement is also typically absent in PTFL.² *BCL2* expression may be observed in approximately 40% to 50% of cases, and *BCL6* expression can be seen in the majority of cases.⁸⁻¹¹


Histologically, PTFL is associated with large expansive follicles with a “starry sky” pattern, effacement of architecture, absence of diffuse area, high histologic grade (grade 3), and a high proliferation index.⁹⁻¹¹ In young patients with *BCL2*-negative localized disease, the diagnosis of PTFL may be considered. Analysis of *BCL6* rearrangement may be useful for evaluating the diagnosis of PTFL.



In adult patients without *BCL2* rearrangement but with high proliferation index, PTFL has a highly indolent disease course (stage I disease with no disease progression or relapse) compared to PTFL with *BCL2* rearrangement and/or low proliferation index (Ki-67 <30%; stage III or IV disease and majority of patients experience disease progression or recurrence).¹⁰

PTFL without *BCL2* rearrangements is generally associated with favorable prognosis with only rare instances of disease progression or relapse and is primarily managed with excision (preferred) or ISRT. RCHOP is recommended for patients with extensive local disease who are not candidates for excision or ISRT.¹⁴⁷

Restaging with PET/CT is recommended for patients treated with ISRT or RCHOP. No further treatment is necessary following excision or CR to ISRT or RCHOP. There are no data to support maintenance therapy. If patients have an excellent prognosis, no surveillance imaging is necessary. Patients with less responsive disease (<CR) should be managed as described for progressive FL.





NCCN Guidelines Version 2.2020

B-Cell Lymphomas

References

1. Al-Hamadani M, Habermann TM, Cerhan JR, et al. Non-Hodgkin lymphoma subtype distribution, geodemographic patterns, and survival in the US: A longitudinal analysis of the National Cancer Data Base from 1998 to 2011. *Am J Hematol* 2015;90:790-795. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26096944>.
2. Swerdlow SH, Harris NL, Jaffe ES, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Revised 4th ed. Lyon, France: IARC; 2017.
3. Hans CP, Weisenburger DD, Vose JM, et al. A significant diffuse component predicts for inferior survival in grade 3 follicular lymphoma, but cytologic subtypes do not predict survival. *Blood* 2003;101:2363-2367. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12424193>.
4. Katzenberger T, Ott G, Klein T, et al. Cytogenetic alterations affecting BCL6 are predominantly found in follicular lymphomas grade 3B with a diffuse large B-cell component. *Am J Pathol* 2004;165:481-490. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15277222>.
5. Katzenberger T, Kalla J, Leich E, et al. A distinctive subtype of t(14;18)-negative nodal follicular non-Hodgkin lymphoma characterized by a predominantly diffuse growth pattern and deletions in the chromosomal region 1p36. *Blood* 2009;113:1053-1061. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18978208>.
6. Siddiqi IN, Friedman J, Barry-Holson KQ, et al. Characterization of a variant of t(14;18) negative nodal diffuse follicular lymphoma with CD23 expression, 1p36/TNFRSF14 abnormalities, and STAT6 mutations. *Mod Pathol* 2016;29:570-581. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26965583>.
7. Salaverria I, Philipp C, Oschlies I, et al. Translocations activating IRF4 identify a subtype of germinal center-derived B-cell lymphoma affecting predominantly children and young adults. *Blood* 2011;118:139-147. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21487109>.
8. Lorbach RB, Shay-Seymore D, Moore J, et al. Clinicopathologic analysis of follicular lymphoma occurring in children. *Blood* 2002;99:1959-1964. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11877266>.
9. Oschlies I, Salaverria I, Mahn F, et al. Pediatric follicular lymphoma--a clinico-pathological study of a population-based series of patients treated within the Non-Hodgkin's Lymphoma--Berlin-Frankfurt-Munster (NHL-BFM) multicenter trials. *Haematologica* 2010;95:253-259. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19679882>.
10. Louissaint A, Jr., Ackerman AM, Dias-Santagata D, et al. Pediatric-type nodal follicular lymphoma: an indolent clonal proliferation in children and adults with high proliferation index and no BCL2 rearrangement. *Blood* 2012;120:2395-2404. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22855608>.
11. Liu Q, Salaverria I, Pittaluga S, et al. Follicular lymphomas in children and young adults: a comparison of the pediatric variant with usual follicular lymphoma. *Am J Surg Pathol* 2013;37:333-343. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23108024>.
12. Louissaint A, Jr., Schafernack KT, Geyer JT, et al. Pediatric-type nodal follicular lymphoma: a biologically distinct lymphoma with frequent MAPK pathway mutations. *Blood* 2016;128:1093-1100. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27325104>.
13. Schmidt J, Gong S, Marafioti T, et al. Genome-wide analysis of pediatric-type follicular lymphoma reveals low genetic complexity and recurrent alterations of TNFRSF14 gene. *Blood* 2016;128:1101-1111. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27257180>.
14. Schmidt J, Ramis-Zaldivar JE, Nadeu F, et al. Mutations of MAP2K1 are frequent in pediatric-type follicular lymphoma and result in ERK pathway activation. *Blood* 2017;130:323-327. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28533310>.
15. Cong P, Raffeld M, Teruya-Feldstein J, et al. In situ localization of follicular lymphoma: description and analysis by laser capture



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

microdissection. Blood 2002;99:3376-3382. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11964306>.

16. Sotomayor EA, Shah IM, Sanger WG, Mark HF. In situ follicular lymphoma with a 14;18 translocation diagnosed by a multimodal approach. Exp Mol Pathol 2007;83:254-258. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17482162>.

17. Carbone A, Gloghini A, Santoro A. In situ follicular lymphoma: pathologic characteristics and diagnostic features. Hematol Oncol 2012;30:1-7. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21560142>.

18. Henopp T, Quintanilla-Martinez L, Fend F, Adam P. Prevalence of follicular lymphoma in situ in consecutively analysed reactive lymph nodes. Histopathology 2011;59:139-142. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21771030>.

19. Jegalian AG, Eberle FC, Pack SD, et al. Follicular lymphoma in situ: clinical implications and comparisons with partial involvement by follicular lymphoma. Blood 2011;118:2976-2984. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21768298>.

20. Pillai RK, Surti U, Swerdlow SH. Follicular lymphoma-like B-cells of uncertain significance (in situ follicular lymphoma) may infrequently progress, but precedes follicular lymphoma, is associated with other overt lymphomas and mimics follicular lymphoma in flow cytometric studies. Haematologica 2013;98:1571-1580. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23831923>.

21. Bonzheim I, Salaverria I, Haake A, et al. A unique case of follicular lymphoma provides insights to the clonal evolution from follicular lymphoma in situ to manifest follicular lymphoma. Blood 2011;118:3442-3444. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21940830>.

22. U.S. National Library of Medicine Key MEDLINE® Indicators Available at: http://www.nlm.nih.gov/bsd/bsd_key.html.

23. Wang SA, Wang L, Hochberg EP, et al. Low histologic grade follicular lymphoma with high proliferation index: morphologic and clinical features. Am J Surg Pathol 2005;29:1490-1496. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16224216>.

24. Koster A, Tromp HA, Raemaekers JM, et al. The prognostic significance of the intra-follicular tumor cell proliferative rate in follicular lymphoma. Haematologica 2007;92:184-190. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17296567>.

25. Solal-Celigny P, Roy P, Colombat P, et al. Follicular lymphoma international prognostic index. Blood 2004;104:1258-1265. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15126323>.

26. Nooka AK, Nabhan C, Zhou X, et al. Examination of the follicular lymphoma international prognostic index (FLIPI) in the National LymphoCare study (NLCS): a prospective US patient cohort treated predominantly in community practices. Ann Oncol 2013;24:441-448. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23041589>.

27. Federico M, Bellei M, Marcheselli L, et al. Follicular lymphoma international prognostic index 2: a new prognostic index for follicular lymphoma developed by the international follicular lymphoma prognostic factor project. J Clin Oncol 2009;27:4555-4562. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19652063>.

28. Press OW, Unger JM, Rimsza LM, et al. A comparative analysis of prognostic factor models for follicular lymphoma based on a phase III trial of CHOP-rituximab versus CHOP + 131iodine--tositumomab. Clin Cancer Res 2013;19:6624-6632. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24130072>.

29. Bain BJ. Bone marrow trephine biopsy. J Clin Pathol 2001;54:737-742. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11577117>.

30. Bishop PW, McNally K, Harris M. Audit of bone marrow trephines. J Clin Pathol 1992;45:1105-1108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1479037>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

31. Blum RH, Seymour JF, Wirth A, et al. Frequent impact of [18F]fluorodeoxyglucose positron emission tomography on the staging and management of patients with indolent non-Hodgkin's lymphoma. Clin Lymphoma 2003;4:43-49. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12837154>.
32. Karam M, Novak L, Cyriac J, et al. Role of fluorine-18 fluoro-deoxyglucose positron emission tomography scan in the evaluation and follow-up of patients with low-grade lymphomas. Cancer 2006;107:175-183. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16721817>.
33. Wohrer S, Jaeger U, Kletter K, et al. 18F-fluoro-deoxy-glucose positron emission tomography (18F-FDG-PET) visualizes follicular lymphoma irrespective of grading. Ann Oncol 2006;17:780-784. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16497824>.
34. Bishu S, Quigley JM, Bishu SR, et al. Predictive value and diagnostic accuracy of F-18-fluoro-deoxy-glucose positron emission tomography treated grade 1 and 2 follicular lymphoma. Leuk Lymphoma 2007;48:1548-1555. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17701586>.
35. Schoder H, Noy A, Gonen M, et al. Intensity of 18fluorodeoxyglucose uptake in positron emission tomography distinguishes between indolent and aggressive non-Hodgkin's lymphoma. J Clin Oncol 2005;23:4643-4651. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15837966>.
36. Noy A, Schoder H, Gonen M, et al. The majority of transformed lymphomas have high standardized uptake values (SUVs) on positron emission tomography (PET) scanning similar to diffuse large B-cell lymphoma (DLBCL). Ann Oncol 2009;20:508-512. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19139176>.
37. Zinzani PL, Musuraca G, Alinari L, et al. Predictive role of positron emission tomography in the outcome of patients with follicular lymphoma. Clin Lymphoma Myeloma 2007;7:291-295. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17324337>.
38. Janikova A, Bolcak K, Pavlik T, et al. Value of [18F]fluorodeoxyglucose positron emission tomography in the management of follicular lymphoma: the end of a dilemma? Clin Lymphoma Myeloma 2008;8:287-293. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18854283>.
39. Le Dortz L, De Guibert S, Bayat S, et al. Diagnostic and prognostic impact of 18F-FDG PET/CT in follicular lymphoma. Eur J Nucl Med Mol Imaging 2010;37:2307-2314. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20717826>.
40. Trotman J, Fournier M, Lamy T, et al. Positron emission tomography-computed tomography (PET-CT) after induction therapy is highly predictive of patient outcome in follicular lymphoma: analysis of PET-CT in a subset of PRIMA trial participants. J Clin Oncol 2011;29:3194-3200. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21747087>.
41. Dupuis J, Berriolo-Riedinger A, Julian A, et al. Impact of [18F]Fluorodeoxyglucose Positron Emission Tomography Response Evaluation in Patients With High-Tumor Burden Follicular Lymphoma Treated With Immunochemotherapy: A Prospective Study From the Groupe d'Etudes des Lymphomes de l'Adulte and GOELAMS. J Clin Oncol 2012;30:4317-4322. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23109699>.
42. Luminari S, Biasoli I, Versari A, et al. The prognostic role of post-induction FDG-PET in patients with follicular lymphoma: a subset analysis from the FOLL05 trial of the Fondazione Italiana Linfomi (FIL). Ann Oncol 2014;25:442-447. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24412823>.
43. Oh YK, Ha CS, Samuels BI, et al. Stages I-III follicular lymphoma: role of CT of the abdomen and pelvis in follow-up studies. Radiology 1999;210:483-486. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10207433>.
44. Zinzani PL, Stefoni V, Tani M, et al. Role of [18F]fluorodeoxyglucose positron emission tomography scan in the follow-up of lymphoma. J Clin



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Oncol 2009;27:1781-1787. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19273712>.

45. Vaughan Hudson B, Vaughan Hudson G, MacLennan KA, et al. Clinical stage 1 non-Hodgkin's lymphoma: long-term follow-up of patients treated by the British National Lymphoma Investigation with radiotherapy alone as initial therapy. *Br J Cancer* 1994;69:1088-1093. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/8198975>.

46. Kamath SS, Marcus RB, Jr., Lynch JW, Mendenhall NP. The impact of radiotherapy dose and other treatment-related and clinical factors on in-field control in stage I and II non-Hodgkin's lymphoma. *Int J Radiat Oncol Biol Phys* 1999;44:563-568. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/10348285>.

47. Mac Manus MP, Hoppe RT. Is radiotherapy curative for stage I and II low-grade follicular lymphoma? Results of a long-term follow-up study of patients treated at Stanford University. *J Clin Oncol* 1996;14:1282-1290. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8648385>.

48. Wilder RB, Jones D, Tucker SL, et al. Long-term results with radiotherapy for Stage I-II follicular lymphomas. *Int J Radiat Oncol Biol Phys* 2001;51:1219-1227. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/11728680>.

49. Campbell BA, Voss N, Woods R, et al. Long-term outcomes for patients with limited stage follicular lymphoma: involved regional radiotherapy versus involved node radiotherapy. *Cancer* 2010;116:3797-3806. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/20564082>.

50. Brady JL, Binkley MS, Hajj C, et al. Definitive radiotherapy for localized follicular lymphoma staged by (18)F-FDG PET-CT: a collaborative study by ILROG. *Blood* 2019;133:237-245. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/30446493>.

51. Yahalom J, Varsos G, Fuks Z, et al. Adjuvant cyclophosphamide, doxorubicin, vincristine, and prednisone chemotherapy after radiation therapy in stage I low-grade and intermediate-grade non-Hodgkin

lymphoma. Results of a prospective randomized study. *Cancer* 1993;71:2342-2350. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/8453557>.

52. Seymour JF, Pro B, Fuller LM, et al. Long-term follow-up of a prospective study of combined modality therapy for stage I-II indolent non-Hodgkin's lymphoma. *J Clin Oncol* 2003;21:2115-2122. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/12775737>.

53. Guadagnolo BA, Li S, Neuberg D, et al. Long-term outcome and mortality trends in early-stage, Grade 1-2 follicular lymphoma treated with radiation therapy. *Int J Radiat Oncol Biol Phys* 2006;64:928-934. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16243446>.

54. Witzens-Harig M, Hensel M, Unterhalt M, Herfarth K. Treatment of limited stage follicular lymphoma with Rituximab immunotherapy and involved field radiotherapy in a prospective multicenter Phase II trial-MIR trial. *BMC Cancer* 2011;11:87. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/21352561>.

55. Janikova A, Bortlicek Z, Campr V, et al. Radiotherapy with rituximab may be better than radiotherapy alone in first-line treatment of early-stage follicular lymphoma: is it time to change the standard strategy? *Leuk Lymphoma* 2015;56:2350-2356. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25426666>.

56. Sancho JM, Garcia O, Mercadal S, et al. The long term follow-up of early stage follicular lymphoma treated with radiotherapy, chemotherapy or combined modality treatment. *Leuk Res* 2015;39:853-858. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26122511>.

57. Ruella M, Filippi AR, Bruna R, et al. Addition of Rituximab to Involved-Field Radiation Therapy Prolongs Progression-free Survival in Stage I-II Follicular Lymphoma: Results of a Multicenter Study. *Int J Radiat Oncol Biol Phys* 2016;94:783-791. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26972651>.

58. MacManus M, Fisher R, Roos D, et al. Randomized Trial of Systemic Therapy After Involved-Field Radiotherapy in Patients With Early-Stage



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Follicular Lymphoma: TROG 99.03. J Clin Oncol 2018;36:2918-2925. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29975623>.

59. Friedberg JW, Byrtek M, Link BK, et al. Effectiveness of first-line management strategies for stage I follicular lymphoma: analysis of the National LymphoCare Study. J Clin Oncol 2012;30:3368-3375. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22915662>.

60. Michallet AS, Lebras LL, Bauwens DD, et al. Early stage follicular lymphoma: what is the clinical impact of the first-line treatment strategy? J Hematol Oncol 2013;6:45. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23816219>.

61. Advani R, Rosenberg S, Horning S. Stage I and II follicular non-Hodgkin's lymphoma: long-term follow-up of no initial therapy. J Clin Oncol 2004;22:1454-1459. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15024027>.

62. Young RC, Longo DL, Glatstein E, et al. The treatment of indolent lymphomas: watchful waiting v aggressive combined modality treatment. Semin Hematol 1988;25:11-16. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2456618>.

63. Brice P, Bastion Y, Lepage E, et al. Comparison in low-tumor-burden follicular lymphomas between an initial no-treatment policy, prednimustine, or interferon alfa: a randomized study from the Groupe d'Etude des Lymphomes Folliculaires. Groupe d'Etude des Lymphomes de l'Adulte. J Clin Oncol 1997;15:1110-1117. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9060552>.

64. Ardeschna KM, Smith P, Norton A, et al. Long-term effect of a watch and wait policy versus immediate systemic treatment for asymptomatic advanced-stage non-Hodgkin lymphoma: a randomised controlled trial. Lancet 2003;362:516-522. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12932382>.

65. Solal-Celigny P, Bellei M, Marcheselli L, et al. Watchful waiting in low-tumor burden follicular lymphoma in the rituximab era: results of an

F2-study database. J Clin Oncol 2012;30:3848-3853. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23008294>.

66. Ardeschna KM, Qian W, Smith P, et al. Rituximab versus a watch-and-wait approach in patients with advanced-stage, asymptomatic, non-bulky follicular lymphoma: an open-label randomised phase 3 trial. The Lancet Oncology 2014;15:424-435. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24602760>.

67. Nastoupil LJ, Sinha R, Byrtek M, et al. Outcomes following watchful waiting for stage II-IV follicular lymphoma patients in the modern era. Br J Haematol 2016;172:724-734. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26729445>.

68. Rummel MJ, Niederle N, Maschmeyer G, et al. Bendamustine plus rituximab versus CHOP plus rituximab as first-line treatment for patients with indolent and mantle-cell lymphomas: an open-label, multicentre, randomised, phase 3 non-inferiority trial. Lancet 2013;381:1203-1210. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23433739>.

69. Rummel MJ, Maschmeyer G, Ganser A, et al. Bendamustine plus rituximab (B-R) versus CHOP plus rituximab (CHOP-R) as first-line treatment in patients with indolent lymphomas: Nine-year updated results from the StiL NHL1 study [abstract]. J Clin Oncol 2017;35 (15_suppl):Abstract 7501. Available at: http://ascopubs.org/doi/abs/10.1200/JCO.2017.35.15_suppl.7501.

70. Flinn IW, van der Jagt R, Kahl B, et al. First-Line Treatment of Patients With Indolent Non-Hodgkin Lymphoma or Mantle-Cell Lymphoma With Bendamustine Plus Rituximab Versus R-CHOP or R-CVP: Results of the BRIGHT 5-Year Follow-Up Study. J Clin Oncol 2019;37:984-991. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30811293>.

71. Marcus R, Davies A, Ando K, et al. Obinutuzumab for the First-Line Treatment of Follicular Lymphoma. N Engl J Med 2017;377:1331-1344. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28976863>.

72. Fowler NH, Davis RE, Rawal S, et al. Safety and activity of lenalidomide and rituximab in untreated indolent lymphoma: an open-label,



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

phase 2 trial. *Lancet Oncol* 2014;15:1311-1318. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25439689>.

73. Martin P, Jung SH, Pitcher B, et al. A phase II trial of lenalidomide plus rituximab in previously untreated follicular non-Hodgkin's lymphoma (NHL): CALGB 50803 (Alliance). *Ann Oncol* 2017;28:2806-2812. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28945884>.

74. Morschhauser F, Fowler NH, Feugier P, et al. Rituximab plus Lenalidomide in Advanced Untreated Follicular Lymphoma. *N Engl J Med* 2018;379:934-947. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30184451>.

75. Colombat P, Salles G, Brousse N, et al. Rituximab (anti-CD20 monoclonal antibody) as single first-line therapy for patients with follicular lymphoma with a low tumor burden: clinical and molecular evaluation. *Blood* 2001;97:101-106. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11133748>.

76. Witzig TE, Vukov AM, Habermann TM, et al. Rituximab therapy for patients with newly diagnosed, advanced-stage, follicular grade I non-Hodgkin's lymphoma: a phase II trial in the North Central Cancer Treatment Group. *J Clin Oncol* 2005;23:1103-1108. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15657404>.

77. Lockmer S, Ostenstad B, Hagberg H, et al. Chemotherapy-Free Initial Treatment of Advanced Indolent Lymphoma Has Durable Effect With Low Toxicity: Results From Two Nordic Lymphoma Group Trials With More Than 10 Years of Follow-Up. *J Clin Oncol* 2018;36:JCO1800262. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30285560>.

78. Ghielmini M, Schmitz SH, Cogliatti SB, et al. Prolonged treatment with rituximab in patients with follicular lymphoma significantly increases event-free survival and response duration compared with the standard weekly x 4 schedule. *Blood* 2004;103:4416-4423. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14976046>.

79. Martinelli G, Hsu Schmitz SF, Utiger U, et al. Long-term follow-up of patients with follicular lymphoma receiving single-agent rituximab at two

different schedules in trial SAKK 35/98. *J Clin Oncol* 2010;28:4480-4484. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20697092>.

80. Taverna C, Martinelli G, Hitz F, et al. Rituximab Maintenance for a Maximum of 5 Years After Single-Agent Rituximab Induction in Follicular Lymphoma: Results of the Randomized Controlled Phase III Trial SAKK 35/03. *J Clin Oncol* 2016;34:495-500. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26712227>.

81. Barta SK, Li H, Hochster HS, et al. Randomized phase 3 study in low-grade lymphoma comparing maintenance anti-CD20 antibody with observation after induction therapy: A trial of the ECOG-ACRIN Cancer Research Group (E1496). *Cancer* 2016;122:2996-3004. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27351685>.

82. Kahl BS, Hong F, Williams ME, et al. Rituximab extended schedule or re-treatment trial for low-tumor burden follicular lymphoma: eastern cooperative oncology group protocol e4402. *J Clin Oncol* 2014;32:3096-3102. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25154829>.

83. Salles G, Seymour JF, Offner F, et al. Rituximab maintenance for 2 years in patients with high tumour burden follicular lymphoma responding to rituximab plus chemotherapy (PRIMA): a phase 3, randomised controlled trial. *Lancet* 2011;377:42-51. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21176949>.

84. Salles GA, Seymour JF, Feugier P, et al. Long Term Follow-up of the PRIMA Study: Half of Patients Receiving Rituximab Maintenance Remain Progression Free at 10 Years [abstract]. *Blood* 2017;130 (Suppl 1):Abstract 486. Available at: http://www.bloodjournal.org/content/130/Suppl_1/486.abstract.

85. Hill BT, Nastoupil L, Winter AM, et al. Maintenance Rituximab or Observation after Frontline Treatment with Bendamustine-Rituximab (BR) for Follicular Lymphoma: A Real World Analysis Across 13 US Cancer Centers [abstract]. *Blood* 2017;130:Abstract 2779. Available at: http://www.bloodjournal.org/content/130/Suppl_1/2779.abstract.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

86. Morschhauser F, Radford J, Van Hoof A, et al. 90Yttrium-Ibritumomab Tiuxetan Consolidation of First Remission in Advanced-Stage Follicular Non-Hodgkin Lymphoma: Updated Results After a Median Follow-Up of 7.3 Years From the International, Randomized, Phase III First-Line Indolent Trial. *J Clin Oncol* 2013;31:1977-1983. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23547079>.

87. Casulo C, Byrtek M, Dawson KL, et al. Early Relapse of Follicular Lymphoma After Rituximab Plus Cyclophosphamide, Doxorubicin, Vincristine, and Prednisone Defines Patients at High Risk for Death: An Analysis From the National LymphoCare Study. *J Clin Oncol* 2015;33:2516-2522. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26124482>.

88. Maurer MJ, Bachy E, Ghesquieres H, et al. Early event status informs subsequent outcome in newly diagnosed follicular lymphoma. *Am J Hematol* 2016;91:1096-1101. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27465588>.

89. Casulo C, Le-Rademacher J, Dixon J, et al. Validation of POD24 As a Robust Early Clinical Endpoint of Poor Survival in Follicular Lymphoma: Results from the Follicular Lymphoma Analysis of Surrogacy Hypothesis (FLASH) Investigation Using Individual Data from 5,453 Patients on 13 Clinical Trials [abstract]. *Blood* 2017;130:Abstract 412. Available at: http://www.bloodjournal.org/content/130/Suppl_1/412.abstract.

90. Casulo C, Friedberg JW, Ahn KW, et al. Autologous Transplantation in Follicular Lymphoma with Early Therapy Failure: A National LymphoCare Study and Center for International Blood and Marrow Transplant Research Analysis. *Biol Blood Marrow Transplant* 2018;24:1163-1171. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29242111>.

91. Jurinovic V, Metzner B, Pfreundschuh M, et al. Autologous Stem Cell Transplantation for Patients with Early Progression of Follicular Lymphoma: A Follow-Up Study of 2 Randomized Trials from the German Low Grade Lymphoma Study Group. *Biol Blood Marrow Transplant* 2018;24:1172-1179. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29605716>.

92. Smith SM, Godfrey J, Ahn KW, et al. Autologous transplantation versus allogeneic transplantation in patients with follicular lymphoma experiencing early treatment failure. *Cancer* 2018;124:2541-2551. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29645093>.

93. Rummel MJ, Al-Batran SE, Kim SZ, et al. Bendamustine plus rituximab is effective and has a favorable toxicity profile in the treatment of mantle cell and low-grade non-Hodgkin's lymphoma. *J Clin Oncol* 2005;23:3383-3389. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15908650>.

94. Robinson KS, Williams ME, van der Jagt RH, et al. Phase II multicenter study of bendamustine plus rituximab in patients with relapsed indolent B-cell and mantle cell non-Hodgkin's lymphoma. *J Clin Oncol* 2008;26:4473-4479. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18626004>.

95. Rummel M, Kaiser U, Balser C, et al. Bendamustine plus rituximab versus fludarabine plus rituximab for patients with relapsed indolent and mantle-cell lymphomas: a multicentre, randomised, open-label, non-inferiority phase 3 trial. *Lancet Oncol* 2016;17:57-66. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26655425>.

96. Radford J, Davies A, Cartron G, et al. Obinutuzumab (GA101) plus CHOP or FC in relapsed/refractory follicular lymphoma: results of the GAUDI study (BO21000). *Blood* 2013;122:1137-1143. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23843495>.

97. Sehn LH, Chua N, Mayer J, et al. Obinutuzumab plus bendamustine versus bendamustine monotherapy in patients with rituximab-refractory indolent non-Hodgkin lymphoma (GADOLIN): a randomised, controlled, open-label, multicentre, phase 3 trial. *Lancet Oncol* 2016;17:1081-1093. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27345636>.

98. Cheson BD, Chua N, Mayer J, et al. Overall Survival Benefit in Patients With Rituximab-Refractory Indolent Non-Hodgkin Lymphoma Who Received Obinutuzumab Plus Bendamustine Induction and Obinutuzumab Maintenance in the GADOLIN Study. *J Clin Oncol*



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

2018;36:2259-2266. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29584548>.

99. Flinn I, Jagt Rvd, Chang JE, et al. First-line treatment of iNHL or MCL patients with BR or R-CHOP/R-CVP: Results of the BRIGHT 5-year follow-up study [abstract]. J Clin Oncol 2017;35:Abstract 7500. Available at: http://ascopubs.org/doi/abs/10.1200/JCO.2017.35.15_suppl.7500.

100. McLaughlin P, Grillo-Lopez AJ, Link BK, et al. Rituximab chimeric anti-CD20 monoclonal antibody therapy for relapsed indolent lymphoma: half of patients respond to a four-dose treatment program. J Clin Oncol 1998;16:2825-2833. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9704735>.

101. Witzig TE, Wiernik PH, Moore T, et al. Lenalidomide oral monotherapy produces durable responses in relapsed or refractory indolent non-Hodgkin's Lymphoma. J Clin Oncol 2009;27:5404-5409. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19805688>.

102. Leonard JP, Jung SH, Johnson J, et al. Randomized Trial of Lenalidomide Alone Versus Lenalidomide Plus Rituximab in Patients With Recurrent Follicular Lymphoma: CALGB 50401 (Alliance). J Clin Oncol 2015;33:3635-3640. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26304886>.

103. Leonard JP, Trneny M, Izutsu K, et al. AUGMENT: A Phase III Study of Lenalidomide Plus Rituximab Versus Placebo Plus Rituximab in Relapsed or Refractory Indolent Lymphoma. J Clin Oncol 2019;37:1188-1199. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30897038>.

104. Witzig TE, Gordon LI, Cabanillas F, et al. Randomized controlled trial of yttrium-90-labeled ibritumomab tiuxetan radioimmunotherapy versus rituximab immunotherapy for patients with relapsed or refractory low-grade, follicular, or transformed B-cell non-Hodgkin's lymphoma. J Clin Oncol 2002;20:2453-2463. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12011122>.

105. Gordon LI, Witzig T, Molina A, et al. Yttrium 90-labeled ibritumomab tiuxetan radioimmunotherapy produces high response rates and durable remissions in patients with previously treated B-cell lymphoma. Clin Lymphoma 2004;5:98-101. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15453924>.

106. Gopal AK, Kahl BS, de Vos S, et al. PI3Kdelta inhibition by idelalisib in patients with relapsed indolent lymphoma. N Engl J Med 2014;370:1008-1018. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24450858>.

107. Salles G, Schuster SJ, de Vos S, et al. Efficacy and safety of idelalisib in patients with relapsed, rituximab- and alkylating agent-refractory follicular lymphoma: a subgroup analysis of a phase 2 study. Haematologica 2017;102:e156-e159. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27979923>.

108. Dreyling M, Santoro A, Mollica L, et al. Phosphatidylinositol 3-Kinase Inhibition by Copanlisib in Relapsed or Refractory Indolent Lymphoma. J Clin Oncol 2017;35:3898-3905. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28976790>.

109. Dreyling M, Santoro A, Mollica L, et al. Long-Term Efficacy and Safety from the Copanlisib CHRONOS-1 Study in Patients with Relapsed or Refractory Indolent B-Cell Lymphoma [abstract]. Blood 2018;132:Abstract 1595. Available at: http://www.bloodjournal.org/content/132/Suppl_1/1595.abstract.

110. Flinn IW, Miller CB, Ardeshtna KM, et al. DYNAMO: A Phase II Study of Duvelisib (IPI-145) in Patients With Refractory Indolent Non-Hodgkin Lymphoma. J Clin Oncol 2019;37:912-922. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30742566>.

111. Gopal AK, Kahl BS, Flowers CR, et al. Idelalisib is effective in patients with high-risk follicular lymphoma and early relapse after initial chemoimmunotherapy. Blood 2017;129:3037-3039. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28325864>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

112. Forstpointner R, Unterhalt M, Dreyling M, et al. Maintenance therapy with rituximab leads to a significant prolongation of response duration after salvage therapy with a combination of rituximab, fludarabine, cyclophosphamide, and mitoxantrone (R-FCM) in patients with recurring and refractory follicular and mantle cell lymphomas: Results of a prospective randomized study of the German Low Grade Lymphoma Study Group (GLSG). *Blood* 2006;108:4003-4008. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16946304>.

113. van Oers MHJ, Van Glabbeke M, Giurgea L, et al. Rituximab maintenance treatment of relapsed/resistant follicular non-Hodgkin's lymphoma: long-term outcome of the EORTC 20981 phase III randomized intergroup study. *J Clin Oncol* 2010;28:2853-2858. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20439641>.

114. Hainsworth JD, Litchy S, Shaffer DW, et al. Maximizing therapeutic benefit of rituximab: maintenance therapy versus re-treatment at progression in patients with indolent non-Hodgkin's lymphoma--a randomized phase II trial of the Minnie Pearl Cancer Research Network. *J Clin Oncol* 2005;23:1088-1095. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15657401>.

115. Schouten HC, Qian W, Kvaloy S, et al. High-dose therapy improves progression-free survival and survival in relapsed follicular non-Hodgkin's lymphoma: results from the randomized European CUP trial. *J Clin Oncol* 2003;21:3918-3927. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/14517188>.

116. Rohatiner AZS, Nadler L, Davies AJ, et al. Myeloablative therapy with autologous bone marrow transplantation for follicular lymphoma at the time of second or subsequent remission: long-term follow-up. *J Clin Oncol* 2007;25:2554-2559. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17515573>.

117. Sebban C, Brice P, Delarue R, et al. Impact of rituximab and/or high-dose therapy with autotransplant at time of relapse in patients with follicular lymphoma: a GELA study. *J Clin Oncol* 2008;26:3614-3620. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18559872>.

118. Le Gouill S, De Guibert S, Planche L, et al. Impact of the use of autologous stem cell transplantation at first relapse both in naive and previously rituximab exposed follicular lymphoma patients treated in the GELA/GOELAMS FL2000 study. *Haematologica* 2011;96:1128-1135. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21486862>.

119. Evens AM, Vanderplas A, LaCasce AS, et al. Stem cell transplantation for follicular lymphoma relapsed/refractory after prior rituximab: a comprehensive analysis from the NCCN lymphoma outcomes project. *Cancer* 2013;119:3662-3671. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23921646>.

120. Peniket AJ, Ruiz de Elvira MC, Taghipour G, et al. An EBMT registry matched study of allogeneic stem cell transplants for lymphoma: allogeneic transplantation is associated with a lower relapse rate but a higher procedure-related mortality rate than autologous transplantation. *Bone Marrow Transplant* 2003;31:667-678. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12692607>.

121. van Besien K, Loberiza FR, Bajorunaite R, et al. Comparison of autologous and allogeneic hematopoietic stem cell transplantation for follicular lymphoma. *Blood* 2003;102:3521-3529. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12893748>.

122. Martinelli G, Montoro J, Vanazzi A, et al. Chlorambucil-rituximab as first-line therapy in patients affected by follicular non-Hodgkin's lymphoma: a retrospective single-centre study. *Hematol Oncol* 2015;33:129-135. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25047267>.

123. Peterson BA, Petroni GR, Frizzera G, et al. Prolonged single-agent versus combination chemotherapy in indolent follicular lymphomas: a study of the cancer and leukemia group B. *J Clin Oncol* 2003;21:5-15. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12506163>.

124. Scholz CW, Pinto A, Linkesch W, et al. (90)Yttrium-ibritumomab-tiuxetan as first-line treatment for follicular lymphoma: 30 months of follow-up data from an international multicenter phase II clinical trial. *J Clin Oncol* 2013;31:308-313. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23233718>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

125. Al-Tourah AJ, Gill KK, Chhanabhai M, et al. Population-based analysis of incidence and outcome of transformed non-Hodgkin's lymphoma. *J Clin Oncol* 2008;26:5165-5169. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18838711>.

126. Conconi A, Ponzio C, Lobetti-Bodoni C, et al. Incidence, risk factors and outcome of histological transformation in follicular lymphoma. *Br J Haematol* 2012;157:188-196. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22348437>.

127. Link BK, Maurer MJ, Nowakowski GS, et al. Rates and outcomes of follicular lymphoma transformation in the immunochemotherapy era: a report from the University of Iowa/MayoClinic Specialized Program of Research Excellence Molecular Epidemiology Resource. *J Clin Oncol* 2013;31:3272-3278. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23897955>.

128. Sarkozy C, Maurer MJ, Link BK, et al. Cause of Death in Follicular Lymphoma in the First Decade of the Rituximab Era: A Pooled Analysis of French and US Cohorts. *J Clin Oncol* 2019;37:144-152. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30481079>.

129. Wagner-Johnston ND, Link BK, Byrtek M, et al. Outcomes of transformed follicular lymphoma in the modern era: a report from the National LymphoCare Study (NLCS). *Blood* 2015;126:851-857. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26105149>.

130. Gleeson M, Hawkes EA, Peckitt C, et al. Outcomes for transformed follicular lymphoma in the rituximab era: the Royal Marsden experience 2003-2013. *Leuk Lymphoma* 2017;58:1805-1813. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27931133>.

131. Montoto S, Davies AJ, Matthews J, et al. Risk and clinical implications of transformation of follicular lymphoma to diffuse large B-cell lymphoma. *J Clin Oncol* 2007;25:2426-2433. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17485708>.

132. Sarkozy C, Trneny M, Xerri L, et al. Risk Factors and Outcomes for Patients With Follicular Lymphoma Who Had Histologic Transformation

After Response to First-Line Immunochemotherapy in the PRIMA Trial. *J Clin Oncol* 2016;34:2575-2582. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27298402>.

133. Mei M, Wondergem MJ, Palmer JM, et al. Autologous transplantation for transformed non-Hodgkin lymphoma using an yttrium-90 ibritumomab tiuxetan conditioning regimen. *Biol Blood Marrow Transplant* 2014;20:2072-2075. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25079874>.

134. Smith SD, Bolwell BJ, Advani AS, et al. High rate of survival in transformed lymphoma after autologous stem cell transplant: pathologic analysis and comparison with de novo diffuse large B-cell lymphoma. *Leuk Lymphoma* 2009;50:1625-1631. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19672775>.

135. Ban-Hoefen M, Kelly JL, Bernstein SH, et al. High-dose therapy and autologous stem cell transplant for transformed non-Hodgkin lymphoma in the rituximab era. *Leuk Lymphoma* 2012;53:830-835. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22023518>.

136. Reddy N, Oluwole O, Greer JP, et al. Superior long-term outcome of patients with early transformation of non-Hodgkin lymphoma undergoing stem cell transplantation. *Clin Lymphoma Myeloma Leuk* 2012;12:406-411. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22981964>.

137. Ban-Hoefen M, Vanderplas A, Crosby-Thompson AL, et al. Transformed non-Hodgkin lymphoma in the rituximab era: analysis of the NCCN outcomes database. *Br J Haematol* 2013;163:487-495. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24111533>.

138. Villa D, Crump M, Panzarella T, et al. Autologous and allogeneic stem-cell transplantation for transformed follicular lymphoma: a report of the Canadian blood and marrow transplant group. *J Clin Oncol* 2013;31:1164-1171. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23401459>.



NCCN Guidelines Version 2.2020 B-Cell Lymphomas

139. Villa D, Crump M, Keating A, et al. Outcome of patients with transformed indolent non-Hodgkin lymphoma referred for autologous stem-cell transplantation. *Ann Oncol* 2013;24:1603-1609. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23425946>.

140. Wirk B, Fenske TS, Hamadani M, et al. Outcomes of hematopoietic cell transplantation for diffuse large B cell lymphoma transformed from follicular lymphoma. *Biol Blood Marrow Transplant* 2014;20:951-959. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24641828>.

141. Madsen C, Pedersen MB, Vase MO, et al. Outcome determinants for transformed indolent lymphomas treated with or without autologous stem-cell transplantation. *Ann Oncol* 2015;26:393-399. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25411416>.

142. Clavert A, Le Gouill S, Brissot E, et al. Reduced-intensity conditioning allogeneic stem cell transplant for relapsed or transformed aggressive B-cell non-Hodgkin lymphoma. *Leuk Lymphoma* 2010;51:1502-1508. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20583964>.

143. Kridel R, Mottok A, Farinha P, et al. Cell of origin of transformed follicular lymphoma. *Blood* 2015;126:2118-2127. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26307535>.

144. Neelapu SS, Locke FL, Bartlett NL, et al. Axicabtagene Ciloleucel CAR T-Cell Therapy in Refractory Large B-Cell Lymphoma. *N Engl J Med* 2017;377:2531-2544. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29226797>.

145. Locke FL, Ghobadi A, Jacobson CA, et al. Long-term safety and activity of axicabtagene ciloleucel in refractory large B-cell lymphoma (ZUMA-1): a single-arm, multicentre, phase 1-2 trial. *Lancet Oncol* 2019;20:31-42. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30518502>.

146. Schuster SJ, Bishop MR, Tam CS, et al. Tisagenlecleucel in Adult Relapsed or Refractory Diffuse Large B-Cell Lymphoma. *N Engl J Med* 2019;380:45-56. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30501490>.

147. Kumar R, Galardy PJ, Dogan A, et al. Rituximab in combination with multiagent chemotherapy for pediatric follicular lymphoma. *Pediatr Blood Cancer* 2011;57:317-320. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21462303>.



Marginal Zone Lymphomas

Marginal zone lymphomas (MZLs) originate in the marginal zone of lymphoid follicles found in the mucosa-associated lymphoid tissues (MALT), spleen, and lymph nodes.¹ Extranodal MZLs of MALT (MALT lymphomas), nodal MZL (NMZL), and splenic MZL (SMZL) are the three distinct subtypes of MZLs. In a Surveillance, Epidemiology, and End Results (SEER) database analysis that assessed the incidence rates (IRs) of different subtypes of MZLs in the United States (2001–2009), NMZL and extranodal MZL were diagnosed in 4,081 (IR = 5.7) and 8,821 (IR = 12.3) individuals, respectively.² The most common sites were stomach (IR = 3.8), spleen (IR = 1.6), ocular adnexa (IR = 1.4), and lung, skin, and salivary glands (IR = 0.9–1.0).

The etiology of MZLs has been associated with chronic immune stimulation due to infectious agents or inflammation. *Helicobacter pylori* (*H. pylori*) has been implicated in the pathogenesis of gastric MALT lymphoma and other infectious agents such as *Chlamydia psittaci* (*C. psittaci*), *Campylobacter jejuni* (*C. jejuni*), and *Borrelia burgdorferi* (*B. burgdorferi*) have also been implicated in the pathogenesis of MZLs.³ MZLs are also characterized by a high prevalence of hepatitis C virus (HCV) infection and HCV seroprevalence has been reported in about 22% to 35% of patients with NMZL, SMZL, and nongastric MALT lymphomas.^{4,5}

MZLs are also characterized by clinical and pathologic features that overlap with Waldenström's macroglobulinemia/lymphoplasmacytic lymphoma (WM/LPL).⁶ Recent studies have confirmed that the *MYD88* L265P somatic mutational analysis could be useful in differentiating WM/LPL from other B-cell malignancies with overlapping clinical and pathologic features.⁷⁻¹⁰ In a retrospective analysis of 123 patients with a diagnosis of MZLs and WM/LPL, *MYD88* mutation was found in 67% of patients with WM/LPL (18 of 27) compared to 4% of patients with SMZL (2 out of 53), 7% of patients with MALT lymphomas (2 out of 28), and 0% of

patients with NMZL.⁸ Immunoglobulin heavy chain variable (*IGHV*) gene sequencing clearly distinguished SMZL and WM/LPL. SMZL was characterized by overrepresentation of *IGHV1-2* gene rearrangements with low or intermediate mutation rates, whereas WM/LPL was associated with overrepresentation of *IGHV3-23* rearrangements and high mutation rates.⁸ In selected circumstances when plasmacytic differentiation is present, *MYD88* mutational analysis should be considered to differentiate MZLs from WM/LPL.⁷⁻¹⁰

The NCCN Guidelines provide recommendations for the diagnosis, workup, and treatment for MALT lymphomas, NMZL, and SMZL.

Literature Search Criteria and Guidelines Update Methodology

Prior to the update of this version of the NCCN Guidelines® for B-Cell Lymphomas an electronic search of the PubMed database was performed to obtain key literature in MZLs published since the previous Guidelines update, using the following search terms: MALT lymphoma, gastric MALT lymphoma, nongastric MALT lymphoma, nodal marginal zone lymphoma and splenic marginal zone lymphoma. The PubMed database was chosen as it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.¹¹

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.

The data from key PubMed articles deemed as relevant to these Guidelines and discussed by the panel have been included in this version of the Discussion section (eg, e-publications ahead of print, meeting abstracts). Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.



The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

MALT Lymphomas

The gastrointestinal (GI) tract is the most common site of involvement with the stomach being the most common primary site in gastric MALT lymphomas; salivary/parotid glands, skin, ocular adnexa, head and neck, lung, thyroid, and breast are the other common sites of involvement in nongastric MALT lymphoma.² Bone marrow involvement has been reported in about 15% to 20% of MALT lymphomas.^{12,13}

Although MALT lymphomas are localized in most patients, about a third of patients present with disseminated disease and localized disease is more frequently observed with gastric subtype than with nongastric subtype.^{12,13} MALT lymphomas tend to be indolent, with similar long-term outcomes reported between gastric and nongastric subtypes and between patients with disseminated and localized disease.^{12,14} More recent retrospective data, however, reported higher progression-free survival (PFS) outcomes in patients with gastric MALT lymphomas and in patients with localized disease.¹⁵

t(11;18), t(1;14), t(14;18), and t(3;14) are the most common genetic alterations implicated in the pathogenesis of MALT lymphomas.¹⁶ t(11;18), resulting in the formation of the chimeric fusion gene, *API2-MALT1*, is frequently detected in gastric and pulmonary MALT lymphomas.^{17,18} t(1;14) results in the overexpression of BCL10 protein and occurs in 1% to 2% of MALT lymphomas, usually detected in MALT lymphomas of the stomach, lung, and skin.¹⁹ The presence of both t(11;18) and BCL10 overexpression is associated with locally advanced disease, which is less likely to respond to *H. pylori* eradication with antibiotic therapy.²⁰ t(14;18) results in the deregulated expression of *MALT1* gene and has been reported to occur in 15% to 20% of MALT lymphomas, most frequently

detected in MALT lymphomas of the liver, skin, ocular adnexa, and the salivary gland.^{18,21} t(3;14) results in the upregulation of the *FOXP1* gene and is associated with the MALT lymphomas of thyroid, ocular adnexa, and skin.²² The clinical significance of t(14;18) and t(3;14) is unknown.

Gastric MALT Lymphoma

Diagnosis

An endoscopic biopsy is required to establish the diagnosis of gastric MALT lymphoma, as fine-needle aspiration (FNA) is not adequate for diagnosis. Endoscopy may reveal erythema, erosions, or ulcerations.²³ Adequate hematopathology review of biopsy material and immunophenotyping are needed to establish a diagnosis. The recommended markers for an immunohistochemistry (IHC) panel include CD20, CD3, CD5, CD10, CD21 or CD23, kappa/lambda, CCND1, BCL2, and BCL6; the recommended markers for flow cytometry analysis include CD19, CD20, CD5, CD23, and CD10. The typical immunophenotype for MALT lymphoma is CD5-, CD10-, CD20+, CD23-/+ , CD43 -/+ , cyclin D1-, and BCL2- follicles.

H. pylori infection has a critical role in the pathogenesis of gastric MALT lymphomas and its eradication can lead to tumor remission.²³⁻²⁵ Therefore, staining for detection of *H. pylori* should be performed. However, *H. pylori* infection is not evident in approximately 5% to 10% of patients with gastric MALT lymphomas and the translocation t(11;18) was reported to occur at a high frequency in *H. pylori*-negative patients with gastric MALT lymphomas.²⁶ This chromosomal abnormality has been associated with disseminated disease and resistance to antibiotic treatment in patients with gastric MALT lymphoma.^{27,28} Molecular analysis by polymerase chain reaction (PCR) or fluorescence in situ hybridization (FISH) for the evaluation of t(11;18) is recommended. In some cases, molecular analysis for the detection of antigen receptor gene rearrangements and cytogenetic or FISH evaluation for t(3;14), t(1;14) and t(14;18) may also be useful.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Workup

A comprehensive physical examination should be performed with attention to nongastric sites such as the eyes and skin, and performance status should be assessed. Laboratory evaluations should include a complete blood count (CBC) with differentials and platelets, comprehensive metabolic panel, and measurement of serum lactate dehydrogenase (LDH) levels. Evaluation of bone marrow biopsy, with or without aspirates, may be useful under certain circumstances. Testing for HCV should be performed for all patients and testing for hepatitis B virus (HBV) is indicated for patients being considered for treatment with rituximab-containing regimens due to the risk of viral reactivation. Appropriate imaging studies include chest/abdomen/pelvis CT scan with contrast of diagnostic quality and/or whole body PET/CT (especially if involved-site radiation therapy [ISRT] is anticipated). A multigated acquisition (MUGA) scan/echocardiogram should be performed if the patient is being considered for treatment with regimens containing anthracycline or anthracenedione.

Special aspects of the workup for gastric MALT lymphoma include direct endoscopic assessment of the GI tract and additional evaluation of the tumor specimen for the presence of *H. pylori*. If the *H. pylori* infection status is negative based on histopathologic evaluation, other noninvasive testing methods may be employed to confirm negative status (ie, stool antigen test, urea breath test, blood antibody test). Nondiagnostic atypical lymphoid infiltrates that are *H. pylori* positive should be rebiopsied to confirm or exclude lymphoma prior to treatment of *H. pylori*. At some NCCN Member Institutions, endoscopic ultrasound (EUS) is used to complement conventional endoscopy at the time of the initial workup and at follow-up. EUS provides information regarding the depth of involvement in the gastric wall, which provides essential information for some of the currently used staging systems; it also helps to distinguish benign lymphoid aggregates from lymphoma associated with *H. pylori* infection.²⁹

In addition, EUS staging is also useful in predicting the efficacy of *H. pylori* eradication therapy.^{30,31} EUS with multiple biopsies of anatomic sites is particularly useful for *H. pylori*-positive patients because the likelihood of tumor response to antibiotic therapy is related to depth of tumor invasion.

Staging

It is unknown whether staging for MALT lymphomas should follow the standard Ann Arbor staging systems used for nodal lymphomas. The widely used Lugano staging system for GI lymphomas is a modification of the original Ann Arbor staging system.³² Ann Arbor stage III has been removed. The TNM staging system corresponds to the staging in gastric cancer and the depth of the lymphoma infiltration is measured by EUS. Stage I refers to disease confined to the GI tract (single primary or multiple non-contiguous lesions (stage I₁, the infiltration is limited to mucosa or submucosa and in stage I₂, infiltration is present in the muscularis propria, serosa, or both). Stage II refers to disease extending into the abdomen (stage II₁, localized nodal involvement [perigastric lymph nodes] and in stage II₂, distant nodal involvement). Stage IIE refers to penetration of serosa to involve adjacent organs or tissues; subscripts (1 or 2) may be added to the designation if both the lymph nodes and adjacent organs are involved. Stage IV refers to disseminated extranodal involvement or concomitant supradiaphragmatic nodal involvement, and cases with disseminated nodal involvement appear to behave more like NMZL or follicular lymphoma (FL).

Treatment

H. pylori infection plays a central role in the pathogenesis of localized gastric MALT lymphoma and the treatment approach depends on the *H. pylori* infection status. In retrospective and prospective clinical studies, *H. pylori* eradication with antibiotic therapy resulted in lymphoma regression in 70% to 95% of patients with localized disease.³³⁻⁴⁰ In a multicenter cohort follow-up study of 420 patients with gastric MALT lymphoma, the response rate was 77% for *H. pylori* eradication therapy and the estimated



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

10-year freedom from treatment failure (FFTF), overall survival (OS), and event-free survival (EFS) rates were 90%, 95%, and 86%, respectively.⁴⁰ The presence of t(11;18) and *H. pylori*-negativity and submucosal invasion were independent predictors of resistance to *H. pylori* eradication. *H. pylori* eradication therapy generally comprises a proton pump inhibitor (eg, omeprazole or other agents such as lansoprazole or rabeprazole) along with a combination of antibiotics including clarithromycin and amoxicillin (or metronidazole for patients allergic to penicillin).²³

Long-term follow-up data from clinical studies suggest that RT is an effective treatment modality in gastric MALT lymphoma after failure with *H. pylori* eradication therapy or as an initial treatment for patients with *H. pylori*-negative disease.⁴¹⁻⁴⁶

In the randomized controlled study of 241 patients with localized gastric MALT lymphomas, after a median follow-up of 8 years, the 10-year EFS rates were 52%, 52%, and 87%, respectively, for patients treated with surgery, RT, and chemotherapy ($P < .01$).⁴¹ The 10-year OS rate was not significantly different between the treatment groups (80% vs. 75% vs. 87%, respectively). In an analysis of registry data from a German multicenter study in patients with localized gastric lymphomas, extended-field RT (30 Gy followed by 10 Gy boost) alone resulted in an EFS and OS rate of 88% and 93%, respectively, in the subgroup of patients with gastric MALT lymphomas.⁴² These outcomes were not significantly different from patients with gastric MALT lymphomas who received combined modality therapy with surgery and RT (EFS and OS rates of 72% and 83%, respectively).⁴² This study had also included patients with gastric MALT lymphomas who experienced treatment failure with *H. pylori* eradication therapy. In a retrospective study of 192 patients with localized (stage I-II) MALT lymphomas, the complete response (CR) rate was 99% in the group of patients treated with involved-field RT (IFRT) (30–35 Gy) alone.⁴⁴ After a median follow-up of 7 years, the estimated

10-year relapse-free survival (RFS) rate, disease-free survival (DFS) rate, and OS rate were 76%, 68%, and 87%, respectively. Patients with thyroid and gastric MALT lymphomas had better outcome than patients with MALT lymphomas diagnosed at other sites ($P = .004$). The 10-year RFS rate was 95% for thyroid, 92% for stomach, 68% for salivary glands, and 67% for orbit. In a recent prospective randomized study, reduced-dose RT (25 Gy) was also effective for the treatment of *H. pylori*-negative stage I–II disease or *H. pylori*-positive disease that is refractory to *H. pylori* eradication therapy.⁴⁶ The presence of the t(11,18) translocation was a predictor for persistent lymphoma after *H. pylori* eradication therapy.

Rituximab has demonstrated activity in patients not eligible for eradication therapy (*H. pylori*-negative disease) or in those with gastric MALT lymphoma relapsed or refractory to *H. pylori* eradication therapy.^{47,48} In a prospective study of 27 patients with relapsed/refractory disease or *H. pylori*-negative disease (majority of patients [81%] had stage I or II₁ disease), rituximab resulted in an overall response rate (ORR) of 77% (46% CR). At a median follow-up of 28 months from start of treatment, all patients were alive and 54% of patients were disease free.⁴⁷ In a retrospective study that evaluated treatment options for patients with persistent disease despite *H. pylori* eradication therapy or *H. pylori*-negative disease ($n = 106$; 28 patients were treated with rituximab), rituximab resulted in an ORR of 73% (64% CR).⁴⁸ After a median follow-up of 5 years, the 5-year PFS and OS rates were 70% and 95%, respectively.

Stage I–II

H. Pylori-positive Stage I₁, or I₂ or Stage II₁ Disease

Antibiotic therapy to eradicate the *H. Pylori* infection is recommended, regardless of t(11;18) status. However, it should be noted that t(11;18) is a predictor for lack of response to antibiotic therapy.^{28,40} Patients with



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

involvement of submucosa or regional lymph nodes are also much less likely to respond to antibiotic therapy.⁴⁰

Evaluation with endoscopy and biopsy is recommended after 3 months following antibiotic therapy. In symptomatic patients after antibiotic therapy, restaging can be done earlier than 3 months. Clinical follow-up every 3 to 6 months for 5 years and then yearly or as clinically indicated is recommended for patients with disease responding to initial antibiotic therapy (*H. pylori* negative and lymphoma negative). There is increasing evidence that late relapses can occur after antibiotic treatment and a long duration of follow-up is appropriate.

For patients with t(11;18)-negative disease or t(11;18) status unknown, ISRT is recommended for those who are *H. pylori* negative with persistent lymphoma.^{41-44,46} Alternatively, asymptomatic patients can be observed for another 3 months. CRs may be observed as early as 3 months after antibiotic treatment but can take longer to achieve (up to 18 months). If re-evaluation suggests slowly responding disease or asymptomatic nonprogression, continued observation may be warranted (category 2B). Second-line antibiotic treatment is recommended for patients with persistent *H. pylori* and regressing or stable lymphoma. Lastly, patients who are *H. pylori* positive with progressive or symptomatic lymphoma should be treated with ISRT and second-line antibiotics.

Follow-up endoscopy is recommended after 3 months following completion of treatment (ISRT or second-line antibiotic treatment). Biopsy is recommended to rule out evidence of diffuse large B-cell lymphoma (DLBCL). Any area of DLBCL should be treated according to recommendations for DLBCL. Clinical follow-up every 3 to 6 months for 5 years and then yearly thereafter (or as clinically indicated) is recommended for patients with a CR. The optimal interval for follow-up endoscopy and imaging is not known. Follow-up endoscopy and imaging at NCCN Member Institutions are performed as clinically indicated based

on symptoms. Locoregional RT is indicated for patients with no response to second-line antibiotic therapy and chemoimmunotherapy is recommended for patients with no response to ISRT.

For patients with t(11;18)-positive disease, those with persistent lymphoma after antibiotic therapy should be treated with ISRT or rituximab (if ISRT is contraindicated).^{41-44,46,47} Evaluation with endoscopy and biopsy is recommended after 3 to 6 months following ISRT or rituximab. Clinical follow-up every 3 to 6 months for 5 years and then yearly or as clinically indicated is recommended for patients with responsive disease (*H. pylori* negative and lymphoma negative). Antibiotic treatment can be considered for patients with persistent *H. pylori* and regressing lymphoma. However, patients with persistent lymphoma (regardless of *H. pylori* status) following ISRT or rituximab should be managed with chemoimmunotherapy.

Suggested treatment regimens for first-line therapy are discussed in *Chemoimmunotherapy for Marginal Zone Lymphomas*.

H. pylori Negative Stage I₁, or I₂ or Stage II₁ Disease

ISRT is preferred for *H. pylori*-negative disease and rituximab is an option for patients with contraindications to ISRT.^{41-44,46,47} Evaluation with endoscopy and biopsy is recommended after 3 to 6 months following ISRT or rituximab. Additional treatment options are similar to that described above for patients with t(11;18)-positive disease.

Treatment for Relapsed Disease

Locoregional ISRT is recommended for local recurrence following antibiotic therapy. Treatment for recurrence following ISRT or systemic recurrence following antibiotic therapy is based on the presence of indications for treatment. Asymptomatic patients without indications for treatment are observed. Chemoimmunotherapy is recommended for patients with indications for treatment.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Suggested treatment regimens for second-line therapy are discussed in the section *Chemoimmunotherapy for Marginal Zone Lymphomas*.

Stage IIE, II₂, or IV

In patients with distant nodal, advanced stage disease (which is uncommon), treatment is similar to that described for patients with advanced stage FL. Asymptomatic patients without indications for treatment are observed. The decision to treat is guided by end-organ dysfunction or the presence of symptoms (such as GI bleeding, early satiety), bulky disease at presentation, steady progression of disease, or patient preference. Chemoimmunotherapy or palliative ISRT is recommended for patients with indications for treatment. Suggested treatment regimens for first-line therapy are discussed in *Chemoimmunotherapy for Marginal Zone Lymphomas*. Surgical resection is generally limited to specific clinical situations such as life-threatening hemorrhage. Although disease control is excellent with total gastrectomy, the long-term morbidity has precluded routine surgical resection.

Evaluation with endoscopy is recommended after completion of treatment. Recurrent disease should be managed with treatment regimens for second-line therapy as discussed in *Chemoimmunotherapy for Marginal Zone Lymphomas*.

Nongastric MALT Lymphomas

Diagnosis

Adequate hematopathology review of biopsy materials and immunophenotyping are needed to establish a diagnosis. The recommended markers for an IHC panel include CD20, CD3, CD5, CD10, CD21 or CD23, kappa/lambda, CCND1, and BCL2; the recommended markers for flow cytometry analysis include CD19, CD20, CD5, CD23, and CD10. The typical immunophenotype for MALT lymphoma is CD5-, CD10-, CD20+, CD23-/+ , CD43 -/+ , cyclin D1-, and BCL2-. Molecular analysis to detect antigen receptor gene rearrangement or t(11;18) may be

useful in certain cases. In addition, cytogenetics or FISH for t(11;18), t(3;14), t(11;14), and t(14;18) may also be considered under certain circumstances.

Workup

The workup for nongastric MALT lymphoma is similar to the workup for other non-Hodgkin's lymphomas (NHLs). A comprehensive physical examination should be performed and performance status should be assessed. Laboratory evaluations should include a CBC with differentials and platelets, comprehensive metabolic panel, and measurement of serum LDH levels. Evaluation of bone marrow biopsy, with or without aspirates, may be useful for patients with multifocal disease. In addition, endoscopy with multiple biopsies of anatomical sites may be useful in selected cases. Upper GI endoscopy should be considered where primary site is thought to be in head/neck or lungs.⁴⁹ Testing for HCV should be performed for all patients and testing for HBV is indicated for patients being considered for treatment with rituximab-containing regimens due to the risk of viral reactivation. Infectious pathogens such as *C. jejuni* and *C. psittaci* have been associated with nongastric MALT lymphomas, but testing for these pathogens is not required for disease workup or management.³

Appropriate imaging studies include chest/abdomen/pelvis CT scan with contrast of diagnostic quality and/or whole body PET/CT. A MUGA scan/echocardiogram should be performed if the patient is being considered for treatment with regimens containing anthracycline or anthracenedione. MRI with contrast should be done for neurologic evaluation or if CT with contrast is contraindicated.

Treatment

ISRT is an effective initial treatment for patients with nongastric MALT lymphoma.^{44,45,50} In a retrospective study of 167 patients with localized MALT lymphomas treated with RT with or without chemotherapy



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

(nongastric MALT lymphoma, $n = 142$), the group who received IFRT alone ($n = 144$; dose range 25–35 Gy; 25 Gy for orbit) had an estimated 10-year RFS rate and OS rate of 74% and 89%, respectively.⁴⁴ The 10-year RFS rates for patients with primary involvement of the thyroid, salivary gland, and orbital adnexa were 95%, 68%, and 67%, respectively. In another retrospective study of 208 patients with nongastric MALT lymphomas (Ann Arbor stage III–IV in 44%), the ORR among patients treated with chemotherapy, RT, or surgery were 65%, 76%, and 90%, respectively.⁵⁰ After a median follow-up of 3 years, the estimated 5-year EFS and OS rates were 37% and 83%, respectively. The 5-year OS rates were significantly higher among patients with Ann Arbor stage I–II disease compared with those with stage III–IV disease (94% vs. 69%; $P = .001$). On multivariate analysis, bone marrow involvement was the only significant independent predictor of inferior EFS and OS.⁵⁰

Rituximab has demonstrated activity in patients with nongastric MALT lymphoma.^{51–53} The IELSG evaluated the clinical activity of single-agent rituximab in a phase II study in patients with untreated as well as relapsed MALT lymphomas (35 patients; 20 patients with nongastric MALT lymphoma).⁵¹ Among patients with nongastric MALT lymphoma, treatment with rituximab resulted in an ORR of 80% (55% CR and 25% partial response [PR]). For the entire study population, the ORR was significantly higher in the chemotherapy-naïve patients than in previously treated patients (87% and 45% respectively; $P = .03$).

ISRT (24–30 Gy) is the preferred treatment for patients with stage I–II disease and is also recommended for patients presenting with stage IV disease.^{44,45,50} In patients with stage IV disease, definitive treatment of multiple sites and palliative treatment of symptomatic sites may be indicated. RT dose is site dependent, with lower doses usually reserved for orbital involvement. Based on anecdotal responses to antibiotics in ocular and cutaneous MZLs, some physicians may give an empiric course

of doxycycline prior to initiating therapy. Observation may be considered for patients whose diagnostic biopsy was excisional or in whom RT could result in significant morbidity.

Rituximab is an option for selected patients with stage I–II disease.^{51–53} Surgical excision may be appropriate treatment for certain extranodal sites (e., lung, thyroid, colon, small intestine, breast) in patients with stage I–II disease.¹³ Observation is recommended if there is no residual disease following surgery and locoregional RT should be considered for patients with positive surgical margins.

Clinical follow-up (including repeat diagnostic tests and imaging based on the site of disease and as clinically indicated) should be conducted every 3 to 6 months for 5 years and then annually thereafter (or as clinically indicated). Local recurrence may be treated with ISRT (if not previously received) or managed according to recommendations for advanced-stage NMZL. In patients with systemic recurrence, observation is recommended for patients without any indications for treatment. Chemoimmunotherapy is recommended for patients with indications for treatment. Suggested treatment regimens for first-line therapy (for treatment-naïve patients) or second-line therapy (in patients who have received prior treatment with rituximab) are discussed in *Chemoimmunotherapy for Marginal Zone Lymphomas*.

Nodal Marginal Zone Lymphoma

In patients with NMZL, peripheral lymphadenopathy is present in nearly all cases (>95%); thoracic or abdominal lymph nodes may also be involved in about 50% of cases and involvement of bone marrow and peripheral blood may be seen in about 30% to 40% and 10% of cases, respectively.^{54,55} Although about two-thirds of patients with newly diagnosed NMZL present with advanced-stage disease, most tumors are non-bulky and B symptoms are present in only about 15% of patients.^{54,55} NMZL has an indolent



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

disease course, but long-term survival outcomes appear less favorable compared with MALT lymphomas.^{54,56} In a retrospective analysis of data from patients with MZL, the OS rate (56% vs. 81%) and 5-year failure-free survival rate (28% vs. 65%) were lower in the subgroup of patients with NMZL compared with those with MALT lymphoma.⁵⁴ A SEER database analysis also reported more favorable 5-year survival outcomes for patients with MALT lymphoma (89%) compared to those with SMZL (80%) or NMZL (77%).⁵⁶ A more recent cohort analysis has reported more favorable survival outcomes for 56 patients with NMZL treated in the rituximab era (the majority of patients [79%] presented with advanced-stage disease).⁵⁷ After a median follow-up of 38 months, the PFS was 42 months and the estimated OS at 120 months after diagnosis was 72%.

Diagnosis

Adequate hematopathology review of biopsy materials and immunophenotyping are needed to establish a diagnosis. NMZL is rare and occurs most commonly as disseminated disease from extranodal MALT lymphoma. The recommended markers for an IHC panel include CD20, CD3, CD5, CD10, CD21 or CD23, kappa/lambda, CCND1, and BCL2; the recommended markers for flow cytometry include CD19, CD20, CD5, CD23, and CD10. The typical immunophenotype for NMZL is CD5-, CD10-, CD20+, CD23-/+ , CD43 -/+, cyclin D1-, and BCL2-. Pediatric NMZL should be considered with localized disease in young patients. Molecular analysis to detect antigen receptor gene rearrangement or t(11;18) (by PCR) may be useful in certain cases. In addition, cytogenetics or FISH for t(11;18), t(3;14), t(11;14), t(14;18), del(13q), and del(7q) may also be considered under certain circumstances.

Workup

NMZL occurs primarily in the lymph nodes, although involvements of additional extranodal sites are common.^{54,55} The diagnosis of NMZL requires careful evaluation to rule out extranodal sites of primary disease

and must be distinguished from nodal FL, mantle cell lymphoma (MCL), chronic lymphocytic leukemia (CLL), and WM/LPL. A comprehensive physical examination should be performed and performance status should be assessed. Laboratory evaluations should include a CBC with differentials and platelets, comprehensive metabolic panel, and measurement of serum LDH levels. Evaluation of bone marrow biopsy with aspirates should be performed to document clinical stage I–II disease. Bone marrow biopsy may be deferred until treatment is indicated. Testing for HCV should be performed for all patients, and testing for HBV is indicated for patients being considered for treatment with rituximab-containing regimens due to the risk of viral reactivation.

Appropriate imaging studies include chest/abdomen/pelvis CT scan with contrast of diagnostic quality and/or whole body PET/CT. A MUGA scan/echocardiogram should be performed if the patient is being considered for treatment with regimens containing anthracycline or anthracenedione.

First-line Treatment

Stage I–II

ISRT (24–30 Gy) is recommended for patients with non-bulky stage I–II disease. Clinical follow-up (complete history and physical exam and laboratory assessment is recommended every 3–6 months for the first 5 years, and then annually thereafter or as clinically indicated) is recommended for patients achieving CR or PR. Surveillance imaging with CT scans can be performed no more than every 6 months up to the first 2 years following completion of treatment, and then no more than annually thereafter. Patients with disease not responding to ISRT should be managed as described for stage III or IV disease.

Anti-CD20 monoclonal antibody (MAB) with or without chemotherapy with or without ISRT is recommended for patients with bulky disease (stage I or contiguous stage II) or non-contiguous stage II disease. The addition of



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

ISRT is included with a category 2B recommendation. Observation may be appropriate in circumstances where potential toxicity of ISRT outweighs potential clinical benefit. Clinical follow-up with a complete history and physical exam, laboratory assessment, and surveillance imaging (as described above) is recommended for patients with CR or PR after completion of treatment. Patients with non-responsive disease should be managed as described below for stage III or IV disease.

Stage III–IV

Observation is recommended for patients with no indications for treatment. Treatment should only be initiated when a patient presents with indications for treatment (based on the modified GELF criteria). The modified criteria used to determine treatment initiation include: symptoms attributable to FL (not limited to B symptoms); threatened end-organ function; cytopenia secondary to lymphoma; bulky disease (single mass >7 cm or 3 or more masses >3 cm); splenomegaly; and steady progression over at least 6 months.

Suggested regimens for first-line therapy are discussed in *Chemoimmunotherapy for Marginal Zone Lymphomas*.

First-line Consolidation or Extended Therapy

Observation is recommended for patients achieving CR or PR to first-line therapy.

Rituximab maintenance (single-dose rituximab every 3 months until treatment failure) or retreatment with rituximab (rituximab weekly x 4 at the time of each progression until treatment failure) was evaluated in an exploratory sub-study in patients with low-tumor-burden, non-follicular, indolent B-cell NHL responding to induction therapy with single-agent rituximab.⁵⁸ The ORR after induction therapy with rituximab was 40% (52% for patients with MZL). After a median of 4 years from randomization, maintenance rituximab significantly improved time to

treatment failure (TTTF) compared to rituximab retreatment (5 years and 1 year respectively; $P = .012$). Consolidation with rituximab (375 mg/m² one dose every 12 weeks) is recommended for patients initially treated with single-agent rituximab.

Second-line and Subsequent Therapy

Frequently, patients with disease relapse or progression of disease after first-line therapy will benefit from a second period of observation. Considerations and indications for treatment of relapsed/refractory or progressive disease include, among other factors, the modified GELF criteria, which include: symptoms attributable to MZL (not limited to B symptoms); threatened end-organ function; significant cytopenia secondary to lymphoma; bulky disease; splenomegaly; and steady progression over at least 6 months.

Progressive disease should be histologically documented to exclude transformation, especially in the presence of raising LDH levels, disproportional growth in one area, development of extranodal disease, or development of new constitutional symptoms. Areas of high standardized uptake value (SUV) should raise suspicion for the presence of transformation. However, a positive PET/CT scan does not replace a biopsy; rather, results of the PET/CT scan should be used to direct a biopsy to enhance the diagnostic yield from the biopsy.

Suggested regimens for second-line therapy are discussed under *Chemoimmunotherapy for Marginal Zone Lymphomas*.

Second-line Consolidation or Extended Dosing

Obinutuzumab maintenance (1 g every 8 weeks for a total of 12 doses) is included as an option for patients with rituximab-refractory disease treated with bendamustine + obinutuzumab based on the results of the GADOLIN study.⁵⁹



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

High-dose therapy followed by autologous stem cell rescue (HDT/ASCR) has been associated with survival benefit in patients with relapsed or refractory disease.⁶⁰⁻⁶² HDT/ASCR is included as an option for consolidative therapy for patients with disease responding to second-line therapy. Allogeneic hematopoietic cell transplant (HCT) may also be considered for highly selected patients.^{61,63}

Splenic Marginal Zone Lymphoma

SMZL is characterized by the presence of splenomegaly in all cases, which may become symptomatic when massive or when associated with cytopenias.^{55,64} Peripheral lymph nodes are generally not involved while splenic hilar lymph nodes are often involved.⁶⁴ Involvement of thoracic or abdominal lymph nodes may also be seen in about a third of patients with SMZL.⁵⁵ In addition, bone marrow involvement is present in the majority of patients (about 85%) and involvement of peripheral blood occurs in 30% to 50% of patients.^{55,64} The disease course of SMZL is generally indolent, although most patients present with advanced-stage disease. In a retrospective study that evaluated the clinical outcomes of 124 patients with non-MALT-type MZL, the median time to progression (TTP) and median OS were 7 years and 9 years, respectively, for the subgroup of 59 patients with SMZL.⁵⁵ In a cohort analysis of 64 patients with SMZL treated in the rituximab era, the estimated median PFS and OS were 53 months and 156 months, respectively, after a median follow-up of 38 months.⁶⁵

Diagnosis

Adequate hematopathology review of biopsy materials and immunophenotyping are needed to establish a diagnosis. The diagnosis of SMZL requires bone marrow involvement with or without peripheral blood involvement by small lymphoid cells with immunoglobulin (Ig) light chain restriction that lack characteristic features of other small B-cell neoplasms (ie, CD5, CD10, cyclin D1). The recommended markers for an IHC panel

include CD20, CD3, CD5, CD10, CD21 or CD23, CD43, kappa/lambda, IgD, CCND1, BCL2, and annexin A1; the recommended markers for flow cytometry analysis include CD19, CD20, CD5, CD23, CD10, CD43, and CD103. The typical immunophenotype for SMZL is CD5-, CD10-, CD20+, CD23-/+, CD43-, cyclin D1-, BCL2 follicles-, annexinA1-, CD103-, and with expression of both IgM and IgD. This lymphoma is distinguished from CLL by the absence of CD5 expression, strong CD20 expression, and variable CD23 expression, and from hairy cell leukemia (HCL) by the absence of CD103 expression.

SMZL is most definitively diagnosed at splenectomy, since the immunophenotype is nonspecific and morphologic features on the bone marrow may not be diagnostic. However, in a patient with splenomegaly (small or no M component) and a characteristic intra sinusoidal lymphocytic infiltration of the bone marrow, the diagnosis can strongly be suggested on bone marrow biopsy, if the immunophenotype is consistent. Plasmacytoid differentiation with cytoplasmic Ig detectable on paraffin sections may occur. In such cases, the differential diagnosis may include WM/LPL. *MYD88* mutation analysis⁷⁻¹⁰ and *BRAF* mutation analysis can be useful in selected cases for differentiating SMZL from WM/LPL and HCL, respectively.

Workup

The initial workup for SMZL is similar to the other indolent lymphomas. A comprehensive physical examination should be performed and performance status should be assessed. Laboratory evaluations should include a CBC with differentials and platelets, comprehensive metabolic panel, and measurement of serum LDH levels. Serum protein electrophoresis (SPEP) and/or measurement of quantitative Ig levels should be performed. If elevated Ig or monoclonal Ig is detected, further characterization by immunofixation of blood may be useful. Evaluation of bone marrow biopsy with or without aspirates should be performed. Testing for HCV should be performed for all patients and testing for HBV is



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

indicated for patients being considered for treatment with rituximab-containing regimens due to the risk of viral reactivation. Other useful evaluations may include cryoglobulin testing for detection of abnormal proteins frequently associated with hepatitis C, and direct Coombs test for evaluation of autoimmune hemolytic anemia.

Appropriate imaging studies include chest/abdomen/pelvis CT scan with contrast of diagnostic quality and/or whole body PET/CT. A MUGA scan/echocardiogram should be performed if the patient is being considered for treatment with regimens containing anthracycline or anthracenedione.

Treatment

Asymptomatic patients with no splenomegaly or progressive cytopenias can be observed until indications for treatment develop.^{66,67} Patients presenting with splenomegaly should be treated depending on their HCV serology status.

Hepatology evaluation is recommended for patients with HCV-positive disease and appropriate antiviral therapy should be initiated for patients without contradictions for treatment of hepatitis. Interferon (IFN)-based antiviral therapy has been shown to induce virologic and hematologic responses in patients with HCV-positive MZLs.⁶⁸⁻⁷³ In a retrospective series of 134 patients with HCV-positive indolent B-cell NHLs (36 patients with SMZL), among the patients who received antiviral therapy with IFN or pegylated-IFN, with or without ribavirin as first-line therapy (n = 100; 23 patients with SMZL), the ORR and CR rates were 77% (65% for patients with SMZL) and 47%, respectively, and a sustained virologic response was observed in 78% of patients. The median duration of response was 33 months. After a median follow-up of 4 years, the 5-year PFS and OS rates were 63% and 92%, respectively. IFN-free antiviral therapy with a combination of direct-acting antivirals (DAAs) has also been reported to be effective, resulting in high lymphoproliferative disease response rates in

patients with HCV-positive MZLs.⁷⁴ Patients with no response to antiviral therapy or contraindications for treatment of hepatitis should be managed as described below for patients with HCV-negative disease.

Patients with HCV-negative disease can be observed if they are asymptomatic. Rituximab monotherapy (with or without maintenance rituximab) is associated with high response rates with durable remissions and is the preferred treatment for patients who are symptomatic (cytopenias or symptoms of splenomegaly, weight loss, early satiety, or abdominal pain).⁷⁵⁻⁷⁹ In a retrospective study that assessed treatment with rituximab in 43 patients with SMZL, rituximab monotherapy resulted in an ORR of 100% with a CR in 79% of patients and was associated with less toxicity.⁷⁷ The 3-year DFS was more favorable with rituximab-containing therapy, with or without splenectomy (79%) compared with splenectomy alone (29%) or chemotherapy alone (25%).⁷⁷ In this small retrospective study, patients who underwent splenectomy had a higher likelihood of attaining a CR with rituximab (100%) than those who did not undergo splenectomy (67%).

Splenectomy is recommended if the disease is not responsive to rituximab. In retrospective studies, splenectomy with or without chemotherapy has demonstrated favorable outcomes with a median OS exceeding 10 years and a 10-year OS rate of 61% to 84%.^{4,67,80-84} Pneumococcal and meningococcal vaccination should be given at least 2 weeks before splenectomy.

Patients should be monitored on a regular basis following treatment. Clinical follow-up (including repeat diagnostic tests and imaging studies, as clinically indicated) should be performed every 3 to 6 months for 5 years and then annually or as clinically indicated thereafter. In patients with recurrence, observation is recommended for patients without any indications for treatment. Splenectomy, ISRT, or chemoimmunotherapy are recommended for patients with indications for treatment. Suggested



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

treatment regimens for first-line therapy (for treatment-naïve patients) or second-line therapy (in patients who have received prior treatment with rituximab) are discussed in *Chemoimmunotherapy for Marginal Zone Lymphomas*.

Chemoimmunotherapy for Marginal Zone Lymphomas

The NCCN B-Cell Lymphomas Panel stratified all the regimens into 3 categories (based on the evidence, efficacy, toxicity, preexisting comorbidities, and in some cases access to certain agents): preferred regimens, other recommended regimens, and useful under certain circumstances.

First-line Therapy: Preferred Regimens

Chemoimmunotherapy with anti-CD20 monoclonal antibody

Bendamustine + rituximab (BR), RCHOP (cyclophosphamide, doxorubicin, vincristine, prednisone), or RCVP (cyclophosphamide, vincristine, prednisone) are included as preferred regimens.

In the multicenter randomized phase III study (StiL NHL1) that included 67 patients with MZL, BR was superior to RCHOP in terms of PFS in all histologic subtypes of indolent lymphomas except MZL ($P = .32$).^{85,86} The phase III randomized study (BRIGHT) also showed that BR was noninferior to RCHOP or RCVP in terms of PFS in patients with previously untreated indolent lymphoma or MCL.^{87,88} This study included 46 patients with MZL (28 patients were randomized to BR and 18 patients were randomized to RCHOP or RCVP). The ORRs in the subgroup of patients with MZL were 92% (20% CR) and 71% (24% CR), for BR and RCHOP/RCVP, respectively.⁸⁷

BR and RCVP have also demonstrated efficacy as first-line therapy in small clinical studies that have evaluated these regimens specifically in patients with MZL.⁸⁹⁻⁹⁴ In a multicenter phase II trial (MALT 2008-01) of 60 patients with previously untreated MALT lymphoma (20 patients with

gastric MALT lymphoma; 34 patients with nongastric MALT lymphoma), after 3 cycles of treatment, BR resulted in an ORR of 100% (75% CR) and the CR rates were significantly higher for patients with gastric MALT lymphoma (90% vs. 62%; $P = .023$).⁸⁹ After completion of 6 cycles, the ORR and CR rates were 100% and 98%, respectively, with no significant differences according to primary site of disease. After a median observation of 82 months, the 7-year EFS rate was 88% for all patients (90% for patients with gastric MALT lymphoma and 84% for those with nongastric MALT lymphoma). In a phase II study (BRISMA) of 56 patients with SMZL treated with BR as first-line therapy, the ORR and CR rates were 91% and 73%, respectively. The 3-year PFS and OS rates were 90% and 96%, respectively.⁹² In a phase II study of 40 patients with advanced-stage MZL, RCVP resulted in an ORR of 88% (60% CR).⁹³ After a median follow-up of 38 months, the estimated 3-year PFS and OS rates were 59% and 95%, respectively. RCVP regimen is also effective for the treatment of patients ineligible for *H. pylori* eradication therapy or those with gastric MALT lymphoma that is resistant to *H. pylori* eradication therapy.⁹⁴

In the clinical trial that evaluated the efficacy and safety of obinutuzumab compared to rituximab when used in combination with chemotherapy (bendamustine, CHOP, or CVP) in patients with previously untreated advanced-stage MZL (195 patients), after a median follow-up of 38 months, there were no clinically relevant differences in ORR and PFS between the study arms.⁹⁵ However, grade 3 to 5 adverse events were higher with obinutuzumab than with rituximab. Obinutuzumab-based chemoimmunotherapy is not recommended as first-line therapy for patients with MZL.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Rituximab

As discussed earlier, rituximab monotherapy has demonstrated activity in MALT lymphomas^{47,48,51-53} and SMZL.⁷⁵⁻⁷⁹ It is the preferred initial treatment for patients with SMZL.

First-line Therapy: Other Recommended Regimens

Ibritumomab tiuxetan and lenalidomide + rituximab are included with a category 2B recommendation.

Ibritumomab Tiuxetan

In a phase II study of 16 patients with previously untreated MZL, ibritumomab tiuxetan resulted in an ORR of 88% (50% CR; 31% PR).⁹⁶ With a median follow-up of 66 months, the median PFS was 48 months and the median OS was not reached. The 5-year PFS and OS rates were 40% and 72%, respectively.

If ibritumomab tiuxetan is considered, bilateral cores are recommended and the pathologist should provide the percent of overall cellular elements involved in the marrow. As of 2010, updates suggest a trend towards an increased risk of MDS with the use of ibritumomab tiuxetan. Therefore, karyotype ± FISH for known MDS markers should be done.

Lenalidomide + Rituximab

In the phase II trial that evaluated lenalidomide + rituximab in patients with untreated, advanced-stage, indolent lymphoma (n = 110; 30 patients with MZL), the ORR was 89% (67% CR; 22% PR) and the median PFS was 54 months among the subgroup of patients with MZL.⁹⁷

Lenalidomide monotherapy or in combination with rituximab also has demonstrated activity in patients with MALT lymphoma, resulting in ORRs of 61% (33% CR; 28% PR) and 80% (54% CR; 26% PR), respectively.^{98,99}

Second-line and Subsequent Therapy

Alternate non-cross-resistant chemoimmunotherapy with anti-CD20 MAB (bendamustine with obinutuzumab or rituximab, RCHOP, or RCVP),^{59,100,101} lenalidomide ± rituximab,^{102,103} rituximab monotherapy,⁵¹ or ibritumomab tiuxetan¹⁰⁴ are included as options for relapsed/refractory disease. Ibrutinib and phosphatidylinositol 3-kinase (PI3K) inhibitors (idelalisib, copanlisib, and duvelisib) have shown promising activity in the treatment of relapsed or refractory MZL.¹⁰⁵⁻¹⁰⁸ Ibrutinib is also FDA approved for the treatment of relapsed/refractory MZL.

Bendamustine + Obinutuzumab

The GADOLIN study (413 patients: 47 patients had MZL; 28 patients were randomized to bendamustine + obinutuzumab; 19 patients were randomized to bendamustine monotherapy) confirmed that the combination of bendamustine + obinutuzumab followed by obinutuzumab maintenance significantly prolongs PFS compared to bendamustine monotherapy in patients with rituximab-refractory indolent NHL.⁵⁹ After a median follow-up of 32 months, the median PFS was significantly longer with bendamustine + obinutuzumab than with bendamustine monotherapy for the overall intent-to-treat study population (26 months vs. 14 months; $P < .001$).

Lenalidomide + Rituximab

In the MAGNIFY trial that is evaluating lenalidomide + rituximab followed by maintenance therapy in patients with relapsed/refractory indolent NHL (155 patients; 27 patients with MZL), induction therapy with lenalidomide + rituximab resulted in an ORR of 55% (45% CR) in the subgroup of patients with MZL.¹⁰³

In the multicenter, double-blind, randomized phase III study (AUGMENT) that compared lenalidomide + rituximab versus rituximab monotherapy in patients with relapsed/refractory indolent lymphoma (n = 358; 63 patients had MZL), although the ORRs were higher with lenalidomide + rituximab



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

(64% [29% CR; 35% PR]) compared to rituximab monotherapy (44% [13% CR; 31% PR]) in the subgroup of patients with MZL, PFS improvements favored lenalidomide + rituximab all the histologic subtypes of indolent lymphomas except MZL (HR:1.0; 95% CI, 0.47–2.13).¹⁰⁹ After a median follow-up of 28 months, the estimated 2-year OS rates were 82% and 94%, respectively for lenalidomide + rituximab and rituximab monotherapy for the subgroup of patients with MZL.

Ibrutinib

In a multicenter phase II study of 60 patients with relapsed or refractory MZL (after prior therapy with an anti-CD20 MAB-based regimen), ibrutinib resulted in an ORR of 48% (3% CR and 45% PR).¹⁰⁵ After a median follow-up of 19 months, the median PFS was 14 months. The median PFS by subtype was 14 months for extranodal MZL, 19 months for SMZL, and 8 months for NMZL. The median OS has not been reached and the 18-month estimated OS rate was 81%. Infections (19%), anemia (14%), pneumonia (8%), and fatigue (6%) were the most common grade ≥3 adverse events.

PI3K Inhibitors

In a post hoc analysis that evaluated the safety and efficacy of idelalisib in the subset of 21 patients with MZL (enrolled in 2 clinical studies that evaluated idelalisib in patients with relapsed/refractory indolent NHL), ibrutinib resulted in an ORR of 33% and 47%. The median PFS was 7 months.¹⁰⁶

In the subset analysis from the CHRONOS-1 trial (142 patients with relapsed/ refractory indolent NHL after ≥2 prior lines of therapy; 23 patients has MZL), copanlisib resulted in an ORR of 70% (13% CR).¹⁰⁷ After a median duration of treatment of 23 weeks, 85% of patients were estimated to be in response at 9 months.

In the phase II study (DYNAMO) evaluating the safety and efficacy of duvelisib in 129 patients with indolent NHL refractory to both rituximab and chemotherapy or radioimmunotherapy, duvelisib resulted in an ORR of 47% (39% for the subgroup of patients with MZL [n = 18]).¹⁰⁸ The estimated median duration of response and median PFS were 10 months and 9.5 months, respectively.

Based on these results, idelalisib, copanlisib, and duvelisib are included as options for relapsed/refractory disease after 2 prior therapies.

Ibritumomab Tiuxetan

Ibritumomab tiuxetan also has demonstrated clinical activity in patients with relapsed or refractory extranodal MZL.¹⁰⁴ In a small study of 30 patients, ibritumomab tiuxetan resulted in an ORR of 90% (77% CR; 13% PR). After a median follow-up of 5 years, 12 patients were still in CR. Ibritumomab tiuxetan is included as an option with a category 2B recommendation.

Suggested Treatment Regimens for Elderly or Infirm Patients

Rituximab monotherapy is the preferred treatment option for untreated as well as relapsed/refractory disease in elderly or infirm patients. Ibritumomab tiuxetan and alkylating agent-based chemotherapy (cyclophosphamide or chlorambucil) with or without rituximab are included as alternative options.¹¹⁰⁻¹¹²

In a study that evaluated chlorambucil + rituximab versus rituximab monotherapy in 49 patients with gastric MALT lymphoma, chlorambucil + rituximab was more efficient than rituximab alone in patients with t(11;18)-positive disease, whereas rituximab alone is as efficient as chlorambucil + rituximab in patients with t(11;18)-negative disease.¹¹⁰ In the international randomized IELSG-19 trial that evaluated the combination of chlorambucil + rituximab versus chlorambucil or rituximab monotherapy in patients with MALT lymphoma not previously treated with



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

systemic anticancer therapy, the addition of rituximab to chlorambucil resulted in significantly better EFS.¹¹¹ At a median follow-up of 7 years, the 5-year EFS rate was 68% for chlorambucil + rituximab compared to 51% and 50%, respectively, for chlorambucil and rituximab monotherapy. The 5-year OS rate was not significantly different between the treatment arms. Cyclophosphamide + rituximab was effective in the treatment of SMZL in frail patients not fit for splenectomy resulting in an ORR of 87% with a median PFS of 20 months.¹¹²

Histologically Transformed Marginal Zone Lymphoma

Histologic transformation of MZL to DLBCL occurs at an annual rate of approximately 3% to 5% across all subtypes and is generally associated with a poor clinical outcome.^{113,114} High-risk IPI scores, elevated LDH, and B symptoms; disease involvement in multiple mucosal sites and >4 nodal sites at the time of diagnosis of MZL; and failure to achieve CR after initial treatment have been reported as risk factors for histologic transformation to DLBCL.¹¹³⁻¹¹⁶

Treatment Options

Histologically transformed MZL is usually managed with chemoimmunotherapy regimens recommended for DLBCL. However, there are no data from randomized studies to support the optimal treatment, since clinical trials have often excluded this group of patients.

Anthracycline-based chemoimmunotherapy (with regimens recommended for first-line therapy for DLBCL, unless contraindicated) with or without ISRT is recommended for patients who have received minimal (ISRT alone or one course of single agent therapy including rituximab) or no chemotherapy prior to transformation. Observation or consolidation therapy with HDT/ASCR or allogeneic HCT (in selected patients), with or without ISRT (if not previously given), are included as treatment options for patients achieving CR or PR. Alternatively, patients achieving PR may be treated with ibritumomab tiuxetan or ISRT for localized residual

disease and/or residual FDG-avid disease not previously irradiated. Ibritumomab tiuxetan, second-therapy regimens recommended for DLBCL, or best supportive care are included as options for patients with non-responsive or progressive disease.

Enrollment in an appropriate clinical trial is the preferred option for patients who have received multiple prior therapies prior to transformation. In the absence of a suitable clinical trial, treatment options include ibritumomab tiuxetan, ISRT chemoimmunotherapy (with regimens for second-line therapy for DLBCL) with or without ISRT, or best supportive care. Observation or consolidation therapy with HDT/ASCR or allogeneic HCT (in selected patients), with or without ISRT (if not previously given), are included as treatment options for patients with disease responding to initial treatment. Patients with nonresponsive or progressive disease can be treated with any of the initial treatment options (not received previously), if they are candidates for additional therapy.

HDT/ASCR and allogeneic HCT may be reasonable treatment options for selected patients with histologically transformed MZL.¹¹⁷ The efficacy of HDT/ASCR or allogeneic HCT has not been confirmed in prospective controlled studies. Additional systemic therapy ± ISRT to induce CR prior to transplant should be considered for patients achieving PR. Repeat biopsy should be strongly considered if PET positive prior to initiating additional therapy. If transformation is coexisting with extensive MZL, rituximab maintenance should be considered for patients achieving CR.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

References

1. Swerdlow SH, Harris NL, Jaffe ES, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Revised 4th ed. Lyon, France: IARC; 2017.
2. Khalil MO, Morton LM, Devesa SS, et al. Incidence of marginal zone lymphoma in the United States, 2001-2009 with a focus on primary anatomic site. *Br J Haematol* 2014;165:67-77. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24417667>.
3. Zucca E, Bertoni F, Vannata B, Cavalli F. Emerging role of infectious etiologies in the pathogenesis of marginal zone B-cell lymphomas. *Clin Cancer Res* 2014;20:5207-5216. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25320370>.
4. Arcaini L, Paulli M, Boveri E, et al. Splenic and nodal marginal zone lymphomas are indolent disorders at high hepatitis C virus seroprevalence with distinct presenting features but similar morphologic and phenotypic profiles. *Cancer* 2004;100:107-115. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14692030>.
5. Arcaini L, Burcheri S, Rossi A, et al. Prevalence of HCV infection in nongastric marginal zone B-cell lymphoma of MALT. *Ann Oncol* 2007;18:346-350. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17071937>.
6. Berger F, Traverse-Glehen A, Felman P, et al. Clinicopathologic features of Waldenstrom's macroglobulinemia and marginal zone lymphoma: are they distinct or the same entity? *Clin Lymphoma* 2005;5:220-224. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15794852>.
7. Treon SP, Xu L, Yang G, et al. MYD88 L265P somatic mutation in Waldenstrom's macroglobulinemia. *N Engl J Med* 2012;367:826-833. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22931316>.
8. Gachard N, Parrens M, Soubeyran I, et al. IGHV gene features and MYD88 L265P mutation separate the three marginal zone lymphoma entities and Waldenstrom macroglobulinemia/lymphoplasmacytic lymphomas. *Leukemia* 2013;27:183-189. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22944768>.
9. Varettoni M, Arcaini L, Zibellini S, et al. Prevalence and clinical significance of the MYD88 (L265P) somatic mutation in Waldenstrom's macroglobulinemia and related lymphoid neoplasms. *Blood* 2013;121:2522-2528. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23355535>.
10. Xu L, Hunter ZR, Yang G, et al. MYD88 L265P in Waldenstrom macroglobulinemia, immunoglobulin M monoclonal gammopathy, and other B-cell lymphoproliferative disorders using conventional and quantitative allele-specific polymerase chain reaction. *Blood* 2013;121:2051-2058. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23321251>.
11. U.S. National Library of Medicine Key MEDLINE® Indicators Available at: http://www.nlm.nih.gov/bsd/bsd_key.html.
12. Thieblemont C, Berger F, Dumontet C, et al. Mucosa-associated lymphoid tissue lymphoma is a disseminated disease in one third of 158 patients analyzed. *Blood* 2000;95:802-806. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10648389>.
13. Zucca E, Conconi A, Pedrinis E, et al. Nongastric marginal zone B-cell lymphoma of mucosa-associated lymphoid tissue. *Blood* 2003;101:2489-2495. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12456507>.
14. Thieblemont C, Bastion Y, Berger F, et al. Mucosa-associated lymphoid tissue gastrointestinal and nongastrointestinal lymphoma behavior: analysis of 108 patients. *J Clin Oncol* 1997;15:1624-1630. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9193362>.
15. Ueda K, Terui Y, Yokoyama M, et al. Non-gastric advanced mucosa-associated lymphoid tissue (MALT) lymphoma has worse prognosis than gastric MALT lymphoma even when treated with rituximab-containing chemotherapy. *Leuk Lymphoma* 2013;54:1928-1933. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23216271>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

16. Farinha P, Gascoyne RD. Molecular pathogenesis of mucosa-associated lymphoid tissue lymphoma. *J Clin Oncol* 2005;23:6370-6378. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16155022>.

17. Auer IA, Gascoyne RD, Connors JM, et al. t(11;18)(q21;q21) is the most common translocation in MALT lymphomas. *Ann Oncol* 1997;8:979-985. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9402171>.

18. Murga Penas EM, Hinz K, Roser K, et al. Translocations t(11;18)(q21;q21) and t(14;18)(q32;q21) are the main chromosomal abnormalities involving MLT/MALT1 in MALT lymphomas. *Leukemia* 2003;17:2225-2229. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12931213>.

19. Willis TG, Jadayel DM, Du MQ, et al. Bcl10 is involved in t(1;14)(p22;q32) of MALT B cell lymphoma and mutated in multiple tumor types. *Cell* 1999;96:35-45. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9989495>.

20. Liu H, Ye H, Dogan A, et al. T(11;18)(q21;q21) is associated with advanced mucosa-associated lymphoid tissue lymphoma that expresses nuclear BCL10. *Blood* 2001;98:1182-1187. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11493468>.

21. Streubel B, Lamprecht A, Dierlamm J, et al. T(14;18)(q32;q21) involving IGH and MALT1 is a frequent chromosomal aberration in MALT lymphoma. *Blood* 2003;101:2335-2339. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12406890>.

22. Streubel B, Vinatzer U, Lamprecht A, et al. T(3;14)(p14.1;q32) involving IGH and FOXP1 is a novel recurrent chromosomal aberration in MALT lymphoma. *Leukemia* 2005;19:652-658. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15703784>.

23. Kahl B, Yang D. Marginal zone lymphomas: management of nodal, splenic, and MALT NHL. *Hematology Am Soc Hematol Educ Program*

2008:359-364. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19074110>.

24. Isaacson PG, Spencer J. Gastric lymphoma and *Helicobacter pylori*. *Important Adv Oncol* 1996:111-121. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8791131>.

25. Wotherspoon AC. Gastric lymphoma of mucosa-associated lymphoid tissue and *Helicobacter pylori*. *Annu Rev Med* 1998;49:289-299. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9509264>.

26. Ye H, Liu H, Raderer M, et al. High incidence of t(11;18)(q21;q21) in *Helicobacter pylori*-negative gastric MALT lymphoma. *Blood* 2003;101:2547-2550. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12517817>.

27. Morgner A, Bayerdorffer E, Neubauer A, Stolte M. *Helicobacter pylori* associated gastric B cell MALT lymphoma: predictive factors for regression. *Gut* 2001;48:290-292. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11171813>.

28. Liu H, Ye H, Ruskone-Fourmesttraux A, et al. T(11;18) is a marker for all stage gastric MALT lymphomas that will not respond to *H. pylori* eradication. *Gastroenterology* 2002;122:1286-1294. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11984515>.

29. Pavlick AC, Gerdes H, Portlock CS. Endoscopic ultrasound in the evaluation of gastric small lymphocytic mucosa-associated lymphoid tumors. *J Clin Oncol* 1997;15:1761-1766. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9164183>.

30. Sackmann M, Morgner A, Rudolph B, et al. Regression of gastric MALT lymphoma after eradication of *Helicobacter pylori* is predicted by endosonographic staging. MALT Lymphoma Study Group. *Gastroenterology* 1997;113:1087-1090. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9322502>.

31. Nakamura S, Matsumoto T, Suekane H, et al. Predictive value of endoscopic ultrasonography for regression of gastric low grade and high



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

grade MALT lymphomas after eradication of *Helicobacter pylori*. Gut 2001;48:454-460. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11247887>.

32. Rohatiner A, d'Amore F, Coiffier B, et al. Report on a workshop convened to discuss the pathological and staging classifications of gastrointestinal tract lymphoma. Ann Oncol 1994;5:397-400. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/8075046>.

33. Wotherspoon AC, Doglioni C, Diss TC, et al. Regression of primary low-grade B-cell gastric lymphoma of mucosa-associated lymphoid tissue type after eradication of *Helicobacter pylori*. Lancet 1993;342:575-577.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8102719>.

34. Steinbach G, Ford R, Globler G, et al. Antibiotic treatment of gastric lymphoma of mucosa-associated lymphoid tissue. An uncontrolled trial. Ann Intern Med 1999;131:88-95. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10419446>.

35. Montalbán C, Santon A, Boixeda D, et al. Treatment of low grade gastric mucosa-associated lymphoid tissue lymphoma in stage I with *Helicobacter pylori* eradication. Long-term results after sequential histologic and molecular follow-up. Haematologica 2001;86:609-617.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11418369>.

36. Chen L, Lin J, Tai J, et al. Long-term results of anti-*Helicobacter pylori* therapy in early-stage gastric high-grade transformed MALT lymphoma. J Natl Cancer Inst 2005;97:1345-1353. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16174856>.

37. Wundisch T, Thiede C, Morgner A, et al. Long-term follow-up of gastric MALT lymphoma after *Helicobacter pylori* eradication. J Clin Oncol 2005;23:8018-8024. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16204012>.

38. Stathis A, Chini C, Bertoni F, et al. Long-term outcome following *Helicobacter pylori* eradication in a retrospective study of 105 patients with localized gastric marginal zone B-cell lymphoma of MALT type. Ann Oncol

2009;20:1086-1093. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19193705>.

39. Andriani A, Miedico A, Tedeschi L, et al. Management and long-term follow-up of early stage H. pylori-associated gastric MALT-lymphoma in clinical practice: an Italian, multicentre study. Dig Liver Dis 2009;41:467-473. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18945654>.

40. Nakamura S, Sugiyama T, Matsumoto T, et al. Long-term clinical outcome of gastric MALT lymphoma after eradication of *Helicobacter pylori*: a multicentre cohort follow-up study of 420 patients in Japan. Gut 2012;61:507-513. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21890816>.

41. Aviles A, Nambo MJ, Neri N, et al. Mucosa-associated lymphoid tissue (MALT) lymphoma of the stomach: results of a controlled clinical trial. Med Oncol 2005;22:57-62. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15750197>.

42. Koch P, Probst A, Berdel WE, et al. Treatment results in localized primary gastric lymphoma: data of patients registered within the German multicenter study (GIT NHL 02/96). J Clin Oncol 2005;23:7050-7059.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16129843>.

43. Vrieling C, de Jong D, Boot H, et al. Long-term results of stomach-conserving therapy in gastric MALT lymphoma. Radiother Oncol 2008;87:405-411. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18343513>.

44. Goda JS, Gospodarowicz M, Pintilie M, et al. Long-term outcome in localized extranodal mucosa-associated lymphoid tissue lymphomas treated with radiotherapy. Cancer 2010;116:3815-3824. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20564130>.

45. Teckie S, Qi S, Chelius M, et al. Long-term outcome of 487 patients with early-stage extra-nodal marginal zone lymphoma. Ann Oncol 2017;28:1064-1069. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28327924>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

46. Schmelz R, Miehlke S, Thiede C, et al. Sequential H. pylori eradication and radiation therapy with reduced dose compared to standard dose for gastric MALT lymphoma stages IE & IIIE: a prospective randomized trial. J Gastroenterol 2018. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30327875>.

47. Martinelli G, Laszlo D, Ferreri AJM, et al. Clinical activity of rituximab in gastric marginal zone non-Hodgkin's lymphoma resistant to or not eligible for anti-Helicobacter pylori therapy. J Clin Oncol 2005;23:1979-1983. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15668468>.

48. Amiot A, Levy M, Copie-Bergman C, et al. Rituximab, alkylating agents or combination therapy for gastric mucosa-associated lymphoid tissue lymphoma: a monocentric non-randomised observational study. Aliment Pharmacol Ther 2014;39:619-628. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24467480>.

49. Mazloom A, Rodriguez A, Ha CS, et al. Incidence of gastric involvement in patients with nongastrointestinal extranodal marginal zone lymphoma. Cancer 2011;117:2461-2466. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24048794>.

50. Arcaini L, Burcheri S, Rossi A, et al. Nongastric marginal-zone B-cell MALT lymphoma: prognostic value of disease dissemination. Oncologist 2006;11:285-291. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16549813>.

51. Conconi A, Martinelli G, Thieblemont C, et al. Clinical activity of rituximab in extranodal marginal zone B-cell lymphoma of MALT type. Blood 2003;102:2741-2745. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12842999>.

52. Okamura I, Imai H, Mori K, et al. Rituximab monotherapy as a first-line treatment for pulmonary mucosa-associated lymphoid tissue lymphoma. Int J Hematol 2015;101:46-51. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25378228>.

53. Tuncer S, Tanyildiz B, Basaran M, et al. Systemic Rituximab Immunotherapy in the Management of Primary Ocular Adnexal

Lymphoma: Single Institution Experience. Curr Eye Res 2015;40:780-785. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25247376>.

54. Nathwani BN, Anderson JR, Armitage JO, et al. Marginal zone B-cell lymphoma: A clinical comparison of nodal and mucosa-associated lymphoid tissue types. Non-Hodgkin's Lymphoma Classification Project. J Clin Oncol 1999;17:2486-2492. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10561313>.

55. Berger F, Felman P, Thieblemont C, et al. Non-MALT marginal zone B-cell lymphomas: a description of clinical presentation and outcome in 124 patients. Blood 2000;95:1950-1956. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10706860>.

56. Olszewski AJ, Castillo JJ. Survival of patients with marginal zone lymphoma: analysis of the Surveillance, Epidemiology, and End Results database. Cancer 2013;119:629-638. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22893605>.

57. Starr AG, Caimi PF, Fu P, et al. Dual institution experience of nodal marginal zone lymphoma reveals excellent long-term outcomes in the rituximab era. Br J Haematol 2016;175:275-280. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27443247>.

58. Williams ME, Hong F, Gascoyne RD, et al. Rituximab extended schedule or retreatment trial for low tumour burden non-follicular indolent B-cell non-Hodgkin lymphomas: Eastern Cooperative Oncology Group Protocol E4402. Br J Haematol 2016;173:867-875. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26970533>.

59. Cheson BD, Chua N, Mayer J, et al. Overall Survival Benefit in Patients With Rituximab-Refractory Indolent Non-Hodgkin Lymphoma Who Received Obinutuzumab Plus Bendamustine Induction and Obinutuzumab Maintenance in the GADOLIN Study. J Clin Oncol 2018;36:2259-2266. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29584548>.

60. Li L, Bierman P, Vose J, et al. High-dose therapy/autologous hematopoietic stem cell transplantation in relapsed or refractory marginal



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

zone non-Hodgkin lymphoma. Clin Lymphoma Myeloma Leuk 2011;11:253-256. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21658651>.

61. Shimoni A. The role of stem-cell transplantation in the treatment of marginal zone lymphoma. Best Pract Res Clin Haematol 2017;30:166-171. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28288712>.

62. Avivi I, Arcaini L, Ferretti VV, et al. High-dose therapy and autologous stem cell transplantation in marginal zone lymphomas: a retrospective study by the EBMT Lymphoma Working Party and FIL-GITMO. Br J Haematol 2018;182:807-815. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29984825>.

63. Dholaria BR, Hammond WA, Roy V, et al. Allogeneic hematopoietic cell transplant for relapsed-refractory, marginal zone lymphoma: a single-center experience. Leuk Lymphoma 2018;59:2727-2730. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29566563>.

64. Thieblemont C. Clinical presentation and management of marginal zone lymphomas. Hematology Am Soc Hematol Educ Program 2005:307-313. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16304396>.

65. Starr AG, Caimi PF, Fu P, et al. Splenic marginal zone lymphoma: excellent outcomes in 64 patients treated in the rituximab era. Hematology 2017;22:405-411. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28105889>.

66. Thieblemont C, Felman P, Berger F, et al. Treatment of splenic marginal zone B-cell lymphoma: an analysis of 81 patients. Clin Lymphoma 2002;3:41-47. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12141954>.

67. Iannitto E, Ambrosetti A, Ammatuna E, et al. Splenic marginal zone lymphoma with or without villous lymphocytes. Hematologic findings and outcomes in a series of 57 patients. Cancer 2004;101:2050-2057. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15389479>.

68. Hermine O, Lefrere F, Bronowicki JP, et al. Regression of splenic lymphoma with villous lymphocytes after treatment of hepatitis C virus infection. N Engl J Med 2002;347:89-94. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12110736>.

69. Kelaidi C, Rollot F, Park S, et al. Response to antiviral treatment in hepatitis C virus-associated marginal zone lymphomas. Leukemia 2004;18:1711-1716. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15284859>.

70. Vallisa D, Bernuzzi P, Arcaini L, et al. Role of anti-hepatitis C virus (HCV) treatment in HCV-related, low-grade, B-cell, non-Hodgkin's lymphoma: a multicenter Italian experience. J Clin Oncol 2005;23:468-473. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15659492>.

71. Alric L, Besson C, Lapidus N, et al. Antiviral Treatment of HCV-Infected Patients with B-Cell Non-Hodgkin Lymphoma: ANRS HC-13 Lympho-C Study. PLoS One 2016;11:e0162965. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27749916>.

72. Arcaini L, Vallisa D, Rattotti S, et al. Antiviral treatment in patients with indolent B-cell lymphomas associated with HCV infection: a study of the Fondazione Italiana Linfomi. Ann Oncol 2014;25:1404-1410. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24799461>.

73. Michot JM, Canioni D, Driss H, et al. Antiviral therapy is associated with a better survival in patients with hepatitis C virus and B-cell non-Hodgkin lymphomas, ANRS HC-13 lympho-C study. Am J Hematol 2015;90:197-203. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25417909>.

74. Arcaini L, Besson C, Frigeni M, et al. Interferon-free antiviral treatment in B-cell lymphoproliferative disorders associated with hepatitis C virus infection. Blood 2016;128:2527-2532. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27605512>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

75. Bennett M, Sharma K, Yegena S, et al. Rituximab monotherapy for splenic marginal zone lymphoma. *Haematologica* 2005;90:856-858.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15951303>.

76. Tsimberidou AM, Catovsky D, Schlette E, et al. Outcomes in patients with splenic marginal zone lymphoma and marginal zone lymphoma treated with rituximab with or without chemotherapy or chemotherapy alone. *Cancer* 2006;107:125-135. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16700034>.

77. Else M, Marin-Niebla A, de la Cruz F, et al. Rituximab, used alone or in combination, is superior to other treatment modalities in splenic marginal zone lymphoma. *Br J Haematol* 2012;159:322-328. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23016878>.

78. Olszewski AJ, Ali S. Comparative outcomes of rituximab-based systemic therapy and splenectomy in splenic marginal zone lymphoma. *Ann Hematol* 2014;93:449-458. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24057925>.

79. Kalpadakis C, Pangalis GA, Sachanas S, et al. Rituximab monotherapy in splenic marginal zone lymphoma: prolonged responses and potential benefit from maintenance. *Blood* 2018;132:666-670.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29914978>.

80. Chacon JI, Mollejo M, Munoz E, et al. Splenic marginal zone lymphoma: clinical characteristics and prognostic factors in a series of 60 patients. *Blood* 2002;100:1648-1654. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12176884>.

81. Milosevic R, Todorovic M, Balint B, et al. Splenectomy with chemotherapy vs surgery alone as initial treatment for splenic marginal zone lymphoma. *World J Gastroenterol* 2009;15:4009-4015. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19705496>.

82. Lenglet J, Traulle C, Mounier N, et al. Long-term follow-up analysis of 100 patients with splenic marginal zone lymphoma treated with splenectomy as first-line treatment. *Leuk Lymphoma* 2014;55:1854-1860.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24206091>.

83. Xing KH, Kahlon A, Skinnider BF, et al. Outcomes in splenic marginal zone lymphoma: analysis of 107 patients treated in British Columbia. *Br J Haematol* 2015;169:520-527. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25854936>.

84. Pata G, Bartoli M, Damiani E, et al. Still a role for surgery as first-line therapy of splenic marginal zone lymphoma? Results of a prospective observational study. *Int J Surg* 2017;41:143-149. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28366762>.

85. Rummel MJ, Niederle N, Maschmeyer G, et al. Bendamustine plus rituximab versus CHOP plus rituximab as first-line treatment for patients with indolent and mantle-cell lymphomas: an open-label, multicentre, randomised, phase 3 non-inferiority trial. *Lancet* 2013;381:1203-1210.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23433739>.

86. Rummel MJ, Maschmeyer G, Ganser A, et al. Bendamustine plus rituximab (B-R) versus CHOP plus rituximab (CHOP-R) as first-line treatment in patients with indolent lymphomas: Nine-year updated results from the StiL NHL1 study [abstract]. *J Clin Oncol* 2017;35 (15_suppl):Abstract 7501. Available at:

http://ascopubs.org/doi/abs/10.1200/JCO.2017.35.15_suppl.7501.

87. Flinn IW, van der Jagt R, Kahl BS, et al. Randomized trial of bendamustine-rituximab or R-CHOP/R-CVP in first-line treatment of indolent NHL or MCL: the BRIGHT study. *Blood* 2014;123:2944-2952.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24591201>.

88. Flinn IW, van der Jagt R, Kahl B, et al. First-Line Treatment of Patients With Indolent Non-Hodgkin Lymphoma or Mantle-Cell Lymphoma With Bendamustine Plus Rituximab Versus R-CHOP or R-CVP: Results of the BRIGHT 5-Year Follow-Up Study. *J Clin Oncol* 2019;37:984-991.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30811293>.

89. Salar A, Domingo-Domenech E, Panizo C, et al. Long-term results of a phase 2 study of rituximab and bendamustine for mucosa-associated lymphoid tissue lymphoma. *Blood* 2017;130:1772-1774. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28801448>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

90. Laribi K, Tempescul A, Ghnaya H, et al. The bendamustine plus rituximab regimen is active against primary nodal marginal zone B-cell lymphoma. *Hematol Oncol* 2017;35:536-541. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27443419>.

91. Castelli R, Bergamaschini L, Deliliers GL. First-line treatment with bendamustine and rituximab, in patients with intermediate-/high-risk splenic marginal zone lymphomas. *Med Oncol* 2017;35:15. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29288421>.

92. Iannitto E, Bellei M, Amorim S, et al. Efficacy of bendamustine and rituximab in splenic marginal zone lymphoma: results from the phase II BRISMA/IELSG36 study. *Br J Haematol* 2018;183:755-765. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30407629>.

93. Kang HJ, Kim WS, Kim SJ, et al. Phase II trial of rituximab plus CVP combination chemotherapy for advanced stage marginal zone lymphoma as a first-line therapy: Consortium for Improving Survival of Lymphoma (CISL) study. *Ann Hematol* 2012;91:543-551. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21922208>.

94. Aguiar-Bujanda D, Llorca-Martinez I, Rivero-Vera JC, et al. Treatment of gastric marginal zone B-cell lymphoma of the mucosa-associated lymphoid tissue with rituximab, cyclophosphamide, vincristine and prednisone. *Hematol Oncol* 2014;32:139-144. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24123108>.

95. Herold M, Hoster E, Janssens A, et al. Immunochemotherapy with obinutuzumab or rituximab in a subset of patients in the randomised gallium trial with previously untreated marginal zone lymphoma (MZL). *Hematology Oncology* 2017;35:146-147. Available at: <https://onlinelibrary.wiley.com/doi/abs/10.1002/hon.2437> 136.

96. Lossos IS, Fabregas JC, Koru-Sengul T, et al. Phase II study of (90)Y Ibritumomab tiuxetan (Zevalin) in patients with previously untreated marginal zone lymphoma. *Leuk Lymphoma* 2015;56:1750-1755. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25315074>.

97. Fowler NH, Davis RE, Rawal S, et al. Safety and activity of lenalidomide and rituximab in untreated indolent lymphoma: an open-label, phase 2 trial. *Lancet Oncol* 2014;15:1311-1318. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25439689>.

98. Kiesewetter B, Troch M, Dolak W, et al. A phase II study of lenalidomide in patients with extranodal marginal zone B-cell lymphoma of the mucosa associated lymphoid tissue (MALT lymphoma). *Haematologica* 2013;98:353-356. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22899582>.

99. Kiesewetter B, Willenbacher E, Willenbacher W, et al. A phase 2 study of rituximab plus lenalidomide for mucosa-associated lymphoid tissue lymphoma. *Blood* 2017;129:383-385. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27879257>.

100. Kiesewetter B, Mayerhoefer ME, Lukas J, et al. Rituximab plus bendamustine is active in pretreated patients with extragastric marginal zone B cell lymphoma of the mucosa-associated lymphoid tissue (MALT lymphoma). *Ann Hematol* 2014;93:249-253. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23925930>.

101. Raderer M, Wohrer S, Streubel B, et al. Activity of rituximab plus cyclophosphamide, doxorubicin/mitoxantrone, vincristine and prednisone in patients with relapsed MALT lymphoma. *Oncology* 2006;70:411-417. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17220639>.

102. Sacchi S, Marcheselli R, Bari A, et al. Safety and efficacy of lenalidomide in combination with rituximab in recurrent indolent non-follicular lymphoma: final results of a phase II study conducted by the Fondazione Italiana Linfomi. *Haematologica* 2016;101:e196-199. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26858355>.

103. Coleman M, Andorsky DJ, Yacoub A, et al. Phase IIIB study of lenalidomide plus rituximab followed by maintenance in relapsed or refractory NHL: analysis of marginal zone lymphoma [abstract]. *Hematology Oncology* 2017;35:Abstract 139. Available at: <https://onlinelibrary.wiley.com/doi/abs/10.1002/hon.2437> 138.



104. Vanazzi A, Grana C, Crosta C, et al. Efficacy of (9)(0)Yttrium-ibritumomab tiuxetan in relapsed/refractory extranodal marginal-zone lymphoma. *Hematol Oncol* 2014;32:10-15. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23696416>.

105. Noy A, de Vos S, Thieblemont C, et al. Targeting Bruton tyrosine kinase with ibrutinib in relapsed/refractory marginal zone lymphoma. *Blood* 2017;129:2224-2232. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28167659>.

106. Martin P, Armas A, Gopal AK, et al. Idelalisib Monotherapy and Durable Responses in Patients with Relapsed or Refractory Marginal Zone Lymphoma (MZL) [abstract]. *Blood* 2015;126:Abstract 1543. Available at: <http://www.bloodjournal.org/content/126/23/1543.abstract>.

107. Dreyling M, Panayiotidis P, Egyed M, et al. Efficacy of Copanlisib Monotherapy in Patients with Relapsed or Refractory Marginal Zone Lymphoma: Subset Analysis from the CHRONOS-1 Trial [abstract]. *Blood* 2017;130:Abstract 4053. Available at: http://www.bloodjournal.org/content/130/Suppl_1/4053.abstract.

108. Flinn IW, Miller CB, Ardeschna KM, et al. DYNAMO: A Phase II Study of Duvelisib (IPI-145) in Patients With Refractory Indolent Non-Hodgkin Lymphoma. *J Clin Oncol* 2019:[Epub ahead of print]. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30742566>.

109. Leonard JP, Trneny M, Izutsu K, et al. AUGMENT: A Phase III Study of Lenalidomide Plus Rituximab Versus Placebo Plus Rituximab in Relapsed or Refractory Indolent Lymphoma. *J Clin Oncol* 2019;37:1188-1199. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30897038>.

110. Levy M, Copie-Bergman C, Amiot A, et al. Rituximab and chlorambucil versus rituximab alone in gastric mucosa-associated lymphoid tissue lymphoma according to t(11;18) status: a monocentric non-randomized observational study. *Leuk Lymphoma* 2013;54:940-944. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22978684>.

111. Zucca E, Conconi A, Martinelli G, et al. Final Results of the IELSG-19 Randomized Trial of Mucosa-Associated Lymphoid Tissue Lymphoma: Improved Event-Free and Progression-Free Survival With Rituximab Plus Chlorambucil Versus Either Chlorambucil or Rituximab Monotherapy. *J Clin Oncol* 2017;35:1905-1912. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28355112>.

112. Cervetti G, Ghio F, Cecconi N, et al. How to treat splenic marginal zone lymphoma (SMZL) in patients unfit for surgery or more aggressive therapies: experience in 30 cases. *J Chemother* 2017;29:126-129. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27454143>.

113. Meyer AH, Stroux A, Lerch K, et al. Transformation and additional malignancies are leading risk factors for an adverse course of disease in marginal zone lymphoma. *Ann Oncol* 2014;25:210-215. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24356632>.

114. Conconi A, Franceschetti S, Aprile von Hohenstaufen K, et al. Histologic transformation in marginal zone lymphomas. *Ann Oncol* 2015;26:2329-2335. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26400898>.

115. Alderuccio JP, Zhao W, Desai A, et al. Risk Factors for Transformation to Higher-Grade Lymphoma and Its Impact on Survival in a Large Cohort of Patients With Marginal Zone Lymphoma From a Single Institution. *J Clin Oncol* 2018;JCO1800138. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30312133>.

116. Alderuccio JP, Zhao W, Desai A, et al. Short survival and frequent transformation in extranodal marginal zone lymphoma with multiple mucosal sites presentation. *Am J Hematol* 2019. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30784098>.

117. Villa D, George A, Seymour JF, et al. Favorable outcomes from allogeneic and autologous stem cell transplantation for patients with transformed nonfollicular indolent lymphoma. *Biol Blood Marrow Transplant* 2014;20:1813-1818. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25042735>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Mantle Cell Lymphoma

Mantle cell lymphoma (MCL) comprises about 3% of all newly diagnosed cases of non-Hodgkin's lymphomas (NHL).¹ MCL is thought to possess the unfavorable characteristics of both indolent and aggressive NHL subtypes owing to the incurability of disease with conventional chemotherapy and a typically more aggressive disease course compared to indolent lymphomas.² MCL is characterized by the reciprocal chromosomal translocation t(11;14), juxtaposing the cyclin D1 locus with the immunoglobulin heavy chain (*IGH*) gene locus, resulting in the overexpression of cyclin D1 and the diagnosis of MCL generally requires the expression of cyclin D1.³

Cyclin D1-negative MCL with otherwise typical immunophenotype has also been reported, though rare (<5% of cases).^{4,5} In the rare instances of MCL negative for cyclin D1 and t(11;14), overexpression of cyclin D2 or cyclin D3 may be observed.^{6,7} Immunohistochemistry (IHC) for cyclin D2 or cyclin D3 is not helpful in establishing the diagnosis of cyclin D1-negative MCL, as these proteins are also expressed in other B-cell malignancies. A recent study of cyclin D1-negative MCL showed rearrangements involving the *CCND2* gene in 55% of cases, which was associated with high expression of cyclin D2 mRNA.⁸ Gene expression profiling (GEP) and miRNA profiling showed that the genomic signatures of cyclin D1-negative MCL were similar to those of cyclin D1-positive MCL.^{5,6,8} The pathologic features and clinical characteristics of cyclin D1-negative MCL appear to be similar to those of cyclin D1-positive MCL.^{6,8} Thus, in the absence of data suggesting otherwise, cyclin D1-negative MCL should not be managed differently than cyclin D1-positive MCL.

Literature Search Criteria and Guidelines Update Methodology

Prior to the update of this version of the NCCN Guidelines® for B-Cell Lymphomas an electronic search of the PubMed database was performed to obtain key literature in MCL published since the previous Guidelines

update. The PubMed database was chosen as it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.⁹

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.

The data from key PubMed articles deemed as relevant to these Guidelines and discussed by the panel have been included in this version of the Discussion section (eg, e-publications ahead of print, meeting abstracts). Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.

The complete details of the development and update of the NCCN Guidelines are available at www.NCCN.org.

Diagnosis

The diagnosis can be established by histologic examination in combination with IHC with a profile consisting of CD5+, CD10-/+ , CD20+, CD23-/+ , CD43+, and cyclin D1+. Some cases of MCL may be CD5- or CD23+. Currently available reagents for IHC evaluation of cyclin D1 are robust and yield good staining; however, in some cases, molecular analysis of *CCND1* rearrangements or cytogenetics or fluorescence in situ hybridization (FISH) for t(11;14) can be helpful for diagnosis.¹⁰ In certain cases, cytogenetics or FISH for t(14;18) and a FISH panel for chronic lymphocytic leukemia (CLL) may also be useful. In addition, Ki-67 should be included in the IHC panel for initial diagnostic workup. Ki-67 proliferation index of less than 30% has been associated with a more favorable prognosis.¹¹⁻¹⁵ However, this should not be used to guide treatment decisions at this time. Nuclear overexpression of the



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

transcription factor SOX11 is observed in nearly all cases of MCL, regardless of cyclin D1 expression level. Immunoglobulin heavy chain variable (*IGHV*) gene sequencing or IHC for SOX11 potentially aid in differentiating cyclin D1-negative MCL cases from other B-cell lymphomas.¹⁶⁻¹⁸

TP53 mutation is associated with poor prognosis in patients treated with conventional therapy, including hematopoietic cell transplantation (HCT). In an analysis that evaluated the prognostic effect of the most common genomic alterations in 183 patients with MCL treated in clinical trials, *TP53* mutations were significantly associated with Ki-67 >30%, blastoid morphology, and inferior responses to both induction chemotherapy and high-dose chemotherapy.¹⁹ *TP53* sequencing would be useful to identify patients with typical MCL with an expected aggressive clinical course, particularly if upfront HCT is anticipated.

GEP studies have identified an indolent, leukemic, and non-nodal MCL subtype that is largely *IGHV* mutated and mostly SOX11-negative, with peripheral blood, bone marrow and splenic involvement, low tumor burden and, a Ki-67 proliferation fraction <10%.²⁰⁻²² This description represents the most common indolent presentation; however, there are some patients with gastrointestinal (GI) or blood/bone marrow involvement only, which may express SOX11 and have an indolent course. The 2017 WHO classification also recognizes the two MCL subtypes with different clinicopathologic and molecular features: the usual MCL subtype (nodal or extranodal) that is SOX11+ and *IGHV* unmutated with an aggressive clinical course; and indolent (leukemic and non-nodal) subtype.²³

Mantle Cell Lymphoma “In Situ”

The presence of MCL-like B-cells in the mantle zones of morphologically reactive lymph nodes (“MCL in situ”) has been described in several case reports (including in patients with lymphoid hyperplasia).^{24,25} “MCL in situ”

is characterized by preservation of the lymph node architecture and presence of cyclin D1-positive B-cells restricted to the mantle zones with minimal expansion of the mantle zone (and with only minimal or no spread of cyclin D1-positive cells in the interfollicular area).²⁴⁻²⁶ An unusual presentation with a scattering of cyclin D1-positive cells in the germinal centers (but not the mantle zones) of a lymph node specimen retrospectively evaluated several years prior to the diagnosis of symptomatic MCL has also been reported.²⁷ “MCL in situ” is now known as “in situ mantle cell neoplasia” (ISMCN).

The occurrence of ISMCN in studies of reactive lymph nodes is very rare.^{26,28} In an analysis of a consecutive series of unselected surgical samples of reactive lymph nodes from patients without a history of lymphoma (n = 131; 1292 samples), no cases of ISMCN were identified.²⁸ Development of overt MCL in patients found to have ISMCN has been reported, although this appears to be very uncommon.²⁶ The significance or potential for malignancy of ISMCN in patients without known MCL remains uncertain. These cases appear to have a very indolent course with long-term survival even without treatment intervention.²⁶ It is therefore important to distinguish cases of ISMCN from cases of overt MCL with a mantle zone pattern. In patients with the former in whom overt MCL can be excluded based on a thorough evaluation (eg, biopsy of additional suspicious nodes, physical examination, peripheral blood flow cytometry, CT scan of neck, chest, abdomen, and pelvis) close follow-up may still be warranted.²⁹ Similar to “follicular lymphoma in situ”, the WHO classification recommends that a diagnosis of MCL not be made in such cases.

Workup

The initial workup should include a thorough physical examination with attention to node-bearing areas, and evaluation of performance status and constitutional symptoms. Laboratory assessments should include standard blood work including CBC with differential and a comprehensive metabolic



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

panel, in addition to measurements of serum lactate dehydrogenase (LDH). Patients with high tumor burden and elevated LDH should be assessed for spontaneous tumor lysis syndrome, including measurements of uric acid level. Measurement of serum beta-2-microglobulin levels may also be useful in some circumstances. HBV testing is recommended due to increased risks of viral reactivation when immunotherapy regimens are being considered for treatment. MCL is a systemic disease with frequent involvement of the bone marrow. For this reason, both the peripheral blood and bone marrow must be carefully evaluated for the presence of malignant cells. Adequate trephine biopsy should be obtained for initial staging evaluation, with or without bone marrow aspiration.

Chest/abdomen/pelvic CT with contrast of diagnostic quality and/or whole-body PET/CT scan are recommended as part of initial diagnostic workup. CT scan of the neck with contrast may be helpful in selected cases. In patients with the blastic variant or for patients presenting with CNS symptoms, a lumbar puncture should be performed to evaluate the cerebral spinal fluid for potential disease involvement. Multigated acquisition (MUGA) scan or echocardiogram is recommended for patients being considered for treatment regimens containing anthracyclines or anthracenediones.

GI involvement has been reported in 15% to 30% of patients with MCL. In two prospective studies, the frequency of GI tract involvement in patients with MCL was higher than that reported in the literature.^{30,31} In the study by Romaguera et al, MCL was histologically present in the lower and upper GI tract in 88% and 43% of patients, respectively, and 26% of patients presented with GI symptoms at the time of diagnosis.³⁰ Despite the high frequency of GI tract involvement (which was primarily observed at the microscopic level), the use of endoscopy with biopsies led to changes in clinical management in only 4% of patients.³⁰ Salar et al reported upper or lower GI tract involvement in 38% and 54% of patients, respectively, at diagnosis.³⁰ The NCCN Guidelines panel does not recommend endoscopy

or colonoscopy as part of routine initial workup, but suggests that it may be useful in certain circumstances. However, endoscopic or colonoscopic evaluation of the GI tract is necessary for confirmation of stage I–II disease and for response assessment to initial therapy.

Stage I–II

Few patients present with localized MCL and the available published literature on the management of localized disease is retrospective and anecdotal. In a retrospective analysis of 26 patients with limited bulk, early-stage (stage IA or IIA) MCL, radiation therapy (RT) with or without chemotherapy was associated with significantly improved progression-free survival (PFS) at 5 years (68% vs. 11%; $P = .002$) and a trend towards improved overall survival (OS).³²

In the absence of a suitable clinical trial, involved-site RT (ISRT) (30–36 Gy) or chemoimmunotherapy with or without ISRT are included as options for induction therapy. In highly selected patients with asymptomatic disease, close observation with deferred therapy is a reasonable option, especially for those with good performance status and lower risk scores on standard International Prognostic Index (IPI).^{33,34} These recommendations are based on treatment principles in the absence of more definitive clinical data.

For patients with a complete response (CR), clinical follow-up should be conducted every 3 to 6 months for the first 5 years, and then on a yearly basis or as clinically indicated. Treatment options for patients with a partial response (PR) or disease progression after induction therapy are similar to that of disease relapse as described below. Disease relapse after an initial CR to chemoimmunotherapy with or without ISRT should be managed with second-line therapy regimens recommended for stage II (bulky) or stage III–IV disease (see sections below). Disease relapse after an initial CR to ISRT alone should be managed with first-line induction therapy with



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

chemoimmunotherapy regimens recommended for stage II (bulky) and stage III-IV disease.

Stage II (bulky) and Stage III-IV

While some investigators have suggested that treatment with less aggressive regimens could yield survival outcomes similar to that achieved with more aggressive regimens, others have reported superior survival outcomes with more aggressive regimens.^{35,36} In a single-center cohort analysis of 111 patients with newly diagnosed MCL, among the 75 patients with available data on treatment regimens, the majority of patients (70%) had received CHOP-like regimen with or without rituximab and only 7% of patients had received rituximab in combination with hyperCVAD (R-hyperCVAD; rituximab, cyclophosphamide, vincristine, doxorubicin, and dexamethasone; alternating with high-dose methotrexate and cytarabine) and/or high-dose therapy with autologous stem cell rescue (HDT/ASCR).³⁵ The median OS from diagnosis was 85 months, and the 5-year OS rate was 66%. An analysis from the NCCN Oncology Outcomes Database (n = 167) reported superior PFS outcomes with R-hyperCVAD alone or with rituximab-containing regimens (eg, RCHOP) followed by HDT/ASCR, compared with RCHOP alone, in the first-line setting for patients <65 years of age.³⁶ However, the median PFS remained 3 to 4 years despite the use of aggressive regimens.

The standard treatment regimen for induction therapy is not yet established. There are no prospective randomized studies comparing the various aggressive regimens, although some randomized data exist for less aggressive regimens. The NCCN B-Cell Lymphomas Panel stratified the regimens for initial induction therapy into 3 categories (based on the available evidence, efficacy, toxicity, preexisting comorbidities, and in some cases access to certain agents): preferred regimens, other recommended regimens, and useful under certain circumstances.

Aggressive Disease

The majority of patients have advanced stage aggressive MCL, requiring systemic therapy. Induction therapy with aggressive regimens is recommended for patients who are candidates for HDT/ASCR, whereas induction therapy with less aggressive regimens is recommended for those who are not candidates for HDT/ASCR. *TP53* mutation has been associated with poor prognosis in patients treated with conventional therapy, including HCT.¹⁹ Participation in prospective clinical trials is strongly suggested for these patients.

Indolent Disease

Among patients with indolent MCL (*IGHV* mutated and mostly SOX11-negative with leukemic and non-nodal presentation [as described earlier]),²⁰⁻²² observation is a reasonable option for patients with asymptomatic disease with no indications for treatment. For patients with symptomatic disease or other indications for treatment, induction therapy with aggressive regimens is recommended for those without *TP53* mutation. Optimal management is unknown for those with *TP53* mutation. These patients could be treated with induction therapy followed by HDT/ASCR or less aggressive regimens.

Aggressive Induction Therapy: Preferred Regimens

Rituximab, Dexamethasone, Cytarabine and Platinum (carboplatin, cisplatin, or oxaliplatin)

In a phase III trial randomized trial that evaluated the role of rituximab maintenance after HDT/ASCR in patients <66 years (discussed below), induction chemoimmunotherapy with rituximab, dexamethasone, and cytarabine (RDHA) with a platinum agent (carboplatin, cisplatin, or oxaliplatin) resulted in an overall response rate (ORR) of 89% (77% CR).³⁷ HDT/ASCR following induction therapy was performed in 257 patients, resulting in a CR rate of 65%. In a subsequent analysis that evaluated the prognostic impact of carboplatin, cisplatin, or oxaliplatin on survival outcomes, the PFS and OS were identical for all three regimens



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

(RDHA-carboplatin, RDHA-cisplatin, and RDHA-oxaliplatin) but there was a trend towards improved PFS and OS with RDHA-oxaliplatin.³⁸ In the intent-to-treat population, the 4-year PFS and OS rates were 87% and 92% for RDHA-oxaliplatin compared to 65% and 76%, respectively, for the combined RDHA-carboplatin and RDHA-cisplatin group.

Alternating RCHOP and RDHAP (rituximab, dexamethasone, cytarabine, and cisplatin)

In the phase III randomized Intergroup trial conducted by the European-MCL Network (497 patients ≤65 years of age with advanced stage MCL were randomized to receive 6 cycles of RCHOP followed by HDT/ASCR using myeloablative radiochemotherapy or 6 cycles of alternating RCHOP and RDHAP followed by HDT/ASCR using high-dose cytarabine-containing myeloablative regimen), alternating RCHOP and RDHAP induction was associated with higher remission rates compared with RCHOP.³⁹ After a median follow-up of 6 years, the ORR (94% vs. 90%; $P = .14$;) and CR rates (38% vs. 25%; $P = .0016$) after induction therapy were higher for alternating RCHOP/R-DHAP than with RCHOP. The median time to treatment failure (TTF) was significantly longer in the RCHOP/RDHAP arm compared with the RCHOP arm (not reached vs. 9 years; $P = .038$) and the 5-year OS rates were 76% and 69%, respectively ($P = .12$). After HDT/ASCR, the ORR (98% vs. 97%) and CR rates (63% vs. 61%), were similar between treatment arms, although RCHOP/RDHAP was associated with longer remission duration (84 months vs. 49 months; $P = .0001$). These results suggest that induction therapy with alternating RCHOP/RDHAP is associated with improved outcomes, in patients <65 years.

In a phase II multicenter trial of the French cooperative group GELA ($n = 60$), induction therapy with 3 cycles each of RCHOP and RDHAP resulted in an ORR of 95% with CR in 57% of patients (age ≤65 years) with previously untreated MCL.⁴⁰ Patients went on to receive HDT/ASCR on this study. After a median follow-up of 67 months, the median event-free

survival (EFS) was 83 months and median OS has not been reached; the 5-year OS was 75%.⁴⁰

Dose-intensified RCHOP (maxi-CHOP) Alternating with High-Dose Cytarabine

In the Nordic MCL2 study ($n = 160$), induction therapy with dose-intensified RCHOP (maxi-CHOP) alternating with high-dose cytarabine resulted in an ORR and CR rate of 96% and 54%, respectively, in patients ≤65 years of age with previously untreated MCL.⁴¹ Patients with responding disease were eligible to proceed to HDT/ASCR. The ORR and 6-year PFS and OS rates were 96% (54% CR), 66%, and 70%, respectively, with no relapses occurring after a median follow-up of approximately 4 years (at the time of the initial report). After a median follow-up of 11 years, the median PFS and OS were 9 years and 13 years, respectively, for all the intent-to-treat patient population.⁴² The median OS was not reached and median PFS was 11 years for the 145 patients who proceeded to HDT/ASCR. The OS and PFS were significantly better for patients who had achieved CR to induction therapy than those who achieved PR ($P = .0038$ for OS; $P < .0001$ for PFS). However, a continuous pattern of relapse and an excess mortality were observed even after prolonged remission (late relapses were reported in 6 patients, who experienced disease relapse more than 10 years after the end of therapy).⁴² In the multivariate analysis, the MCL IPI (MIPI) and Ki-67 expression level were the only independent predictors of survival outcomes.^{42,43} However, in this trial, patients were monitored by disease-specific primers for molecular relapse, and those with disease relapse received rituximab as re-induction but were not considered to have disease relapse unless there was morphologic evidence of disease relapse.

HyperCVAD + Rituximab

R-hyperCVAD has resulted in favorable PFS and OS outcomes in patients with previously untreated MCL.⁴⁴⁻⁴⁷



In a phase II study of 97 patients with previously untreated with MCL, R-hyperCVAD produced 3-year failure-free survival (FFS) and OS rates of 64% and 82%, respectively, with a median follow-up time of 40 months.⁴⁴ After 10 years of follow-up, the median OS had not been reached and the median TTF was 5 years for all patients. Among patients ≤65 years, the median OS had not been reached and the median TTF was 6 years. In the multivariate analysis pre-treatment serum levels of beta-2- microglobulin, IPI score, and MIPI score were predictive of both OS and TTF.⁴⁵ FFS and OS rates were 43% and 60%, respectively; among patients 65 years or younger, the corresponding survival rates were 52% and 68%, respectively.

In the Italian study (60 evaluable patients), R-hyper-CVAD resulted in an ORR of 83% with a CR rate of 72%. The 5-year PFS and OS rates were 61% and 73%, respectively.⁴⁶ However, this regimen was associated with substantial toxicity.

In the SWOG 0213 study of 49 patients with previously untreated MCL (age <70 years), R-hyper-CVAD induced CR/CRu in 58% of patients.⁴⁷ With a median follow-up of 5 years, the median PFS and OS were 5 years (6 years for those ≤65 years) and 7 years, respectively. The 2-year PFS and OS rates were 63% and 76%, respectively.

Rituximab + ibrutinib has been used as a pre-treatment to limit the number of cycles of R-HyperCVAD and rituximab maintenance in young patients.⁴⁸

Aggressive Induction Therapy: Other Recommended Regimen

Bendamustine + rituximab is included as an option with a category 2B recommendation based on the results from 2 phase III randomized studies (discussed below).^{49,50}

Consolidation After Aggressive Induction Therapy

HDT/ASCR as first-line consolidation has demonstrated promising outcomes in a number of studies.⁵¹⁻⁵⁸

In a study conducted by the MD Anderson Cancer Center, HDT/ASCR in patients with MCL (n = 33) in first remission following treatment with hyper-CVAD resulted in 5-year disease-free survival (DFS) and OS rates of 42% and 77%, respectively.⁵² In particular, the subgroup of patients with low serum beta-2 microglobulin levels appeared to benefit most, with a 5-year OS rate of 100% (compared with 22% for patients with elevated beta-2 microglobulin).⁵² In a randomized trial conducted by the European-MCL Network, patients ≤65 years of age with advanced stage MCL (n = 122) in remission after CHOP-like chemotherapy were randomized to HDT/ASCR or maintenance with interferon alfa.⁵³ In this study, HDT/ASCR was associated with a significantly longer median PFS compared with interferon alfa maintenance (39 months vs. 17 months; *P* = .011). The 3-year OS rates were 83% and 77%, respectively, and were not significantly different between consolidation arms.⁵³ In an analysis of long-term outcomes from patients with MCL treated at the MD Anderson Cancer Center, the median PFS and OS were 42 months and 93 months, respectively, for the subgroup of patients treated primarily with hyper-CVAD (with or without rituximab) followed by HDT/ASCR in first remission (n = 50).⁵⁷

Rituximab maintenance after HDT/ASCR has been associated with survival benefit.^{37,59,60}

In a phase III randomized trial, rituximab maintenance after HDT/ASCR prolonged EFS, PFS, and OS compared to observation in patients <66 years.³⁷ In this trial involving 299 patients, 279 patients received induction chemoimmunotherapy with RDHA with a platinum agent (carboplatin, cisplatin, or oxaliplatin) resulting in an ORR of 89% (77% CR). HDT/ASCR



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

was performed in 257 patients following induction therapy. Among these 257 patients, 240 patients were randomized to receive rituximab maintenance or observation after HDT/ASCR. After a median follow-up of 50 months, the 4-year EFS, PFS, and OS rates were 79%, 83%, and 89%, respectively, for patients assigned to rituximab maintenance. The corresponding survival rates were 61%, 64%, and 80%, respectively, for patients assigned to observation. The results of a more recent study also confirmed the survival benefit of rituximab maintenance in patients with newly diagnosed MCL treated with Nordic MCL regimen followed by HDT/ASCR.⁶⁰ After a median follow-up of 4 years, the estimated 5-year PFS and OS rates were both 83% for patients assigned to rituximab maintenance. The corresponding survival rates were 63% and 79%, respectively, for patients assigned to observation.

The panel recommends consolidation with HDT/ASCR for eligible patients in CR following induction therapy with aggressive regimens, although no studies have compared maintenance rituximab with HDT/ASCR for patients in first CR. Rituximab maintenance following HDT/ASCR is included with a category 1 recommendation.^{37,59,60} For patients with only a PR to first-line therapy, additional treatment with second-line therapy regimens may be considered in an effort to improve the quality of a response. If the patient achieves a CR (or improved PR) with additional therapy, consolidation with HDT/ASCR may be considered for eligible patients, as discussed above. For patients who relapse after achieving a remission to first-line therapy, or for patients who experience disease progression during initial therapy, participation in clinical trials is preferred. In the absence of suitable clinical trials, second-line treatment options can be considered.

Less Aggressive Induction Therapy: Preferred Regimens

Bendamustine + Rituximab

The efficacy of bendamustine and rituximab (BR) as first-line therapy for MCL was established in 2 randomized phase III studies.^{49,50}

The randomized phase III study of the StiL (Study Group Indolent Lymphomas) compared BR versus RCHOP as first-line therapy in patients with advanced follicular, indolent, and mantle cell lymphomas (514 evaluable patients; MCL histology comprised 18% of patients).⁴⁹ With a median follow-up time of 45 months, the BR regimen was associated with significantly longer median PFS (primary endpoint) compared with RCHOP (70 months vs. 31 months; $P < .0001$). However, OS outcomes were not significantly different between treatment arms and ORR was similar in both arms (93% with BR vs. 91% with RCHOP), although the CR rate was significantly higher in the BR arm (40% vs. 30%; $P = .021$). Among the subgroup of patients with MCL, the median PFS was significantly higher with BR compared with RCHOP (35 months vs. 22 months; $P = .0044$).⁴⁹ The BR regimen was associated with less frequent serious adverse events (19% vs. 29%) and less grade 3–4 hematologic toxicities compared with RCHOP. Grade 3–4 neutropenia (29% vs. 69%), peripheral neuropathy (all grades; 7% vs. 29%), and infectious complications (all grades; 37% vs. 50%) were less frequent with BR compared with RCHOP was less frequent in the BR arm. Fatal sepsis occurred in 1 patient in the BR arm and 5 patients in the RCHOP arm. Skin toxicities (all grades) including erythema (16% vs. 9%) and allergic reactions (15% vs. 6%) were more frequent with BR than with RCHOP.⁴⁹ Although this phase III randomized trial showed superior PFS outcomes with the BR regimen compared with RCHOP, there may be limitations given that data from more than half of the patients in this trial were censored prior to the minimum follow-up period.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Another randomized phase III study (BRIGHT) demonstrated that BR was noninferior to RCHOP or RCVP (in terms of PFS) as first-line treatment of patients with indolent or MCL (224 patients were randomized to receive BR and 223 patients were randomized to receive RCHOP or RCVP).⁵⁰ At a median follow up of 5 years, the corresponding 5-year PFS rate for the overall study population was 66% and 56% ($P = .0025$), for BR and RCHOP/RCVP, respectively.⁵⁰ The 5-year OS rate was not statistically different between the treatment groups and the incidences of vomiting and drug hypersensitivity reactions, opportunistic infections and secondary malignancies were significantly higher in patients treated with BR.

Bortezomib, Rituximab, Cyclophosphamide, Doxorubicin, and Prednisone (VR-CAP)

A phase III randomized study evaluated the safety and efficacy of bortezomib in combination with rituximab, cyclophosphamide, doxorubicin, and prednisone (VR-CAP) versus RCHOP in patients with newly diagnosed MCL who are not candidates for HDT/ASCR.⁶¹ In this study, 487 patients were randomly assigned to VR-CAP or RCHOP; 268 patients (140 patients in the VR-CAP group and 128 patients in the RCHOP group) were included in the final follow-up analysis. The majority of patients had stage IV disease (74%) and 54% of patients had an IPI ≥ 3 . After median follow-up of 82, the median OS was significantly longer for patients in the VR-CAP group than in the RCHOP group (91 months vs. 56 months; $P = .001$). The incidences of grade ≥ 3 adverse events, although slightly higher with VR-CAP (93% compared to 84% with RCHOP), were manageable.

RCHOP

In the earlier studies, the addition of rituximab to CHOP chemotherapy was associated with high response rates but did not translate to prolonged PFS or OS.^{62,63} In a phase III randomized trial by the European-MCL Network that evaluated rituximab maintenance following induction therapy with R-FC (rituximab, fludarabine, and cyclophosphamide) or RCHOP in

patients >60 years with previously untreated MCL not eligible for HDT/ASCR ($n = 560$; 485 patients evaluable for response), RCHOP was associated with significantly longer median OS (67 months vs. 40 months) and significantly higher 4-year OS rate (62% and 47%, respectively; $P = .005$) than R-FC, although response rates (ORR; 86% vs. 78%; CR; 34% vs. 40%) and median duration of response (37 months vs. 36 months) were similar for both regimens.⁶⁴ Grade 3–4 hematologic toxicities occurred more frequently with R-FC induction.

Lenalidomide + Rituximab

In a multicenter phase II study that evaluated lenalidomide plus rituximab as induction and maintenance therapy for patients with previously untreated MCL ($n=38$), at the median follow-up of 30 months, lenalidomide plus rituximab resulted in an ORR of 87% (61% CR).⁶⁵ At a median follow-up of 64 months, the 3-year PFS and OS rates were 80% and 90%, respectively, with the 5-year estimated PFS and OS of 64% and 77%, respectively.⁶⁶

Modified HyperCVAD + Rituximab

In a small phase II pilot study of 22 patients with previously untreated MCL, a less intensive, modified R-hyper-CVAD regimen (without methotrexate or cytarabine, and with modifications to dose schedule of vincristine and steroids) followed by rituximab maintenance for 5 years resulted in a median PFS of 37 months with median OS not reached; the use of rituximab maintenance appeared to prolong PFS with acceptable toxicity.⁶⁷

In a phase II study (E1405), the addition of bortezomib to the modified R-hyper-CVAD (VcR-CVAD regimen) resulted in an ORR of 95% (68% CR) in patients with previously untreated MCL ($n = 75$).⁶⁸ Following induction therapy, patients proceeded with maintenance rituximab ($n = 44$) or consolidation with HCT off protocol ($n = 22$). After a median follow-up of 4.5 years, the 3-year PFS and OS rates were 72% and 88%, respectively.



No differences in PFS or OS were observed between patients who went on to receive rituximab maintenance or SCT.

Modified R-hyperCVAD is included as an option for patients >65 years.

Less Aggressive Induction Therapy: Other Recommended Regimen

RBAC (rituximab, bendamustine, and cytarabine)

In a phase II study that included 20 patients ≥65 years with previously untreated MCL not eligible for HDT/ASCR, the RBAC regimen was well tolerated resulting in an ORR of 100% (95% CR).⁶⁹ After a median follow-up of 26 months, the 2-year PFS rate was 95%.

Maintenance Rituximab After Less Aggressive Induction Therapy

In the aforementioned phase III randomized trial that evaluated induction therapy R-FC versus RCHOP, patients with disease responding to induction therapy (n = 316) underwent second randomization to receive maintenance therapy (given until progression) with either rituximab or interferon.⁶⁴ The median remission duration was significantly improved with rituximab maintenance compared with interferon alfa (75 months vs. 27 months; $P < .001$). After a median follow-up of 42 months, OS outcomes were not significantly different between the two maintenance arms (4-year OS: 79% with rituximab vs. 67% with interferon alfa).⁶⁴ However, in the subgroup of 184 patients treated with RCHOP induction therapy, the median OS (from end of induction) was significantly longer with rituximab maintenance compared with interferon alfa (not reached vs. 64 months; 4-year OS: 87% vs. 63%; $P = .005$). Moreover, grade 3–4 hematologic toxicities occurred more frequently with interferon alfa. Rituximab was associated with more frequent grade 1–2 infections.⁶⁴

Long-term follow-up data also confirmed that rituximab maintenance after RCHOP induction is associated with substantially prolonged PFS and OS in older patients who are not candidates for HDT/ASCR.⁷⁰ At a median follow-up of 7 years, after response to RCHOP, the 5-year PFS

rates were 51% and 22%, respectively ($P < .0001$) for patients randomized to rituximab maintenance and interferon. The corresponding 5-year OS rates were 79% and 59%, respectively ($P = .0026$).

This study suggests that RCHOP followed by rituximab maintenance as part of induction therapy may offer the best chance to prolong remission duration for patients who are not candidates for HDT/ASCR.^{64,70} Given the positive outcomes reported in this study (with median duration of response exceeding 6 years with rituximab maintenance and a 4-year OS rate of 87% in patients treated with RCHOP and rituximab maintenance), it is unknown whether first-line consolidation with HDT/ASCR provides an advantage over rituximab maintenance in patients of any age. At the present time, no data are available from randomized studies that would allow direct comparison of outcomes with these two different consolidation approaches.

Rituximab maintenance (every 8 weeks until disease progression) is recommended for patients who are not candidates for HDT/ASCR, and who are in remission after induction therapy with less aggressive regimens (category 1 following RCHOP).^{64,70} Rituximab maintenance following induction therapy with modified R-hyper-CVAD regimen has also been reported to provide extended disease control for patients who are not physically fit or not eligible to undergo aggressive first-line treatment regimens and HDT/ASCR.^{67,68,71} Data from a prospective, randomized phase II trial suggest that there is no benefit to rituximab maintenance after induction therapy with the BR regimen.⁷² Rituximab maintenance following induction therapy VR-CAP and RBAC has not been evaluated in clinical trials.

Second-line Therapy

The optimal approach to relapsed or refractory disease remains to be defined. Duration of response to prior chemoimmunotherapy is an



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

important factor in the selection of second-line therapy. Treatment with regimens containing small-molecule inhibitors or lenalidomide (with or without rituximab) should be considered for patients with early relapse (short response duration to prior chemoimmunotherapy), whereas alternate non-cross-resistant chemoimmunotherapy regimens (not previously given) would be more appropriate for patients with late relapse (extended response duration to prior chemoimmunotherapy).

See *Special Considerations for the Use of Small-Molecule Inhibitors* in the algorithm for monitoring and management of adverse reactions associated with small-molecule inhibitors (ibrutinib, acalabrutinib and venetoclax).

Short Response Duration to Prior Chemoimmunotherapy (< expected median PFS)

Preferred Regimens

Bruton Tyrosine Kinase Inhibitors

Acalabrutinib

In a phase II study of 124 patients with relapsed or refractory MCL, at a median follow-up of 15 months, acalabrutinib resulted in an ORR of 81% (40% CR).⁷³ The 12-month PFS and OS rates were 67% and 87%, respectively. Long-term follow-up (>24 months) also confirmed these initial findings.⁷⁴ The median PFS was 20 months and the estimated 24-month OS rate was 72%. Headache (38%), diarrhea (36%), fatigue (28%), cough (22%), bleeding (22%), and myalgia (21%) were the most common grade 1 or 2 adverse events. Anemia (10%), neutropenia (10%), pneumonia (6%), and infections (15%) were the most common grade 3 or 4 adverse events.

Ibrutinib ± Rituximab

Ibrutinib, either as monotherapy or in combination with rituximab results in durable responses with a favorable toxicity profile in patients with relapsed or refractory MCL.⁷⁵⁻⁷⁷ In the multicenter phase II study of 111 patients with

relapsed or refractory MCL after a median of 3 prior therapies, after a median follow-up of 27 months, ibrutinib monotherapy resulted in an ORR of 67% (23% CR) with a median duration of response of 18 months.⁷⁵ The 24-month PFS and OS rates were 31% and 47%, respectively. A phase III randomized trial (RAY) compared ibrutinib (n = 139) and temsirolimus (n = 141) in patients with relapsed or refractory MCL.⁷⁶ The ORR was 77% for ibrutinib and 47% for temsirolimus ($P < .0001$). After a median follow-up of 39 months, the median PFS was significantly longer with ibrutinib (16 months vs. 6 months) and there was also a trend toward improved OS in patients randomized to receive ibrutinib (30 months vs. 24 months; $P = .06$). Diarrhea (33%), fatigue (24%), and cough (23%) were the most common adverse events of any grade in the ibrutinib group. Thrombocytopenia (56%), anemia (44%), and diarrhea (31%) were the most common adverse events in the temsirolimus group. The rate of grade ≥ 3 bleeding events (9% vs. 5%) and atrial fibrillation (5% vs. 1%) were higher with ibrutinib.

In another phase II study of 50 patients with relapsed or refractory MCL after a median of 3 prior therapies, at a median follow-up of 47 months, ibrutinib in combination with rituximab resulted in a CR of 58% and the median PFS was 43 months.⁷⁷ Blastoid morphology, high-risk MIPI score, and high Ki-67 were associated with inferior survival.

The use of ibrutinib has been known to result in an initial transient lymphocytosis, which resolves by a median of 8 weeks after initiation of ibrutinib. Ibrutinib treatment has also been associated with grade ≥ 3 bleeding events and atrial fibrillation. The benefit and risk of ibrutinib should be considered in patients requiring anti-platelet or anticoagulant therapies.

Zanubrutinib

In November 2019, the FDA granted accelerated approval of zanubrutinib (selective and irreversible BTK inhibitor) for the treatment of



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

relapsed or refractory MCL after at least one prior therapy. This approval is based on the efficacy results (ORR as the primary end point) from two multicenter single-arm clinical trials (phase II study, 86 patients with relapsed or refractory MCL and phase I study, 48 patients [37 patients with relapsed or refractory MCL]).^{78,79} Zanubrutinib (160 mg BID given until disease progression or unacceptable toxicity) resulted in an ORR of 85% (77% CR) and 87% (30% CR) in these two trials, respectively. With a median follow-up of 14 to 16 months, treatment discontinuation due to progressive disease was reported in 22% and 33% of patients, respectively. The median PFS was 17 months and 15 months, respectively. Neutropenia (16%), anemia (6%), lung infection (6%), thrombocytopenia (5%) and hypertension (3.5%) were the most common grade 3 adverse events; major hemorrhage (serious or grade 3 bleeding or CNS bleeding of any grade) was reported in 2% of patients.⁷⁸

Lenalidomide With or Without Rituximab

Lenalidomide monotherapy has demonstrated efficacy in patients with relapsed or refractory MCL in multiple multicenter phase II studies.⁸⁰⁻⁸²

In the multicenter MCL-001 (EMERGE) study that evaluated the safety and efficacy of lenalidomide in relapsed or refractory MCL in patients who had received prior bortezomib ($n = 134$), lenalidomide resulted in an ORR of 28% (7.5% CR/CRu) by independent central review.⁸⁰ The median duration of response was 17 months. The median PFS and OS were 4 months and 19 months, respectively. The most common grade 3 or 4 toxicities were neutropenia (43%), thrombocytopenia (28%), anemia (11%), pneumonia (8%), and fatigue (7%).

In the multicenter MCL-002 (SPRINT) study, 254 patients with relapsed/refractory MCL ineligible for intensive chemotherapy or HCT were randomized to receive lenalidomide ($n = 170$) or investigator's choice monotherapy ($n = 84$).⁸¹ At a median follow-up of 16 months, lenalidomide significantly improved PFS compared with investigator's choice

monotherapy (9 months vs. 5 months; $P = .004$). Neutropenia without increased risk of infection (73% vs. 34%), thrombocytopenia (18% vs. 28%), leucopenia (8% vs. 11%), and anemia (8% vs. 7%) were the most common grade 3–4 adverse events.

Lenalidomide plus rituximab is also well tolerated and effective for patients with relapsed or refractory MCL. In a phase I/II study of 52 patients with relapsed or refractory MCL, lenalidomide in combination with rituximab resulted in an ORR of 57% (36% CR).⁸³ The median duration of response, PFS, and OS were 19 months, 11 months, and 24 months, respectively. The most common grade 3 or 4 toxicities included neutropenia (66%) and thrombocytopenia (23%).

Venetoclax

This recommendation is based on the results from a phase I study that demonstrated the safety and efficacy of venetoclax in relapsed or refractory NHL (106 patients; 28 patients had MCL).⁸⁴ Venetoclax resulted in responses across all NHL subtypes (ORR of 44%) and the estimated median PFS for all patients was 6 months. The ORR (75%; CR, 21%) and median PFS (14 months) were higher for patients with MCL than other NHL subtypes. The estimated 12-month OS was 70% for all patients and 82% for those with MCL.

Patients with high tumor burden, particularly those with MCL, are at increased risk for tumor lysis syndrome (TLS) and may be best managed with a starting dose of 20 mg daily for one week, and gradually escalate to target dose of 400 mg daily over 5 weeks to reduce the risk of TLS.

Other Recommended Regimens

Ibrutinib, Lenalidomide, and Rituximab

The panel has included this regimen with a category 2B recommendation based on the preliminary report from a multicenter phase II study of 50 patients with relapsed or refractory MCL in which the combination of



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

ibrutinib, lenalidomide, and rituximab resulted in an ORR of 76% (56% CR; 20% PR) at a median follow-up of 18 months.⁸⁵

Venetoclax + Ibrutinib

The panel has included the combination of venetoclax + ibrutinib as an option with a category 2B recommendation based on the results of a phase II study that showed improved outcomes in patients with MCL and poor prognostic factors (eg, *TP53* aberrations and high-risk prognostic score).⁸⁶ In this study, the CR rate of 42% as assessed by CT scan at 16 weeks was higher than the historical CR rate of 9% with ibrutinib monotherapy ($P < .001$) and 78% of patients with responding disease were estimated to have an ongoing response at 15 months.

Extended Response Duration to Prior Chemoimmunotherapy (> expected median PFS)

Preferred Regimens

Bendamustine ± Rituximab (if not previously given)

Bendamustine, as a single agent or in combination with rituximab (BR), has also shown promising results with acceptable toxicity in patients with relapsed or refractory MCL.^{87,88}

In a phase II multicenter study, BR resulted in an ORR of 92% (41% CR) in patients with relapsed or refractory indolent lymphomas and MCL ($n = 67$).⁸⁷ The median duration of response and PFS was 21 months and 23 months, respectively. The ORR was 92% (42% CR) and the median duration of response was 19 months for the subgroup of patients with MCL ($n = 12$).⁸⁷

The phase III randomized trial from StiL demonstrated superiority of the BR regimen over fludarabine and rituximab in patients with relapsed/refractory follicular or indolent lymphoma or MCL (219 evaluable patients; MCL histology in about 20%).⁸⁸ At a median follow-up of 96

months, the median PFS was 34 months and 12 months, respectively, for BR and fludarabine plus rituximab ($P < .0001$).

Bortezomib ± Rituximab

Bortezomib is a proteasome inhibitor with activity in patients with relapsed or refractory MCL and is currently approved for this indication.^{89,90} In the pivotal phase II PINNACLE trial ($n = 155$; 141 evaluable patients), bortezomib induced an ORR of 33% (CR in 8%), with a median duration of response of 9 months.⁸⁹ Median time to progression (in all patients) was 6 months. Longer follow-up data also confirmed these initial findings; after a median follow-up time of 26 months, the median OS in all patients was 24 months (35 months in patients with responding disease).⁹⁰

The efficacy of bortezomib + rituximab in patients with heavily pretreated relapsed or refractory MCL was demonstrated in a phase II study (11 patients with FL and 14 patients with MCL).⁹¹ The ORR was 40% in all 25 patients (29% in patients with MCL). The estimated 2-year PFS rate was 24% for all patients and 60% in patients with responding disease.

Other Recommended Regimens

The panel has included prednisone, etoposide, procarbazine, and cyclophosphamide (PEP-C) oral combination chemotherapy regimen and bendamustine, bortezomib, and rituximab options with a category 2B recommendation.^{92,93} In a single-center study of 22 patients with heavily pretreated MCL, PEP-C regimen was well tolerated resulting in an ORR of 82% (46% CR; 36% PR).⁹² The efficacy of bendamustine, bortezomib, and rituximab was established in a multicenter phase 2 study of 30 patients (7 patients with MCL)⁹³ and this regimen is currently being evaluated in a larger randomized trial.

Small-molecule inhibitors (as listed above for patients with short response duration to prior chemoimmunotherapy), RCHOP (if not previously given), and VR-CAP (if not previously given) are included as alternate options.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Second-Line Consolidation Therapy

Allogeneic HCT using reduced-intensity conditioning (RIC) has been evaluated as a consolidation strategy for patients in remission following treatment for relapsed/refractory MCL.^{57,94-96}

In an analysis of patients with MCL treated with HCT at the MD Anderson Cancer Center, the subgroup of patients with relapsed/refractory disease treated with RIC allogeneic stem cell transplant (SCT) (n = 35) had favorable long-term outcomes.⁵⁷ Most of these patients (62%) received HCT after achieving remission (31% in second remission). The median PFS was 60 months, and the 6-year PFS and OS rates were 46% and 53%, respectively. The transplant-related mortality (TRM) rates at 3 months and 1 year were 0% and 9%, respectively.⁵⁷

In a small study of 18 patients with relapsed MCL treated with allogeneic HCT using RIC, the 3-year PFS rate and estimated 3-year OS rate were 82% and 86%, respectively; the majority of patients in this study (89%) had chemosensitive disease.⁹⁴ In another study that evaluated allogeneic SCT using RIC in 33 patients with relapsed/refractory MCL (42% of these patients had failed prior HDT/ASCR), the 2-year DFS and OS rates were 60% and 65%, respectively.⁹⁵ The 2-year relapse rate and the 2-year TRM rate were 9% and 24% respectively; moreover, with a median follow-up of nearly 25 months, none of the 13 patients who received HCT after achieving CR experienced disease relapse.

The results of another retrospective analysis (70 patients; 35 in CR, 20 in PR, and 15 in stable or progressive disease) suggest that allogeneic HCT using RIC may be an effective therapy in patients with relapsed or refractory disease that is chemosensitive disease at time of HCT.⁹⁶ At a median follow-up of 24 months, the 2-year EFS and OS rates were 50% and 53%, respectively. Disease status at transplantation was the only significant predictor of EFS and OS.

Remission duration after autologous HCT has been identified as the only significant predictor of outcome following allogeneic HCT in patients with relapsed MCL.⁹⁷ Longer remission duration (>12 months) after autologous HCT was associated with significantly better outcomes compared to early relapse (within 1 year after autologous HCT) or primary refractory disease.

Allogeneic HCT (with myeloablative conditioning or RIC) should be considered for patients with relapsed or refractory disease that is in remission following second-line therapy.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

References

1. Teras LR, DeSantis CE, Cerhan JR, et al. 2016 US lymphoid malignancy statistics by World Health Organization subtypes. *CA Cancer J Clin* 2016;66:443-459. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27618563>.
2. Teodorovic I, Pittaluga S, Kluin-Nelemans JC, et al. Efficacy of four different regimens in 64 mantle-cell lymphoma cases: clinicopathologic comparison with 498 other non-Hodgkin's lymphoma subtypes. *European Organization for the Research and Treatment of Cancer Lymphoma Cooperative Group. J Clin Oncol* 1995;13:2819-2826. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7595744>.
3. Yatabe Y, Suzuki R, Tobinai K, et al. Significance of cyclin D1 overexpression for the diagnosis of mantle cell lymphoma: a clinicopathologic comparison of cyclin D1-positive MCL and cyclin D1-negative MCL-like B-cell lymphoma. *Blood* 2000;95:2253-2261. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10733493>.
4. Hartmann EM, Campo E, Wright G, et al. Pathway discovery in mantle cell lymphoma by integrated analysis of high-resolution gene expression and copy number profiling. *Blood* 2010;116:953-961. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20421449>.
5. Iqbal J, Shen Y, Liu Y, et al. Genome-wide miRNA profiling of mantle cell lymphoma reveals a distinct subgroup with poor prognosis. *Blood* 2012;119:4939-4948. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22490335>.
6. Fu K, Weisenburger DD, Greiner TC, et al. Cyclin D1-negative mantle cell lymphoma: a clinicopathologic study based on gene expression profiling. *Blood* 2005;106:4315-4321. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16123218>.
7. Wlodarska I, Dierickx D, Vanhentenrijk V, et al. Translocations targeting CCND2, CCND3, and MYCN do occur in t(11;14)-negative mantle cell lymphomas. *Blood* 2008;111:5683-5690. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18391076>.
8. Salaverria I, Royo C, Carvajal-Cuenca A, et al. CCND2 rearrangements are the most frequent genetic events in cyclin D1(-) mantle cell lymphoma. *Blood* 2013;121:1394-1402. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23255553>.
9. U.S. National Library of Medicine Key MEDLINE® Indicators Available at: http://www.nlm.nih.gov/bsd/bsd_key.html.
10. Avet-Loiseau H, Garand R, Gaillard F, et al. Detection of t(11;14) using interphase molecular cytogenetics in mantle cell lymphoma and atypical chronic lymphocytic leukemia. *Genes Chromosomes Cancer* 1998;23:175-182. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9739021>.
11. Determann O, Hoster E, Ott G, et al. Ki-67 predicts outcome in advanced-stage mantle cell lymphoma patients treated with anti-CD20 immunochemotherapy: results from randomized trials of the European MCL Network and the German Low Grade Lymphoma Study Group. *Blood* 2008;111:2385-2387. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18077791>.
12. Garcia M, Romaguera JE, Inamdar KV, et al. Proliferation predicts failure-free survival in mantle cell lymphoma patients treated with rituximab plus hyperfractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone alternating with rituximab plus high-dose methotrexate and cytarabine. *Cancer* 2009;115:1041-1048. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19170236>.
13. Hsi ED, Jung S, Lai R, et al. Ki67 and PIM1 expression predict outcome in mantle cell lymphoma treated with high dose therapy, stem cell transplantation and rituximab: a Cancer and Leukemia Group B 59909 correlative science study. *Leuk Lymphoma* 2008;49:2081-2090. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19021050>.
14. Schaffel R, Hedvat CV, Teruya-Feldstein J, et al. Prognostic impact of proliferative index determined by quantitative image analysis and the International Prognostic Index in patients with mantle cell lymphoma. *Ann Oncol* 2010;21:133-139. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20019090>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

15. Tiemann M, Schrader C, Klapper W, et al. Histopathology, cell proliferation indices and clinical outcome in 304 patients with mantle cell lymphoma (MCL): a clinicopathological study from the European MCL Network. *Br J Haematol* 2005;131:29-38. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16173960>.

16. Mozos A, Royo C, Hartmann E, et al. SOX11 expression is highly specific for mantle cell lymphoma and identifies the cyclin D1-negative subtype. *Haematologica* 2009;94:1555-1562. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19880778>.

17. Chen YH, Gao J, Fan G, Peterson LC. Nuclear expression of SOX11 is highly associated with mantle cell lymphoma but is independent of t(11;14)(q13;q32) in non-mantle cell B-cell neoplasms. *Mod Pathol* 2010;23:105-112. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19801969>.

18. Zeng W, Fu K, Quintanilla-Fend L, et al. Cyclin D1-negative blastoid mantle cell lymphoma identified by SOX11 expression. *Am J Surg Pathol* 2012;36:214-219. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22251940>.

19. Eskelund CW, Dahl C, Hansen JW, et al. TP53 mutations identify younger mantle cell lymphoma patients who do not benefit from intensive chemoimmunotherapy. *Blood* 2017;130:1903-1910. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28819011>.

20. Fernandez V, Salameo O, Espinet B, et al. Genomic and gene expression profiling defines indolent forms of mantle cell lymphoma. *Cancer Res* 2010;70:1408-1418. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20124476>.

21. Ondrejka SL, Lai R, Smith SD, Hsi ED. Indolent mantle cell leukemia: a clinicopathological variant characterized by isolated lymphocytosis, interstitial bone marrow involvement, kappa light chain restriction, and good prognosis. *Haematologica* 2011;96:1121-1127. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21508124>.

22. Navarro A, Clot G, Royo C, et al. Molecular subsets of mantle cell lymphoma defined by the IGHV mutational status and SOX11 expression have distinct biologic and clinical features. *Cancer Res* 2012;72:5307-5316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22915760>.

23. Swerdlow SH, Harris NL, Jaffe ES, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Revised 4th ed. Lyon, France: IARC; 2017.

24. Koletsa T, Markou K, Ouzounidou S, et al. In situ mantle cell lymphoma in the nasopharynx. *Head Neck* 2012. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23280758>.

25. Richard P, Vassallo J, Valmary S, et al. "In situ-like" mantle cell lymphoma: a report of two cases. *J Clin Pathol* 2006;59:995-996. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16935977>.

26. Carvajal-Cuenca A, Sua LF, Silva NM, et al. In situ mantle cell lymphoma: clinical implications of an incidental finding with indolent clinical behavior. *Haematologica* 2012;97:270-278. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22058203>.

27. Edlefsen KL, Greisman HA, Yi HS, et al. Early lymph node involvement by mantle cell lymphoma limited to the germinal center: report of a case with a novel "follicular in situ" growth pattern. *Am J Clin Pathol* 2011;136:276-281. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21757601>.

28. Adam P, Schiefer AI, Prill S, et al. Incidence of preclinical manifestations of mantle cell lymphoma and mantle cell lymphoma in situ in reactive lymphoid tissues. *Mod Pathol* 2012;25:1629-1636. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22790016>.

29. Carbone A, Santoro A. How I treat: diagnosing and managing "in situ" lymphoma. *Blood* 2011;117:3954-3960. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21224472>.

30. Romaguera JE, Medeiros LJ, Hagemeister FB, et al. Frequency of gastrointestinal involvement and its clinical significance in mantle cell



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

lymphoma. Cancer 2003;97:586-591. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12548600>.

31. Salar A, Juanpere N, Bellosillo B, et al. Gastrointestinal involvement in mantle cell lymphoma: a prospective clinic, endoscopic, and pathologic study. Am J Surg Pathol 2006;30:1274-1280. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17001159>.

32. Leitch HA, Gascoyne RD, Chhanabhai M, et al. Limited-stage mantle-cell lymphoma. Ann Oncol 2003;14:1555-1561. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/14504058>.

33. Martin P, Chadburn A, Christos P, et al. Outcome of deferred initial therapy in mantle-cell lymphoma. J Clin Oncol 2009;27:1209-1213. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19188674>.

34. Cohen JB, Han X, Jemal A, et al. Deferred therapy is associated with improved overall survival in patients with newly diagnosed mantle cell lymphoma. Cancer 2016;122:2356-2363. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/27153197>.

35. Martin P, Chadburn A, Christos P, et al. Intensive treatment strategies may not provide superior outcomes in mantle cell lymphoma: overall survival exceeding 7 years with standard therapies. Ann Oncol 2008;19:1327-1330. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18349031>.

36. Lacasce AS, Vandergrift JL, Rodriguez MA, et al. Comparative outcome of initial therapy for younger patients with mantle cell lymphoma: an analysis from the NCCN NHL Database. Blood 2012;119:2093-2099. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22234679>.

37. Le Gouill S, Thieblemont C, Oberic L, et al. Rituximab after Autologous Stem-Cell Transplantation in Mantle-Cell Lymphoma. N Engl J Med 2017;377:1250-1260. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28953447>.

38. Le Gouill S, Thieblemont C, Oberic L, et al. R-DHA-Oxaliplatin before Autologous Stem Cell Transplantation Prolongs PFS and OS As

Compared to R-DHA-Carboplatin and R-DHA-Cisplatin in Patients with Mantle Cell Lymphoma, a Subgroup Analysis of the LyMa Trial [abstract]. Blood 2017;130:Abstract 1496. Available at:
http://www.bloodjournal.org/content/130/Suppl_1/1496.abstract.

39. Hermine O, Hoster E, Walewski J, et al. Addition of high-dose cytarabine to immunochemotherapy before autologous stem-cell transplantation in patients aged 65 years or younger with mantle cell lymphoma (MCL Younger): a randomised, open-label, phase 3 trial of the European Mantle Cell Lymphoma Network. Lancet 2016;388:565-575. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27313086>.

40. Delarue R, Haioun C, Ribrag V, et al. CHOP and DHAP plus rituximab followed by autologous stem cell transplantation in mantle cell lymphoma: a phase 2 study from the Groupe d'Etude des Lymphomes de l'Adulte. Blood 2013;121:48-53. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22718839>.

41. Geisler CH, Kolstad A, Laurell A, et al. Long-term progression-free survival of mantle cell lymphoma after intensive front-line immunochemotherapy with in vivo-purged stem cell rescue: a nonrandomized phase 2 multicenter study by the Nordic Lymphoma Group. Blood 2008;112:2687-2693. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18625886>.

42. Eskelund CW, Kolstad A, Jerkeman M, et al. 15-year follow-up of the Second Nordic Mantle Cell Lymphoma trial (MCL2): prolonged remissions without survival plateau. Br J Haematol 2016;175:410-418. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/27378674>.

43. Geisler CH, Kolstad A, Laurell A, et al. The Mantle Cell Lymphoma International Prognostic Index (MIPI) is superior to the International Prognostic Index (IPI) in predicting survival following intensive first-line immunochemotherapy and autologous stem cell transplantation (ASCT). Blood 2010;115:1530-1533. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/20032504>.

44. Romaguera JE, Fayad L, Rodriguez MA, et al. High rate of durable remissions after treatment of newly diagnosed aggressive mantle-cell



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

lymphoma with rituximab plus hyper-CVAD alternating with rituximab plus high-dose methotrexate and cytarabine. *J Clin Oncol* 2005;23:7013-7023. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16145068>.

45. Romaguera JE, Fayad LE, Feng L, et al. Ten-year follow-up after intense chemoimmunotherapy with Rituximab-HyperCVAD alternating with Rituximab-high dose methotrexate/cytarabine (R-MA) and without stem cell transplantation in patients with untreated aggressive mantle cell lymphoma. *Br J Haematol* 2010;150:200-208. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20528872>.

46. Merli F, Luminari S, Ilariucci F, et al. Rituximab plus HyperCVAD alternating with high dose cytarabine and methotrexate for the initial treatment of patients with mantle cell lymphoma, a multicentre trial from Gruppo Italiano Studio Linfomi. *Br J Haematol* 2012;156:346-353. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22145911>.

47. Bernstein SH, Epner E, Unger JM, et al. A phase II multicenter trial of hyperCVAD MTX/Ara-C and rituximab in patients with previously untreated mantle cell lymphoma; SWOG 0213. *Ann Oncol* 2013;24:1587-1593. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23504948>.

48. Wang ML, Lee H, Thirumurthi S, et al. Ibrutinib-rituximab followed by reduced chemo-immunotherapy consolidation in young, newly diagnosed mantle cell lymphoma patients: a window of opportunity to reduce chemo. *Hematological Oncology* 2017;35:142-143. Available at: <https://onlinelibrary.wiley.com/doi/abs/10.1002/hon.2437> 132.

49. Rummel MJ, Niederle N, Maschmeyer G, et al. Bendamustine plus rituximab versus CHOP plus rituximab as first-line treatment for patients with indolent and mantle-cell lymphomas: an open-label, multicentre, randomised, phase 3 non-inferiority trial. *Lancet* 2013;381:1203-1210. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23433739>.

50. Flinn IW, van der Jagt R, Kahl B, et al. First-Line Treatment of Patients With Indolent Non-Hodgkin Lymphoma or Mantle-Cell Lymphoma With Bendamustine Plus Rituximab Versus R-CHOP or R-CVP: Results of the BRIGHT 5-Year Follow-Up Study. *J Clin Oncol* 2019;37:984-991. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30811293>.

51. Khouri IF, Romaguera J, Kantarjian H, et al. Hyper-CVAD and high-dose methotrexate/cytarabine followed by stem-cell transplantation: an active regimen for aggressive mantle-cell lymphoma. *J Clin Oncol* 1998;16:3803-3809. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9850025>.

52. Khouri IF, Saliba RM, Okoroji GJ, et al. Long-term follow-up of autologous stem cell transplantation in patients with diffuse mantle cell lymphoma in first disease remission: the prognostic value of beta2-microglobulin and the tumor score. *Cancer* 2003;98:2630-2635. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14669282>.

53. Dreyling M, Lenz G, Hoster E, et al. Early consolidation by myeloablative radiochemotherapy followed by autologous stem cell transplantation in first remission significantly prolongs progression-free survival in mantle-cell lymphoma: results of a prospective randomized trial of the European MCL Network. *Blood* 2005;105:2677-2684. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15591112>.

54. Thieblemont C, Antal D, Lacotte-Thierry L, et al. Chemotherapy with rituximab followed by high-dose therapy and autologous stem cell transplantation in patients with mantle cell lymphoma. *Cancer* 2005;104:1434-1441. Available at: <http://www.ncbi.nlm.nih.gov/PubMed/16104036>.

55. Ritchie DS, Seymour JF, Grigg AP, et al. The hyper-CVAD-rituximab chemotherapy programme followed by high-dose busulfan, melphalan and autologous stem cell transplantation produces excellent event-free survival in patients with previously untreated mantle cell lymphoma. *Ann Hematol* 2007;86:101-105. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17089127>.

56. Till BG, Gooley TA, Crawford N, et al. Effect of remission status and induction chemotherapy regimen on outcome of autologous stem cell transplantation for mantle cell lymphoma. *Leuk Lymphoma* 2008;49:1062-1073. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18452065>.

57. Tam CS, Bassett R, Ledesma C, et al. Mature results of the M. D. Anderson Cancer Center risk-adapted transplantation strategy in mantle



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

cell lymphoma. Blood 2009;113:4144-4152. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19168784>.

58. van 't Veer MB, de Jong D, MacKenzie M, et al. High-dose Ara-C and beam with autograft rescue in R-CHOP responsive mantle cell lymphoma patients. Br J Haematol 2009;144:524-530. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19036081>.

59. Graf SA, Stevenson PA, Holmberg LA, et al. Maintenance rituximab after autologous stem cell transplantation in patients with mantle cell lymphoma. Ann Oncol 2015;26:2323-2328. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/26347113>.

60. Klener P, Salek D, Pytlik R, et al. Rituximab maintenance significantly prolongs progression-free survival of patients with newly diagnosed mantle cell lymphoma treated with the Nordic MCL2 protocol and autologous stem cell transplantation. Am J Hematol 2019;94:E50-E53. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/30474171>.

61. Robak T, Jin J, Pylypenko H, et al. Frontline bortezomib, rituximab, cyclophosphamide, doxorubicin, and prednisone (VR-CAP) versus rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisone (R-CHOP) in transplantation-ineligible patients with newly diagnosed mantle cell lymphoma: final overall survival results of a randomised, open-label, phase 3 study. Lancet Oncol 2018;19:1449-1458. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/30348538>.

62. Howard OM, Gribben JG, Neuberg DS, et al. Rituximab and CHOP induction therapy for newly diagnosed mantle-cell lymphoma: molecular complete responses are not predictive of progression-free survival. J Clin Oncol 2002;20:1288-1294. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/11870171>.

63. Lenz G, Dreyling M, Hoster E, et al. Immunochemotherapy with rituximab and cyclophosphamide, doxorubicin, vincristine, and prednisone significantly improves response and time to treatment failure, but not long-term outcome in patients with previously untreated mantle cell lymphoma: results of a prospective randomized trial of the German Low Grade

Lymphoma Study Group (GLSG). J Clin Oncol 2005;23:1984-1992. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15668467>.

64. Kluin-Nelemans HC, Hoster E, Hermine O, et al. Treatment of older patients with mantle-cell lymphoma. N Engl J Med 2012;367:520-531. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22873532>.

65. Ruan J, Martin P, Shah B, et al. Lenalidomide plus Rituximab as Initial Treatment for Mantle-Cell Lymphoma. N Engl J Med 2015;373:1835-1844. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26535512>.

66. Ruan J, Martin P, Christos P, et al. Five-year follow-up of lenalidomide plus rituximab as initial treatment for mantle cell lymphoma. Blood 2018;132:2016-2025. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/30181173>.

67. Kahl BS, Longo WL, Eickhoff JC, et al. Maintenance rituximab following induction chemoimmunotherapy may prolong progression-free survival in mantle cell lymphoma: a pilot study from the Wisconsin Oncology Network. Ann Oncol 2006;17:1418-1423. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16766582>.

68. Chang JE, Li H, Smith MR, et al. Phase II study of VcR-CVAD with maintenance rituximab for untreated mantle cell lymphoma: an Eastern Cooperative Oncology Group Study (E1405). Blood 2014;123:1665-1673. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24458437>.

69. Visco C, Finotto S, Zambello R, et al. Combination of rituximab, bendamustine, and cytarabine for patients with mantle-cell non-Hodgkin lymphoma ineligible for intensive regimens or autologous transplantation. J Clin Oncol 2013;31:1442-1449. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23401442>.

70. Hoster E, Kluin-Nelemans H, Hermine O, et al. Rituximab Maintenance after First-Line Immunochemotherapy in Mantle Cell Lymphoma: Long-Term Follow-up of the Randomized European MCL Elderly Trial [abstract]. Blood 2017;130 (Suppl 1):Abstract 153. Available at:
http://www.bloodjournal.org/content/130/Suppl_1/153.abstract.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

71. Chang JE, Carmichael LL, Kim K, et al. VcR-CVAD Induction Chemotherapy Followed by Maintenance Rituximab Produces Durable Remissions in Mantle Cell Lymphoma: A Wisconsin Oncology Network Study. Clin Lymphoma Myeloma Leuk 2018;18:e61-e67. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29191715>.

72. Rummel MJ, Knauf W, Goerner M, et al. Two years rituximab maintenance vs. observation after first-line treatment with bendamustine plus rituximab (B-R) in patients with mantle cell lymphoma: First results of a prospective, randomized, multicenter phase II study (a subgroup study of the StiL NHL7-2008 MAINTAIN trial) [abstract]. J Clin Oncol 2016;34:Abstract 7503. Available at: https://doi.org/10.1200/JCO.2016.34.15_suppl.7503.

73. Wang M, Rule S, Zinzani PL, et al. Acalabrutinib in relapsed or refractory mantle cell lymphoma (ACE-LY-004): a single-arm, multicentre, phase 2 trial. Lancet 2018;391:659-667. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29241979>.

74. Wang M, Rule S, Zinzani PL, et al. Long-Term Follow-up of Acalabrutinib Monotherapy in Patients with Relapsed/Refractory Mantle Cell Lymphoma [abstract]. Blood 2018;132:Abstract 2876. Available at: http://www.bloodjournal.org/content/132/Suppl_1/2876.abstract.

75. Wang ML, Blum KA, Martin P, et al. Long-term follow-up of MCL patients treated with single-agent ibrutinib: updated safety and efficacy results. Blood 2015;126:739-745. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26059948>.

76. Rule S, Jurczak W, Jerkeman M, et al. Ibrutinib versus temsirolimus: 3-year follow-up of patients with previously treated mantle cell lymphoma from the phase 3, international, randomized, open-label RAY study. Leukemia 2018;32:1799-1803. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29572505>.

77. Jain P, Romaguera J, Srour SA, et al. Four-year follow-up of a single arm, phase II clinical trial of ibrutinib with rituximab (IR) in patients with relapsed/refractory mantle cell lymphoma (MCL). Br J Haematol

2018;182:404-411. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29785709>.

78. Song Y, Zhou K, Zou D, et al. Zanubrutinib in patients with relapsed/refractory mantle cell lymphoma [abstract]. Hematological Oncology 2019;37:45-46. Available at: https://onlinelibrary.wiley.com/doi/abs/10.1002/hon.15_2629.

79. Tam CS, Wang M, Simpson D, et al. Updated safety and efficacy data in the phase 1 trial of patients with mantle cell lymphoma (mcl) treated with bruton tyrosine kinase (BTK) inhibitor zanubrutinib (BGB-3111) [abstract]. Hematological Oncology 2019;37:245-247. Available at: https://onlinelibrary.wiley.com/doi/abs/10.1002/hon.55_2630.

80. Goy A, Sinha R, Williams ME, et al. Single-agent lenalidomide in patients with mantle-cell lymphoma who relapsed or progressed after or were refractory to bortezomib: phase II MCL-001 (EMERGE) study. J Clin Oncol 2013;31:3688-3695. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24002500>.

81. Trneny M, Lamy T, Walewski J, et al. Lenalidomide versus investigator's choice in relapsed or refractory mantle cell lymphoma (MCL-002; SPRINT): a phase 2, randomised, multicentre trial. Lancet Oncol 2016;17:319-331. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26899778>.

82. Witzig TE, Luigi Zinzani P, Habermann TM, et al. Long-term analysis of phase II studies of single-agent lenalidomide in relapsed/refractory mantle cell lymphoma. Am J Hematol 2017;92:E575-E583. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28699256>.

83. Wang M, Fayad L, Wagner-Bartak N, et al. Lenalidomide in combination with rituximab for patients with relapsed or refractory mantle-cell lymphoma: a phase 1/2 clinical trial. Lancet Oncol 2012;13:716-723. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22677155>.

84. Davids MS, Roberts AW, Seymour JF, et al. Phase I First-in-Human Study of Venetoclax in Patients With Relapsed or Refractory Non-Hodgkin



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Lymphoma. J Clin Oncol 2017;35:826-833. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28095146>.

85. Jerkeman M, Eskelund CW, Hutchings M, et al. Ibrutinib, lenalidomide, and rituximab in relapsed or refractory mantle cell lymphoma (PHILEMON): a multicentre, open-label, single-arm, phase 2 trial. Lancet Haematol 2018;5:e109-e116. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29396091>.

86. Tam CS, Anderson MA, Pott C, et al. Ibrutinib plus Venetoclax for the Treatment of Mantle-Cell Lymphoma. N Engl J Med 2018;378:1211-1223. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29590547>.

87. Robinson KS, Williams ME, van der Jagt RH, et al. Phase II multicenter study of bendamustine plus rituximab in patients with relapsed indolent B-cell and mantle cell non-Hodgkin's lymphoma. J Clin Oncol 2008;26:4473-4479. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18626004>.

88. Rummel M, Kaiser U, Balser C, et al. Bendamustine plus rituximab versus fludarabine plus rituximab for patients with relapsed indolent and mantle-cell lymphomas: a multicentre, randomised, open-label, non-inferiority phase 3 trial. Lancet Oncol 2016;17:57-66. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26655425>.

89. Fisher RI, Bernstein SH, Kahl BS, et al. Multicenter phase II study of bortezomib in patients with relapsed or refractory mantle cell lymphoma. J Clin Oncol 2006;24:4867-4874. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17001068>.

90. Goy A, Bernstein SH, Kahl BS, et al. Bortezomib in patients with relapsed or refractory mantle cell lymphoma: updated time-to-event analyses of the multicenter phase 2 PINNACLE study. Ann Oncol 2009;20:520-525. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19074748>.

91. Baiocchi RA, Alinari L, Lustberg ME, et al. Phase 2 trial of rituximab and bortezomib in patients with relapsed or refractory mantle cell and

follicular lymphoma. Cancer 2011;117:2442-2451. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24048792>.

92. Coleman M, Martin P, Ruan J, et al. Low-dose metronomic, multidrug therapy with the PEP-C oral combination chemotherapy regimen for mantle cell lymphoma. Leuk Lymphoma 2008;49:447-450. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18297520>.

93. Friedberg JW, Vose JM, Kelly JL, et al. The combination of bendamustine, bortezomib, and rituximab for patients with relapsed/refractory indolent and mantle cell non-Hodgkin lymphoma. Blood 2011;117:2807-2812. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21239695>.

94. Khouri IF, Lee MS, Saliba RM, et al. Nonablative allogeneic stem-cell transplantation for advanced/recurrent mantle-cell lymphoma. J Clin Oncol 2003;21:4407-4412. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14645431>.

95. Maris MB, Sandmaier BM, Storer BE, et al. Allogeneic hematopoietic cell transplantation after fludarabine and 2 Gy total body irradiation for relapsed and refractory mantle cell lymphoma. Blood 2004;104:3535-3542. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15304387>.

96. Le Gouill S, Kroger N, Dhedin N, et al. Reduced-intensity conditioning allogeneic stem cell transplantation for relapsed/refractory mantle cell lymphoma: a multicenter experience. Ann Oncol 2012;23:2695-2703. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22440229>.

97. Dietrich S, Boumendi A, Finel H, et al. Outcome and prognostic factors in patients with mantle-cell lymphoma relapsing after autologous stem-cell transplantation: a retrospective study of the European Group for Blood and Marrow Transplantation (EBMT). Ann Oncol 2014;25:1053-1058. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24585719>.



Diffuse Large B-Cell Lymphoma

Diffuse large B-cell lymphomas (DLBCL) are the most common lymphoid neoplasms in adults, accounting for approximately 30% of non-Hodgkin's lymphomas (NHLs) diagnosed annually.¹ DLBCL not otherwise specified (NOS), follicular lymphoma (FL; grade 3 only), DLBCL coexistent with a low grade lymphoma of any kind (e.g., FL, gastric MALT lymphoma, or non-gastric MALT lymphoma), intravascular large B-cell lymphoma, DLBCL-associated with chronic inflammation, anaplastic lymphoma kinase (ALK)-positive DLBCL, Epstein-Barr virus (EBV)-positive DLBCL in older patients, and T-cell/histiocyte-rich large B-cell lymphoma are also managed according to the DLBCL guidelines.

Literature Search Criteria and Guidelines Update Methodology

Prior to the update of this version of the NCCN Guidelines® for B-cell Lymphomas an electronic search of the PubMed database was performed to obtain key literature in DLBCL published since the previous Guidelines update, using the following search terms: diffuse large B-cell lymphoma, aggressive B-cell lymphoma, primary mediastinal B-cell lymphoma, grey zone lymphoma and high grade B-cell lymphoma. The PubMed database was chosen as it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.²

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.

The data from key PubMed articles deemed as relevant to these Guidelines and discussed by the panel have been included in this version of the Discussion section (eg, e-publications ahead of print, meeting

abstracts). Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.

The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

Diagnosis

Gene expression profiling (GEP) has identified distinct subtypes within DLBCL NOS based on cell of origin (COO): germinal center B-cell (GCB) subtype and activated B-cell (ABC) subtype.^{3,4} Notably, GEP is not currently available in the clinic and remains utilized only in the context of clinical trials. Immunohistochemistry (IHC) algorithms have been developed as surrogates for GEP to assign COO in clinical practice and can generally assign tumors as GCB or non-GCB. The term non-GCB is used when classified by IHC because specific IHC markers for activated B-cells are not routinely available, and thus the non-GCB category is more heterogeneous than true ABC DLBCL when assigned by GEP. The most commonly used IHC algorithms include CD10, BCL6, and IRF4/MUM1 to classify DLBCL into GCB (CD10+; or BCL6+, IRF4/MUM1-) and non-GCB (CD10-, IRF4/MUM1+; or BCL6-, IRF4/MUM1-).⁵ IHC algorithms including GCET1, FOXP1, and LMO2 in addition to CD10, BCL6 and IRF4/MUM1 have been proposed.⁶⁻⁸

GCB DLBCL is associated with an improved outcome compared to ABC DLBCL in patients treated with R-CHOP, and ongoing randomized clinical trials are exploring whether the addition of novel targeted agents to R-CHOP will selectively improve the outcome in patients with ABC DLBCL.⁹⁻¹¹ Presently, the upfront standard of care remains the same for both GCB and ABC subtypes.

In the 2017 WHO classification, double-hit/triple-hit lymphomas (DHL/THL) that harbor *MYC* and *BCL2* and/or *BCL6* translocations have been included in a unique category called high-grade B-cell lymphomas (HGBL)



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

with translocations of *MYC* and *BCL2* and/or *BCL6*.⁴ This subtype is distinct from DLBCL with dual expression of both *MYC* and *BCL2* by IHC but are negative for rearrangements, the so called double expressing lymphomas (DEL). These lymphomas have an inferior prognosis compared to DLBCL as a whole, but not to the same magnitude as true double-hit lymphomas on the basis of gene rearrangements, and do not currently warrant therapy different from typical cases of DLBCL.

Adequate immunophenotyping is required to establish the diagnosis as well as to aid in risk stratification. The IHC panel includes CD20, CD3, CD5, CD10, CD45, BCL2, BCL6, Ki-67, IRF4/MUM1 and *MYC*. Patients with expression of *MYC* and either BCL2 or BCL6 by IHC should undergo FISH or karyotype testing for the detection of *MYC*, *BCL2*, and *BCL6* gene rearrangements. *MYC* gene rearrangements have been reported in 5% to 8% of patients with DLBCL, and often correlate with GCB phenotype and higher risk of progression.¹²⁻¹⁵ Additional markers such as CD138, CD30, cyclin D1, ALK1, SOX11, HHV-8 and EBV in situ hybridization (EBV-ISH) may be useful under certain circumstances to establish the subtype. SOX11 positivity may be useful in differentiating rare cases of cyclin D1-negative pleomorphic or blastoid mantle cell lymphoma (MCL) from CD5-positive DLBCL.^{16,17}

Workup

The initial workup for newly diagnosed patients with DLBCL should include a thorough physical examination with attention to node-bearing areas, and evaluation of performance status (PS) and constitutional symptoms. Laboratory assessments should include standard blood work including complete blood count (CBC) with differential, a comprehensive metabolic panel, and measurements of serum lactate dehydrogenase (LDH). Patients with high tumor burden and elevated LDH should be assessed for spontaneous tumor lysis syndrome (TLS), including measurements of uric acid, potassium, phosphorous, calcium, and renal function. Hepatitis B

virus (HBV) testing prior to initiation of treatment is recommended for all patients who will receive anti-CD20 monoclonal antibody (MAB)-based regimens due to increased risk of HBV reactivation. Human immunodeficiency virus (HIV) testing, hepatitis C virus (HCV) testing and measurement of serum beta-2-microglobulin levels would be useful in selected patients.

PET/CT scans are essential for the initial staging of DLBCL where upstaging resulting in altered therapy occurs about 9% of the time, and for response assessment after treatment because they can distinguish residual fibrotic masses from masses containing viable tumor.¹⁸ Whole body PET/CT scan ± chest/abdomen/pelvic CT with contrast of diagnostic quality is recommended for initial workup. PET/CT has also been reported to be accurate and complementary to bone marrow biopsy for the detection of bone marrow involvement in patients with newly diagnosed DLBCL.^{19,20} Bone marrow biopsy may not be needed if there is clearly positive marrow uptake by PET/CT. Bone marrow biopsy may also be omitted in the absence of any skeletal uptake on the staging PET/CT scan, unless finding another lymphoma subtype (discordant low-grade lymphoma) would be considered important for treatment decisions.

The staging workup is designed to identify all sites of known disease and determine prognosis with known clinical risk factors. International Prognostic Index (IPI) and the revised IPI (R-IPI) identify specific groups of patients who are more or less likely to be cured with standard therapy.^{21,22} IPI scores are based on patient's age, stage of disease, serum LDH level, ECOG PS, and the number of extranodal sites. In patients who are ≤60 years, an age-adjusted IPI uses the prognostic factors of stage, ECOG PS, and serum LDH level.²⁰ An NCCN-IPI has also been developed based on the outcome data from NCCN Member Institutions, and can stratify patients with newly diagnosed DLBCL into 4 different risk groups (low, low-intermediate, high-intermediate, and high) based on quantification of



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

age, LDH, sites of involvement, Ann Arbor stage and ECOG PS).²³ This analysis included 1650 patients identified in the NCCN database that were diagnosed with DLBCL between 2000 and 2010 and treated with rituximab-based therapy. The NCCN-IPI discriminated patients in the low- and high-risk subgroups better (5-year overall survival [OS] rate 96% vs 33%) than the IPI (5 year OS rate 90% vs 54%). The NCCN-IPI was also validated using an independent cohort of 1138 patients from the British Columbia Cancer Agency.²³ While the IPI, R-IPI, and NCCN-IPI predict clinical outcome with high accuracy, R-IPI and NCCN-IPI could also identify a specific subgroup of patients with very good prognosis (3-year progression-free survival [PFS] and OS of 100%).²⁴

Stage I-II

In the SWOG 0014 study that evaluated 3 cycles of R-CHOP followed by involved-field RT (IFRT) in patients with at least one adverse factor (non-bulky stage II disease, age >60 years, ECOG PS 2, or elevated serum LDH) as defined by the stage-modified IPI (n=60), the 4-year PFS rate was 88%, after a median follow-up of 5 years; the corresponding 4-year OS rate was 92%.²⁵ In historical comparison, these results were favorable relative to the survival rates for patients treated without rituximab (4-year PFS and OS rates were 78% and 88%, respectively). A phase III trial (MabThera International Trial [MINT]) compared 6 cycles of CHOP-like chemotherapy to 6 cycles of CHOP-like chemotherapy plus rituximab.²⁶ All patients were younger than 60 years of age and had 0-1 IPI risk factors. Three quarters of patients had limited stage disease, and RT was included for all extranodal sites of disease or any site >7.5 cm. The trial found a benefit to rituximab-based chemotherapy with a 6-year OS rate of 90% versus 80% ($P = .0004$). The 6-year event-free survival (EFS) rate (74% vs. 56%; $P < .0001$) and PFS rate (80% vs. 64%; $P < .0001$) were also significantly higher for patients assigned to chemotherapy plus rituximab compared to chemotherapy alone.²⁶ The results from the

RICOVER-noRT trial also showed a significant advantage to adding RT to initial bulky sites ≥ 7.5 cm.²⁷

R-CHOP (3 cycles) with RT is also associated with reduced short-term toxicity compared to 6-8 cycles of R-CHOP alone. In a SEER-Medicare database analysis of a large cohort of older patients with stage I-II DLBCL, while both treatment options had similar OS, 3 cycles of R-CHOP with RT was associated with significantly lower risk of second-line therapy and lower incidences of neutropenia including those requiring hospitalization.²⁸

The results of a prospective randomized trial showed that R-CHOP-14 was not inferior to R-CHOP-14 plus RT in patients with non-bulky limited-stage DLBCL.²⁹ In this trial, 334 patients were randomized to receive R-CHOP-14 alone (n = 165) or R-CHOP-14 plus RT (n=169). Patients with 0 IPI risk factors received 4 cycles while patients with ≥ 1 risk factor received 6. After a median follow-up of 64 months, the 5-year EFS (89% and 92%, respectively; $P = .18$) and OS (92% and 96% respectively) were not statistically significantly different between the R-CHOP-14 alone and R-CHOP-14 plus RT arms.

In the two GELA studies, intensified chemotherapy [ACVBP (doxorubicin, cyclophosphamide, vindesine, bleomycin and prednisone) followed by consolidation with methotrexate, etoposide, ifosfamide and cytarabine] with or without rituximab was found to be superior to CHOP with or without rituximab (3 cycles) plus RT in patients with low-risk early-stage disease.^{30,31} However, this regimen was also associated with significant toxicity and includes vindesine, which is not available in the United States.

R-CHOP (3 cycles) followed by involved site RT (ISRT) (category 1) or R-CHOP (6 cycles) or R-CHOP-14 (4–6 cycles) with or without ISRT are included as first-line therapy options for patients with non-bulky (<7.5 cm) disease.^{25,26,29} R-CHOP (6 cycles) with or without ISRT is recommended for patients with bulky disease (≥ 7.5 cm).^{26,27} R-mini-CHOP (decreased



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

dose of CHOP with a conventional dose of rituximab) may be substituted for very frail patients and patients >80 years of age with comorbidities to improve tolerability.^{32,33} In a prospective single arm phase II study by the GELA study group that evaluated the safety and efficacy of R-mini-CHOP in 149 patients with >80 years, the 4-year PFS and OS rates were 41% and 49%, respectively.³³ Grade ≥3 neutropenia was the most frequent hematological toxicity observed in 59 patients.

Patients with primary testicular DLBCL are at increased risk of central nervous system (CNS) and contralateral scrotal recurrence, even when presenting with stage I disease. Inclusion of methotrexate for CNS prophylaxis as well as scrotal RT (25–30 Gy) after completion of chemoimmunotherapy is therefore recommended.³⁴ ISRT alone for DLBCL is associated with a high rate of relapse and is only recommended for patients who are not candidates for any chemotherapy.

Stage III-IV

The efficacy of R-CHOP-21 in patients with advanced stage DLBCL has been demonstrated in multiple randomized trials.^{26,35-38} The GELA study (LNH98-5), evaluated 8 cycles of R-CHOP versus CHOP in older patients (age 60–80 years; n=399).³⁵ At a median follow-up of 10 years, the 10-year PFS (37% vs. 20%), disease-free survival (DFS) (64% vs. 43%), and OS (44% vs. 28%) rates were significantly higher for R-CHOP.³⁶ The MInT study (6 cycles of R-CHOP or CHOP) extended these findings to younger patients with 0 or 1 risk factors according to the IPI.²⁶ The Dutch HOVON and Nordic Lymphoma Group study (8 cycles of R-CHOP-14 or CHOP-14) and the ECOG/CALGB study confirmed the findings in patients older than 60 years.^{37,38} The ECOG/CALGB 9703 study also showed that maintenance rituximab in first CR offered no clinical benefit to patients who received R-CHOP as their induction therapy.³⁸

The RICOVER 60-trial showed that the addition of rituximab to 6 or 8 cycles of CHOP-14 (R-CHOP-14) also significantly improved clinical outcomes compared with CHOP-14 alone.^{39,40} In this trial, older patients (age 61–80 years) were randomized to receive CHOP-14 (6 or 8 cycles) with or without 8 cycles of rituximab. RT was administered to sites of initial bulky disease with or without extranodal involvement. With a median follow-up of 78 months, R-CHOP-14 was associated with significantly improved EFS and OS compared to CHOP-14 ($P < .001$).⁴⁰ While there was no difference in clinical benefit, increased toxicity was seen in patients treated with 8 cycles compared with 6 cycles of therapy. The investigators concluded that 6 cycles of R-CHOP-14 in combination with 8 doses of rituximab should be the preferred regimen in this patient population.

Two randomized trials that have compared R-CHOP-21 with R-CHOP-14 showed that while both treatment options are associated with similar OS and PFS, R-CHOP-14 was associated with significantly higher rates of grade 3 or 4 neutropenia.^{41,42} In the phase III randomized trial of 1080 patients with newly diagnosed DLBCL, at a median follow up of 46 months, the 2-year OS rates were 83% and 81% respectively for R-CHOP-14 and R-CHOP-21 ($P = .38$).⁴¹ The corresponding 2-year PFS rates were 75% for both treatment arms ($P = .59$). Notably, there was no difference in outcome between GCB-like and non-GCB-like DLBCL by IHC in this large prospective study. Toxicity was similar, except for a lower rate of grade 3 or 4 neutropenia in the R-CHOP-14 arm (31% vs. 60%), reflecting the fact that all patients in the R-CHOP-14 arm received primary growth factor prophylaxis with G-CSF whereas no primary prophylaxis given with R-CHOP-21.⁴¹ In the phase III LNH03-6B GELA study which compared R-CHOP-14 (8 cycles) with R-CHOP-21 in 602 older patients (age 60–80 years) with untreated DLBCL, after a median follow-up of 56 months, no significant differences were observed in terms of 3-year EFS (56% vs. 60%; $P = .76$), PFS (60% vs. 62%) or OS rates (69% vs 72%) between R-CHOP-14 and R-CHOP-21.⁴² Grade 3 or 4 neutropenia was



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

more frequent in the R-CHOP-14 arm (74% compared to 64% in the R-CHOP 21 arm) despite a higher proportion of patients having received G-CSF (90%) compared with patients in the R-CHOP-21 arm (66%).

The R-MegaCHOEP study reported that the 3-year PFS and OS rates of were 74% and 85% respectively, in young, high-risk patients with DLBCL treated with 8 x CHOEP-14 with 6 infusions of rituximab.⁴³ The results of the dense-R-MegaCHOEP trial showed that doubling the number of rituximab (375 mg/m²) infusions (from 6 to 12) administered with 8 x CHOEP-14 did not result in a significant improvement of EFS and OS in patients with age-adjusted IPI of 2.⁴⁴ After a median follow-up of 24 months, the 2-year EFS and OS rates were 69% and 82% respectively. The results of the PETAL trial also confirmed that the addition of 2 more doses of rituximab to 6 x R-CHOP (6 vs 8 doses of rituximab) did not improve outcome.⁴⁵

Dose-adjusted EPOCH-R (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin and rituximab; DA-EPOCH-R) has shown significant activity in patients with untreated DLBCL with 5-year and 1-year OS rates of 84% and 64%, respectively in a phase II trial.^{46,47} A phase III randomized trial (CALGB 50303) evaluated DA-EPOCH-R versus R-CHOP in untreated patients with DLBCL.⁴⁸ In this study, 524 patients were randomly assigned to 6 cycles of R-CHOP (n = 223) or DA-EPOCH-R (n = 232). The ORR was 89% in both arms, and after a median follow-up of 5 years, there were no statistically significant differences in the 2-year PFS (79% vs. 76%; *P* = .65) or 2-year OS rates (87% and 86%; *P* = .42) between the R-CHOP and DA-EPOCH-R. In addition, R-CHOP also had a more favorable safety and tolerability profile. DA-EPOCH-R was associated with a significantly increased risk of cytopenias and neuropathy.

Collectively, available data from clinical trials discussed above suggest that R-CHOP administered on a 21 day schedule remains the standard

treatment regimen for the majority of patients with newly diagnosed DLBCL.

R-CHOP for a total of 6 cycles (category 1) is the preferred regimen for patients with stage III-IV disease due to reduced toxicities compared to other regimens.^{26,35-38} In selected patients, RT to bulky sites may be beneficial (category 2B). DA-EPOCH-R (category 2B)^{46,47} or dose-dense R-CHOP-14 (category 3)^{41,42} are included as alternative regimens in selected patients. DA-EPOCH-R may be preferred for patients with primary mediastinal large B-cell lymphoma (PMBL) or with HGBL with translocations of *MYC* and *BCL2* and/or *BCL6* (DHL/THL). There is no evidence to suggest that an intensified regimen is better in patients with DLBCL and IHC expression of *MYC* and *BCL2* without chromosomal rearrangements (DEL), for whom standard R-CHOP remains preferred.

R-mini-CHOP may be substituted for very frail patients and for patients >80 years of age with comorbidities to improve tolerability.^{32,33} In patients with bulky disease or impaired renal function, initial therapy should include monitoring and prophylaxis for TLS.

Multiple randomized trials (RICOVER 60, NHL-B2, MInT, and the MegaCHOEP trials) have demonstrated superior outcomes in women relative to men, particularly in older adults with older women benefiting more from the addition of rituximab than men.⁴⁹ This could be explained by a slower clearance rate of rituximab in older women. A prospective non-randomized trial evaluated R-CHOP with rituximab dose of 500 mg/m² in men over the age of 60 with DLBCL.⁵⁰ Rituximab 500 mg/m² was associated with better serum levels and improved OS rates in male patients compared to historical data in older men treated with rituximab 375mg/m². In a planned subgroup analysis, rituximab 500 mg/m² was associated with improved PFS (*P* = .039) with a trend toward better OS (*P* = .076) but was also not more toxic than 375 mg/m² rituximab in elderly male patients. Based on these data, a rituximab dose of 500 mg/m² may



be considered in men >60 years of age treated with R-CHOP. A randomized clinical trial is ongoing.

Prediction and Management of CNS Disease

Age >60 years, elevated LDH, ≥ 2 extranodal sites and involvement of specific extranodal sites (bone marrow, testes, kidney, adrenal gland and paranasal) have been associated with increased risk for developing CNS relapse.⁵¹ Stage IE primary DLBCL of the breast has also been identified as a potential risk factor for CNS relapse.^{52,53}

The German High-Grade Non-Hodgkin Lymphoma Study Group (DSHNHL) has proposed a prognostic model (CNS-IPI) to predict the risk of CNS relapse incorporating the 6 clinical factors (age > 60 years, LDH > normal, ECOG PS >1, stage III or IV disease, extranodal involvement >1 and involvement of the kidney and/or adrenal gland).⁵⁴

The CNS-IPI separated patients into three risk groups based on the rate of developing CNS disease at 2 years: low-risk (0 or 1 risk factor; rate of CNS disease 0.6%), intermediate-risk (2 or 3 risk factors; rate of CNS disease 3%) and high-risk group (4 to 6 risk factors; rate of CNS disease at 10%).⁵⁴ In the multivariate analysis, age >60 years, elevated LDH, ECOG PS >1, and stage III or IV disease were identified as the most significant predictors of CNS relapse. Although involvement of >1 extranodal site was not a significant predictor of CNS relapse, it was retained in the final CNS-IPI for the ease of application. Among the specific extranodal sites, only the involvement of the kidney or adrenal gland were significantly associated with CNS relapse or progression. In another international multicenter retrospectively analysis of 1532 patients with DLBCL treated with chemoimmunotherapy, there was a strong correlation between absolute number of extranodal sites and risk of CNS relapse. The 3-year cumulative incidence of CNS relapse was 15% for

patients with >2 extranodal sites compared with 3% for those with ≤ 2 extranodal sites ($P < 0.001$).⁵⁵

There may be an increased risk of CNS relapse in patients with 4 to 6 risk factors according to CNS-IPI; HIV-associated lymphoma; testicular lymphoma; HGBL with translocations of *MYC* and *BCL2* and/or *BCL6* (DHL/THL); HGBL, NOS; or kidney or adrenal involvement. Lumbar puncture should be considered for these patients, particularly if neurologic symptoms are present. The diagnostic yield is improved if flow cytometric analysis of cerebrospinal fluid is undertaken.

CNS prophylaxis should be considered for patients with risk factors for CNS disease but the optimal method is controversial. Intrathecal methotrexate given at least once per systemic treatment cycle has been used for many years. More recent retrospective studies have suggested that high-dose IV methotrexate-based prophylaxis may be associated with a lower incidence of CNS relapses.⁵⁶⁻⁵⁹ Systemic methotrexate with leucovorin rescue has been safely incorporated into R-CHOP-21, with methotrexate administered on day 15 of the 21-day R-CHOP cycle.⁵⁶ However, other reports suggest that CNS prophylaxis is insufficient to prevent CNS relapse.^{60,61} The NCCN Guidelines currently recommend CNS prophylaxis with 4 to 8 doses of intrathecal methotrexate and/or cytarabine, or systemic methotrexate (3–3.5 g/m²). In selected patients, systemic methotrexate (3–3.5 g/m²) may also be given as consolidation after R-CHOP + intrathecal methotrexate/cytarabine.

For concurrent presentation with systemic and CNS DLBCL, systemic methotrexate (≥ 3.5 g/m²) should be incorporated as part of the treatment plan on day 15 of a 21-day R-CHOP cycle that has been supported by growth factors) for parenchymal involvement. Intrathecal methotrexate/cytarabine and/or systemic methotrexate (3–3.5 g/m²) should be incorporated as part of the treatment plan for leptomeningeal involvement. Ommaya reservoir placement should be considered in



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

patients with leptomeningeal disease. When administering high dose methotrexate, patients must be pretreated with hydration and alkalinization of the urine, and then receive leucovorin rescue beginning 24 hours after the initiation of methotrexate infusion. Renal and hepatic function as well as methotrexate clearance must be monitored. Adequate recovery of blood counts should be confirmed prior to initiating the next cycle of R-CHOP.

Role of HDT/ASCR

Several studies have evaluated high-dose therapy and autologous stem cell rescue (HDT/ASCR) as consolidation therapy for patients in first CR after induction therapy.⁶²⁻⁶⁵

In the randomized GELA LNH87-2 study (performed in pre-rituximab era), patients with DLBCL in first CR after induction therapy received consolidation therapy with either sequential chemotherapy or HDT/ASCR.⁶² Although no difference in outcome was prospectively observed in this trial, a retrospective subset analysis of 236 patients with age-adjusted IPI of 2 (high-intermediate) or 3 (high-risk), found that HDT/ASCR resulted in significantly improved outcomes compared with sequential chemotherapy with regards to both 8-year DFS rate (55% vs. 39%; $P = .02$) and 8-year OS rate (64% vs. 49%; $P = .04$) in the high-intermediate/high-risk subset.⁶²

In the French GOELAMS 075 study, patients aged ≤ 60 years with DLBCL (N=286 evaluable) were randomized to receive 8 cycles of R-CHOP-14 or HDT with rituximab (R-HDT) followed by ASCR.⁶³ The 3-year PFS rate and OS rates were 76% and 83%, respectively, with no significant differences between treatment arms.⁶³

In the SWOG 9704 trial, 253 patients with high-intermediate/high IPI were randomized to receive 3 cycles of R-CHOP or HDT/ASCR, following initial remission with 5 cycles of CHOP or R-CHOP induction.⁶⁴ The 2-year PFS

rate was significantly higher with HDT/ASCR compared with chemoimmunotherapy alone (69% vs. 55%; $P = .005$); the 2-year OS rates were not significantly different (74% vs. 71%, respectively; $P = .30$). In an exploratory subset analysis, HDT/ASCR was associated with an OS benefit for high-risk patients. In this subgroup, the 2-year OS rates were 82% and 63% respectively, for patients treated with HDT/ASCR and chemoimmunotherapy. Notably, in this study a third of the patients did not receive rituximab as part of their induction regimen.

In the randomized DLCL04 trial of the Italian Lymphoma Foundation, 399 patients with DLBCL (aged ≤ 65 years) were randomized to receive rituximab-containing first-line regimens (8 cycles of R-CHOP-14 or 6 cycles of R-MegaCHOP-14) with or without HDT/ASCR.⁶⁵ After a median follow-up of 72 months, the 2-year failure-free survival rate was significantly higher in the HDT/ASCR group compared with the non-HDT/ASCR groups (71% vs. 62%; $P = .012$), but the 5-year OS rate was not different between the two groups (78% and 77% respectively; $P = .91$). In addition, grade ≥ 3 hematologic toxicities were higher in the transplant group (92% compared to 68%).

The above studies, overall, found no benefit to upfront HDT/ASCR as compared with first-line rituximab-based chemoimmunotherapy, except in high-risk IPI patients, but this remains controversial since this finding emerged only on a retrospective subset analysis involving a small number of patients. HDT/ASCR is therefore not routinely recommended.

Response Assessment

Interim restaging is performed to identify patients whose disease has not responded to or has progressed on induction therapy. A negative PET scan after 2 to 4 cycles of induction therapy has been associated with significantly higher EFS and OS rates in several studies.⁶⁶⁻⁶⁹ However, interim PET scans can produce false positive results and many patients



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

treated with chemoimmunotherapy have a favorable long-term outcome despite a positive interim PET scan.^{70,71} In one prospective study, the PFS in patients who were interim PET-positive, biopsy-negative (after 4 cycles of accelerated R-CHOP) was identical to that in patients with a negative interim PET scan.⁷⁰ A retrospective analysis also reported only a minor difference in the 2-year PFS rates between patients with a positive interim PET scan and a negative interim PET scan after treatment with 6 to 8 cycles of R-CHOP (72% and 85% respectively; $P = .0475$).⁷¹ Conversely, the end-of-treatment PET scan was highly predictive of PFS; the 2-year PFS rate was 64% for patients with a final positive PET scan compared to 83% for those with a final negative PET scan ($P < .001$).

More recent reports have also confirmed the limited prognostic value of interim PET scans in patients with DLBCL treated with R-CHOP.⁷²⁻⁷⁵ In a prospective study that evaluated the predictive value of interim PET scans after 2 cycles of R-CHOP in 138 evaluable patients, the 2-year EFS rate was significantly shorter for patients with a positive interim PET-scan compared to those with a negative interim PET scan (48% vs. 74%; $P = .004$); however, the 2-year OS was not significantly different between the two groups (88% vs. 91%; $P = .46$).⁷⁴ The results of the PETAL trial showed that a positive interim PET scan (change in SUVmax of $<66\%$) was associated with significantly inferior EFS and OS, although PET-based treatment intensification did not improve outcome in patients with DLBCL treated with R-CHOP.⁷⁶

Therefore, interim PET imaging is not recommended to be used to guide changes in therapy. If treatment modifications are considered based on interim PET scan results, a repeat biopsy of residual masses should be strongly considered to confirm PET-positivity prior to additional therapy. If the biopsy is negative, the planned course of treatment as recommended for PET-negative guidelines should be completed. Patients should undergo evaluation prior to receiving RT, including all positive studies. If

RT is not planned, interim restaging after 2 to 4 cycles of R-CHOP is appropriate to confirm response. End of treatment restaging is performed upon completion of treatment. The optimal time to end of treatment restaging is not known. However, the panel recommends waiting for 6 to 8 weeks after completion of therapy before repeating PET scans.

Response assessment by PET/CT should be done according to the 5-point scale (5-PS).^{19,77} The 5-PS is based on the visual assessment of fluorodeoxyglucose (FDG) uptake in the involved sites relative to that of the mediastinum and the liver.⁷⁸⁻⁸⁰ A score of 1 denotes no abnormal FDG-avidity, while a score of 2 represents uptake less than the mediastinum. A score of 3 denotes uptake greater than the mediastinum but less than the liver, while scores of 4 and 5 denote uptake greater than the liver, and greater than the liver with new sites of disease, respectively. Different clinical trials have considered scores of either 1 to 2 or 1 to 3 to be PET-negative, but a score of 1 to 3 is now widely considered to be PET negative. Scores of 4 to 5 are universally considered PET-positive. A score of 4 on an interim or end of treatment restaging scan may be consistent with a partial response if the FDG-avidity has declined from initial staging, while a score of 5 denotes progressive disease.

Interim and End-of-Treatment Response Evaluation for Stage I-II

When the treatment plan involves ISRT, restaging should be done after completion of first-line chemoimmunotherapy prior to initiation of ISRT as the dose of RT will be influenced by the result (see *Principles of Radiation Therapy* in the algorithm).

If interim restaging demonstrates CR (PET-negative), the planned course of treatment with the same dose of RT is completed. If the interim restaging demonstrates a PR (PET-positive), treatment with a higher dose of RT is appropriate. It is appropriate to enroll patients with an interim PR on a clinical trial. At the present time, there are no data to suggest that a PR with persistent PET positivity after 3 cycles should prompt a change in



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

treatment. If the PET scan is positive after 6 cycles of R-CHOP or R-CHOP-14, the patient can proceed to HDT/ASCR with or without RT. Patients with primary refractory or progressive disease are managed as refractory or relapsed disease.

After end of treatment restaging, follow-up at regular intervals (every 3–6 months for 5 years and then annually or as clinically indicated thereafter) is recommended for patients with CR. In these patients, follow-up CT scans are recommended only if clinically indicated. Patients with PR and those with no response to treatment or progressive disease are treated as described for relapsed or refractory disease. Palliative RT is recommended for selected patients who are not candidates for chemoimmunotherapy.

Interim and End-of-Treatment Response Evaluation for Stage III–IV

If interim staging (after 2–4 cycles) demonstrates a CR or PR, the planned course of R-CHOP to a total of 6 cycles is completed. End of treatment restaging is performed upon completion of treatment.

The role of ISRT following CR (evaluated by CT criteria) to initial bulky sites ≥ 7.5 cm or extranodal involvement was evaluated in the RICOVER-noRT trial (an amendment to the RICOVER-60 trial).²⁷ In this study, 164 patients with stage III–IV disease were treated with 6 cycles of R-CHOP-14 and RT to bulky sites or extranodal involvement was omitted. The 3-year PFS and OS rates were significantly inferior, compared to the corresponding survival rates in patients from the RICOVER-60 trial treated with the same chemoimmunotherapy with RT to bulky sites.²⁷ The study was therefore discontinued. Similarly, subgroup analyses of the MInT and RICOVER-60 trial showed that patients with skeletal involvement significantly benefitted from RT to sites of skeletal involvement.⁸¹ Although retrospective subgroup analyses may be subjected selection biases, the benefit of RT held up on multivariate analysis in both studies.

After end of treatment restaging, observation is preferred for patients with CR. ISRT to initially bulky disease or isolated skeletal sites can be considered for patients with CR.^{27,81} Lenalidomide maintenance as first-line consolidation therapy for patients 60 to 80 years of age is included with a category 2B recommendation based on the results of the phase III randomized ReMARC trial which showed that lenalidomide maintenance for 24 months significantly prolonged PFS in elderly patients with DLBCL achieving CR or PR to R-CHOP induction therapy, but without impact on OS.⁸²

Patients in CR are followed up at regular intervals (every 3–6 months for 5 years and then annually or as clinically indicated thereafter). In these patients, follow-up imaging CT scans should be performed no more than every 6 months for 2 years after completion of therapy, and then only as clinically indicated thereafter.

Patients with PR (after completion of initial therapy) and those with no response to treatment or progressive disease are treated as described below for relapsed or refractory disease. Palliative RT is recommended for selected patients who are not candidates for chemoimmunotherapy.

Follow-up

Considerable debate remains regarding the routine use of imaging for surveillance in patients who achieve a CR after induction therapy. Although positive scans can help to identify patients with early asymptomatic disease relapse, false positive cases remain common and problematic, and may lead to unnecessary radiation exposure and invasive procedures for patients, as well as increased healthcare costs. In a study that evaluated the use of surveillance CT scans (at 3 and 12 months after completion of chemotherapy) in 117 patients with DLBCL who achieved a CR with induction chemotherapy, 35 patients relapsed, and only 6% of these relapses were detected by follow-up CT scan in



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

asymptomatic patients; 86% of cases of relapse were associated with development of new symptoms or signs of relapse.⁸³ The investigators therefore concluded that routine surveillance with CT scans had limited value in the detection of early relapse in patients with a CR following induction therapy. In a retrospective study evaluating the use of surveillance imaging in 108 patients with relapsed aggressive lymphoma who had a CR to initial chemotherapy, 20% of relapses were detected by imaging in asymptomatic patients.⁸⁴ In the remaining 80% of cases, relapse was identified by clinical signs and/or symptoms. Moreover, the cases of relapse detected by imaging were more likely to represent a population of patients with low-risk disease based on age-adjusted IPI at the time of relapse.⁸⁴ Thus, routine imaging during remission may help to identify patients with more limited disease at the time of relapse, but has not been shown to improve ultimate outcome.

In a prospective study that evaluated the role of PET scans (at 6, 12, 18, and 24 months after completion of induction therapy) in patients with a CR after induction therapy for lymphomas, surveillance using PET scans was found to be useful for detecting early relapse.⁸⁵ Among the cohort of patients with aggressive lymphomas in this study (n=183), follow-up PET scans detected true relapses in 10% of patients at 6 months, 5% at 12 months, and 11% at 18 months; the rate of false-positive scans was low, at 1% (including cohorts of patients with indolent and aggressive NHL).⁸⁵ Inconclusive PET scans were obtained in 4% of patients (8 out of 183), 6 of those had confirmed relapse based on biopsy evaluation. In a retrospective study that evaluated the use of follow-up PET/CT scan in 75 patients with DLBCL who achieved a CR after induction therapy, follow-up PET/CT scan detected relapse in 27 patients, of which 23 patients had confirmed relapse based on biopsy evaluation; thus, the positive predictive value of PET/CT scan for detecting relapse was 0.85.⁸⁶ In this study, patient age (>60 years) and the presence of clinical signs of relapse were significant predictors of disease relapse.

Data from more recent retrospective studies also suggest that routine surveillance with PET or CT scans is of limited utility in the detection of relapse in the majority of patients with DLBCL.⁸⁷⁻⁸⁹ A study comparing the performance of surveillance PET scans in patients with DLBCL treated with CHOP alone versus R-CHOP, found higher false positive results in patients treated with R-CHOP (77% vs. 26%; $P < .001$).⁸⁷ Another study reported a positive predictive value of 56% for surveillance PET/CT scans in patients with an IPI score <3 compared with 80% for patients with an IPI score ≥ 3 , suggesting that surveillance PET/CT has a very limited role in the majority of patients in CR after primary therapy.⁸⁸ Another multiinstitutional retrospective study evaluated the utility of surveillance scans in two independent prospectively enrolled cohorts of patients with DLBCL treated with anthracycline-based chemoimmunotherapy.⁸⁹ In one cohort (n = 680; 552 patients entered post-treatment observation), post treatment surveillance scans detected DLBCL relapse prior to clinical manifestations only in 2% of patients during a planned follow-up visit. In another cohort (n = 261; 222 patients entered post-treatment observation), surveillance imaging detected asymptomatic relapse only in 2% of patients. A population-based study of patients from the Danish and Swedish lymphoma registries also showed that imaging-based surveillance strategy had no impact on survival for patients DLBCL in first complete remission.⁹⁰

A multiinstitutional retrospective study evaluated the EFS at 24 months (EFS24) in two independent prospectively enrolled cohorts of 767 patients with DLBCL treated with anthracycline-based chemoimmunotherapy.⁹¹ Patients who achieved EFS24 had an OS equivalent to that of the age-matched and sex-matched general population ($P = .25$). This was also confirmed in another data set that included 820 patients from a GELA LNH2003B program and the hospital-based registry in France ($P = .71$). These data indicate that EFS24 should be useful for developing strategies



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

for post-therapy surveillance, patient counseling, and as an end point in clinical studies for patients with DLBCL.

In the absence of evidence demonstrating an improved outcome favoring routine surveillance imaging for the detection of relapse, the NCCN Guidelines do not recommend the use of PET or CT for routine surveillance for patients with stage I-II disease who have achieved a CR to initial therapy. For patients with stage III-IV disease who achieve remission to initial therapy, the NCCN Guidelines recommend CT scans no more than once every 6 months for up to 2 years after completion of treatment, with no ongoing routine surveillance imaging after that time, unless it is clinically indicated. When surveillance imaging is performed, CT scan is preferred over PET/CT for the majority of patients. PET/CT may be preferable for patients with primarily osseous presentations, with the caveat that bone remodeling may also be FDG-avid, so a biopsy is recommended for PET positive sites prior to instituting second line therapy.

Relapsed or Refractory Disease

The role of HDT/ASCR in patients with relapsed or refractory disease was demonstrated in an international randomized phase III trial (Parma study) in the pre-rituximab era.⁹² In this study, patients with DLBCL responding to induction DHAP (dexamethasone, cisplatin and cytarabine) chemotherapy after first or second relapse (n=109) were randomized to receive additional DHAP chemotherapy plus RT or RT plus HDT/ASCR. The 5-year EFS rate was significantly higher among the transplant group compared with the non-transplant group (46% vs. 12%; $P=.001$), as was the 5-year OS (53% vs. 32%; $P=.038$).⁹² A recent retrospective analysis based on data from the EBMT registry evaluated the role of HDT/ASCR in patients achieving a second CR after second-line therapy (N=470).⁹³ In this analysis, 25% of patients had received rituximab-based therapy prior to ASCR. The 5-year DFS and OS was 48% and 63% after ASCR for all patients. The median

DFS after ASCR was 51 months, which was significantly longer than the duration of first CR (11 months; $P<.001$). The longer DFS with ASCR compared with first CR was also significant in the subgroup of patients previously treated with rituximab (median not reached vs. 10 months; $P<.001$) and the subgroup who relapsed within 1 year of first-line therapy (median 47 months vs. 6 months; $P<.001$).⁹³

Outcomes of relapsed/refractory DLBCL differ based on the response to initial therapy, timing of relapse and opportunity to undergo HDT/ASCR. In a population based study of 1039 patients treated with anthracycline-based chemoimmunotherapy, 244 patients (23%) had relapsed disease that was subsequently re-treated.⁹⁴ Across all therapies, the 4-year OS rate was 28% and 51% for patients who underwent HDT/ASCR. The 4-year OS rate was 47% for patients with disease relapse after 12 months from initial diagnosis whereas the 4-year OS rate was 13% for those with a transient or no response to initial therapy. Pre-transplantation PET scans have been identified as predictive factors in patients undergoing HDT/ASCR, with positive PET scans following second-line therapy predicting poor outcomes following HDT/ASCR.⁹⁵⁻⁹⁷ The results of studies from the GELTAMO group and ABMTR suggested that HDT/ASCR should be considered for patients who do not achieve a CR but who are still sensitive to chemotherapy.⁹⁸⁻¹⁰⁰

Effective second line regimens for patients who are candidates for HDT/ASCR include R-ICE (rituximab, ifosfamide, carboplatin and etoposide),¹⁰¹⁻¹⁰³ R-DHAP (rituximab, dexamethasone, cytarabine and cisplatin),^{103,104} R-DHAX (rituximab, dexamethasone, cytarabine and oxaliplatin)^{105,106} or R-GDP (gemcitabine, dexamethasone, cisplatin or carboplatin).^{101-103,107,108} The efficacy of second-line therapy is predicted by the second-line age-adjusted IPI.^{109,110}

In the international randomized intergroup study (CORAL study) that evaluated second-line chemoimmunotherapy (R-ICE vs. R-DHAP)



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

followed by ASCR in 469 patients with chemosensitive relapsed or refractory DLBCL no significant difference in outcome was found between treatment arms.¹⁰³ The overall response rates (ORRs) were 63% after R-ICE and 64% after R-DHAP. After a median follow-up of 44 months, the 4-year EFS rate was 26% with R-ICE compared with 34% with R-DHAP ($P = .2$) and the 4-year OS rate was 43% and 51%, respectively ($P = .3$).¹⁰³ Notably, patients relapsing less than 1 year after initial R-CHOP therapy had a particularly poor outcome with 3-year PFS of 23%. A subgroup analysis from the CORAL study (Bio-CORAL) showed that for patients with a GCB phenotype (based on Hans algorithm), R-DHAP resulted in improved PFS (3-year PFS 52% vs. 31% with R-ICE).¹¹¹ This difference was not observed among patients with non-GCB phenotype (3-year PFS 32% with R-DHAP vs. 27% with R-ICE).¹¹¹ Moreover, the subgroup of patients with *MYC* gene rearrangement (with or without concurrent in *BCL2* and/or *BCL6* gene rearrangements) had poor outcomes regardless of treatment arm.¹¹² The 4-year PFS was 18% among patients with *MYC* gene rearrangements compared with 42% in those without ($P = .032$); 4-year OS was 29% and 62%, respectively ($P = .011$). Among patients with *MYC* gene rearrangements, the 4-year PFS was 17% with R-DHAP and 19% with R-ICE; OS was 26% and 31%, respectively.¹¹²

The CORAL study was also designed to evaluate the role of rituximab maintenance (every 2 months for 1 year) following ASCR. Among the 242 patients randomized after ASCR to rituximab maintenance or observation, the 4-year EFS rate was similar between the two groups: 52% with rituximab versus 53% with observation.¹⁰³ The proportion of patients with progression or relapse was also similar between the two groups. In addition, the 4-year OS rate was also not statistically different (61% and 65%, respectively). Serious adverse events were more frequent in the rituximab maintenance arm. Given that this study showed no benefit with rituximab maintenance compared with observation following ASCR,

maintenance therapy cannot be recommended in this setting.^{103 103 103 103}

In a randomized study of 619 patients with relapsed/refractory aggressive lymphoma (419 patients with relapsed/refractory DLBCL randomized to receive GDP [gemcitabine, dexamethasone, cisplatin] or DHAP), GDP was non-inferior to DHAP in terms of ORR (45% vs. 44%) and transplantation rate (52% vs. 49%).¹⁰⁷ After a median follow-up of 53 months, no differences were detected in EFS ($P = .95$) or OS ($P = .78$) between GDP and DHAP. Treatment with GDP was associated with less toxicity ($P < .001$).

Second-line combination chemotherapy is recommended for patients with an intention to proceed to transplant. Rituximab should be included if there is disease relapse after reasonable remission period (>6 months) and biopsy demonstrates continued CD20 expression; however, rituximab should often be omitted in patients with primary refractory disease. Consolidation therapy with HDT/ASCR (category 1 for patients with CR) with or without RT is recommended for patients with CR or PR to second-line therapy, if they are candidates for transplant.^{92,93} ISRT before HDT/ASCR has been shown to result in good local disease control and improved outcome.¹¹³ Additional RT can be given to limited sites with prior positive disease before or after HDT/ASCR. Allogeneic HCT should be considered in selected patients with mobilization failures and persistent bone marrow involvement or lack of adequate response to second line therapy, though patients should be in CR or near CR at the time of transplant.

Patients who are not candidates for transplant should ideally be treated in the context of a clinical trial. Standard options include gemcitabine-based regimens with or without rituximab,¹¹⁴⁻¹¹⁶ bendamustine ± rituximab,¹¹⁷⁻¹¹⁹ or brentuximab vedotin if CD30 positive.¹²⁰ Lenalidomide with or without rituximab,^{9,10} and ibrutinib¹¹ are appropriate options at relapse, particularly



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

for patients with non-GCB DLBCL. Based on the results of the phase II randomized clinical trial that evaluated bendamustine + rituximab (BR) versus polatuzumab vedotin + BR, bendamustine ± rituximab is included with a category 2B recommendation.

Patients who are not candidates for transplant with PR to second-line therapy or those with progressive or refractory disease to second-line therapy (regardless of transplant eligibility) or those with disease relapse following HDT/ASCR or allogeneic HCT should be managed with third-line systemic therapy, palliative ISRT or best supportive care. Repeat biopsy should be strongly considered prior to additional therapy for patients who are non-candidates for transplant achieving PET-positive PR to second-line therapy because PET positivity may represent post-treatment inflammation.

However, patients with progressive disease after ≥2 prior lines of systemic therapy regimens are unlikely to derive additional benefit from currently available systemic therapy options, except for patients who have experienced a long disease-free interval. The results of a multicenter retrospective study (SCHOLAR-1) showed that outcomes are consistently poor among patients with refractory DLBCL.¹²¹ In this study that evaluated the outcome of 636 patients with refractory DLBCL (defined as progressive or stable disease as best response at any point during chemotherapy [>4 cycles of first-line or 2 cycles of later-line therapy] or disease relapse at ≤ 12 months from HDT/ASCR), the pooled ORR was 26% (7% CR) to the next line of therapy, and the median OS was 6 months.¹²¹ The response rates were much lower for patients with primary refractory disease or refractory disease after second-line or subsequent therapy (20% and 26%, respectively) compared to those with disease relapse ≤ 12 months from HDT/ASCR. Data from recent clinical trials (discussed below) suggest that chimeric antigen receptor (CAR) T-cell therapy represents an improved treatment option for this group of patients.

Recently, FDA granted accelerated approval of polatuzumab vedotin (anti-CD79b antibody-drug conjugate) in combination with bendamustine and rituximab (BR) for the treatment relapsed or refractory DLBCL after ≥ 2 prior lines of therapies. This approval is based on the results from the phase II randomized trial that evaluated polatuzumab vedotin + BR (n=40) versus BR (n=40) in patients with relapsed or refractory DLBCL ineligible for HDT/ASCR.¹²² At a median follow-up of 22 months, the ORR were 45% and 18%, for polatuzumab vedotin + BR and BR, respectively. The CR rates were significantly higher for polatuzumab vedotin + BR compared with BR alone (40% vs 18%; $P = .026$). The median PFS (8 months vs. 2 months; $P < .0001$) and OS (12 months vs. 5 months; $P = .0023$) were also significantly longer for polatuzumab vedotin + BR compared with BR. Polatuzumab vedotin + BR was also associated with survival benefit regardless of COO (GCB vs. non-GCB) and double expressor status (MYC and BCL2 overexpression), although the biomarker sample sizes were small. In patients with non-GCB subtype (14 patients in the polatuzumab vedotin + BR arm; 16 patients in the BR arm), the median PFS and OS were 11 months and 14 months, respectively for polatuzumab vedotin + BR. The corresponding median PFS and OS were 3 months and 4 months, respectively for BR. In patients with MYC and BCL2 overexpression (9 patients in the polatuzumab vedotin + BR arm; 6 patients in the BR arm), the median PFS and OS were 7 months and 13 months, respectively for polatuzumab vedotin + BR. The corresponding median PFS and OS were <1 month and 4 months for BR. Polatuzumab vedotin + BR is an appropriate treatment option for patients with relapsed or refractory DLBCL (after ≥ 2 prior lines of therapies) ineligible for HDT/ASCR.

CAR T-Cell Therapy

Axicabtagene ciloleucel and tisagenlecleucel are anti-CD19 CAR T-cell therapies that are FDA-approved for the treatment of adult patients with relapsed/refractory DLBCL, HGBL and transformed follicular lymphoma



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

(TFL) after ≥ 2 prior systemic therapy regimens, based on the results of the ZUMA-1 and JULIET trials (discussed below).¹²³⁻¹²⁵ Axicabtagene ciloleucel is also approved for relapsed or refractory PMBL after ≥ 2 prior systemic therapy regimens.

The multicenter phase II study (ZUMA-1) evaluated axicabtagene ciloleucel in patients with refractory DLBCL (n=81), TFL (n = 30) or PMBL (n = 8).¹²³ Eligible patients had an absolute neutrophil count $>1,000$ cells/ μ L, an absolute lymphocyte count greater than >100 cells/ μ L, a platelet count $>75,000$ cells/ μ L, adequate organ function, no central CNS involvement, and no active infection. In the primary analysis that included 101 patients (78 patients with disease refractory ≥ 2 lines of prior therapy and 21 patients with disease relapse after HDT/ASCR) that were evaluated 6 months after the infusion of axicabtagene ciloleucel, the ORR was 82% (54% CR and 28% PR). With a median follow-up of 15 months, the estimated PFS rate at 15 months and the OS rate at 18 months were 41% and 52%, respectively. The ORRs were 81% (49% CR; 32% PR) and 83% (71% CR; 12% PR), respectively for the subgroup of patients with DLBCL and TFL or PMBL. The 2-year follow-up data also confirmed these initial findings.¹²⁴ After a median follow-up of 27 months, 39% of patients remained in ongoing remission, and 51% remained alive. The median duration of response was 11 months and the median PFS was 6 months.

The multicenter phase II study (JULIET) evaluated tisagenlecleucel in patients with relapsed/refractory DLBCL (after at least two lines of therapy, including rituximab and an anthracycline) and TFL.¹²⁵ Patients who had been previously treated with anti CD19 CAR T-cell therapy or an allogeneic HCT and those with PMBL, or active CNS involvement were excluded. In this study, 115 patients received infusion with tisagenlecleucel. Among the 93 patients included in the efficacy analysis with ≥ 3 months of follow-up or had discontinued participation in the study before 3 months, the best ORR was 52% (40% CR; 12% PR). The

estimated PFS rate at 12 months was 83% for patients who had a CR or PR at 3 months. The estimated OS rate at 12 months was 49% for all patients and 90% for those with a CR.

Axicabtagene ciloleucel and tisagenlecleucel result in responses across all subgroups, including DHL/THL, GCB and non-GCB subtypes, and patients with either relapsed or primary refractory disease. CD19 expression levels did not correlate with response. The NCCN Guidelines recommend CAR T-cell therapy (axicabtagene ciloleucel or tisagenlecleucel) for patients achieving PR following second-line therapy (regardless of their eligibility for transplant) and for those with disease relapse after achieving CR to second-line therapy or progressive disease., Bendamustine should be used with caution (unless immediately prior to CAR T-cell therapy) in patients intended to receive CAR T-cell therapy, since it could impact the success of the patient's T-cell collection.

Guidance for the Treatment of Patients with CAR T-Cell Therapy

Axicabtagene ciloleucel and tisagenlecleucel should only be dispensed and administered in health care facilities that are enrolled in and comply with the Risk Evaluation and Mitigation Strategies (REMS) requirements.

Cytokine-release syndrome (CRS) and neurologic toxicity (also known as CAR-T-cell-related encephalopathy syndrome [CRES]) are the most frequent life-threatening toxicities associated with CAR T-cell therapy. In the ZUMA-1 trial, CRS and neurologic toxicity of any grade were reported in 93% and 64% of patients respectively.^{123,124} The median time from infusion to onset of symptoms was 2 days and 5 days respectively, with the median duration of 8 days and 17 days, respectively. Grade ≥ 3 CRS and neurologic events occurred in 13% and 28% of the patients, respectively. Pyrexia (11%), hypoxia (9%), and hypotension (9%) were the most common symptoms of grade ≥ 3 CRS. The median time to the onset of CRS was 2 days after infusion and the median time until resolution was 8 days. Encephalopathy (21%), confusional state (9%), aphasia (7%), and



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

somnolence (7%) were the most common grade ≥ 3 neurologic events. The incidence of grade ≥ 3 CRS and neurologic events decreased over the course of the study (occurring in 12% and 31% of patients respectively after a median follow-up of 15 months). In the JULIET trial, CRS and neurologic toxicity of any grade were reported in 58% and 21% of patients respectively.¹²⁵ Grade ≥ 3 CRS and neurologic events occurred in 22% and 12% of patients respectively. The median time from infusion to onset of symptoms was 2 days and 6 days respectively, with the median duration of 7 days and 14 days, respectively. Notably, different toxicity scales were used for the grading of CRS in the ZUMA-1 and JULIET trials (grade 3 CRS in JULIET was similar to grade 2 in ZUMA-1), so while rates of any CRS can be compared, the rates specifically of severe (grade 3 or 4) CRS cannot be directly compared with accuracy.

The anti-interleukin-6 (IL-6) receptor monoclonal antibody tocilizumab is highly effective for the management of CRS, inducing rapid reversal of symptoms in most patients.¹²⁶ Tocilizumab is approved for the treatment of CRS occurring after CAR-T-cell therapy and its use has not been shown to have any impact on the efficacy of CAR-T-cell therapy, in terms of response rates or duration of response. Corticosteroids are also an important adjunctive treatment for CRS, in concert with tocilizumab. Corticosteroids are preferred for the management of neurologic toxicity if not associated with CRS whereas tocilizumab in combination with corticosteroids is recommended for management of CRS occurring in tandem with neurologic toxicity.¹²⁶ The use of corticosteroids for the management of toxicities associated with CAR T-cell therapy did not impact the efficacy of CAR T-cell therapy, although their use has been to shown to impair T-cell function. CRS and neurologic toxicity should be managed based on the toxicity grade as outlined in the *Management of CAR T-Cell-Related Toxicities* section of the NCCN Guidelines for the Management of Immunotherapy-Related Toxicities.

Primary Mediastinal Large B-Cell Lymphoma

PMBL is a distinct subtype of NHL that can be histologically indistinguishable from DLBCL that tends to occur in young adults with a median age of 35 years with a slight female predominance.^{127,128} PMBL arises from thymic B-cells with initial locoregional spread to supraclavicular, cervical, hilar nodes and into the mediastinum and lung.¹²⁷ Widespread extranodal disease is uncommon at initial diagnosis, present in approximately one quarter of patients, but can be more common at recurrence.¹²⁸ Clinical symptoms may be related to rapid growth of a mediastinal mass include superior vena cava (SVC) syndrome, pericardial effusions and pleural effusions.

GEP has indicated that PMBL is distinct from DLBCL; the pattern of gene expression in PMBL is more similar to classical Hodgkin lymphoma (CHL).^{129,130} PMBL expresses B-cell antigens and lacks surface immunoglobulin. PMBL is CD19+, CD20+, CD22+, CD21-, IRF4/MUM1+ and CD23+ with a variable expression of BCL2 and BCL6. CD30 is weakly and heterogeneously expressed in more than 80% of cases.¹²⁸ PMBL is also characterized by a low expression of HLA I or II molecules. There have been rare cases of mediastinal gray zone lymphomas with combined features of PMBL and CHL. Cytogenetic abnormalities that are common in PMBL include gains in chromosome 9p24 (involving the *JAK2* in 50–75% of patients) and chromosome 2p15 (involving the *c-REL*, encoding a member of the NF- κ B family of transcription factors) and loss in chromosomes 1p, 3p, 13q, 15q, and 17p.¹²⁸ Age-adjusted IPI is of limited value in determining the prognosis of PMBL at diagnosis.^{127,131,132} In a retrospective analysis of 141 patients, ≥ 2 extranodal sites and the type of initial therapy were predictors of outcome for EFS, whereas only the initial therapy was a predictor for OS.¹³¹

In retrospective analyses from the pre-rituximab era, intensive chemotherapy regimens have appeared more effective than CHOP and



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

the addition of IFRT has been associated with improved PFS.¹³²⁻¹³⁶ The results of subsequent retrospective studies suggest that the addition of rituximab to CHOP improves outcome in patients with PMBL.¹³⁷⁻¹⁴¹

In a subgroup analysis of 87 patients with PMBL from the MInT study, the addition of rituximab significantly improved the CR rate (80% vs. 54% without rituximab; $P=.015$) and 3-year EFS rate (78% vs. 52%; $P=.012$), but not the OS rate (89% vs. 78%; $P=.158$).¹³⁸ The MInT study, however, only included young low-risk patients with IPI scores 0-1. In a more recent retrospective analysis that evaluated the outcome of 80 patients with PMBL treated with a CHOP-based regimen with and without rituximab, the 5-year PFS (95% vs. 67%; $P=.001$) and OS (92% vs. 72%; $P=.001$) rates were significantly higher in the rituximab arm.¹³⁹ In multivariate analysis, only the addition of rituximab to induction chemotherapy and CR after first line therapy had a beneficial effect on both PFS and OS. In another retrospective analysis of 95 consecutive patients treated with chemotherapy (VACOP-B or CHOP) with and without rituximab, the 5-year PFS and OS rates were 79% and 97%, respectively for patients treated with rituximab-based chemotherapy compared with 58% and 88%, respectively for those treated with chemotherapy alone.¹⁴¹ The 5-year PFS rates in patients treated with R-VACOP-B, R-CHOP, VACOP-B, and CHOP were 83%, 69%, 62%, and 20%, respectively.

A more recent retrospective analysis of 63 patients with PMBL treated with R-CHOP reported a 21% rate of primary induction failure, with adverse predictors of outcome being advanced stage and high-risk IPI scores, suggesting that R-CHOP may not be the optimal chemotherapy backbone in PMBL, particularly for high-risk patients.¹⁴² Sequential dose dense R-CHOP followed by ICE consolidation (without RT) was also highly effective in patients with PMBL, with similar outcomes to the above analysis with R-chemotherapy from the MInT study.¹⁴³ At a median follow

up for surviving patients at 3 years, the OS and PFS rates were 88% and 78%, respectively.¹⁴³

DA-EPOCH-R has also been evaluated in small cohorts of patients with PMBL.¹⁴⁴⁻¹⁴⁷ A prospective phase II study from the NCI showed that DA-EPOCH-R is a highly effective regimen in patients with PMBL and obviates the need for RT in the large majority of patients.¹⁴⁴ In this study DA-EPOCH-R for 6 cycles and filgrastim, without RT, was evaluated in 51 patients with previously untreated PMBL. Stage IV disease was present in 29% of patients. After treatment with DA-EPOCH-R, 2 patients showed persistent focal disease and 1 patient had disease progression; 2 of these patients required mediastinal RT while 1 patient was observed after excision biopsy. At a median follow up of 63 months, EFS and OS rates were 93% and 97%, respectively. Grade 4 neutropenia and thrombocytopenia occurred in 50% and 6% of treatment cycles, respectively. Hospitalization for febrile neutropenia occurred in 13% of cycles. In a multicenter retrospective analysis that assessed the outcome of 156 patients with PMBL treated with DA-EPOCH-R (38 children and 118 adults; all patients received at least one cycle of DA-EPOCH-R. RT was administered in 15% of patients). At a median follow-up of 23 months, the estimated 3-year EFS and OS rates were 86% and 95%, respectively. Negative PET at end-of-treatment was associated with improved EFS (95% vs. 55%, $P<.001$).¹⁴⁶ In another multi-center cohort analysis of 132 patients with PMBL (56 patients were treated with R-CHOP and 76 patients were treated with DA-EPOCH-R), both R-CHOP and DA-EPOCH-R were associated with excellent 2-year OS rates (89% and 91% respectively).¹⁴⁷ While DA-EPOCH-R resulted in higher CR rates than R-CHOP, patients treated with DA-EPOCH-R were more likely to experience short-term treatment-related toxicities but were spared long-term risks associated with mediastinal RT.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

In the absence of randomized trials, optimal first-line treatment for patients with PMBL is more controversial than other subtypes of NHL. However, based on the available data, the following regimens are included as options for first-line therapy: DA-EPOCH-R (6 cycles)¹⁴⁴ + RT only for persistent PET-positive local disease; R-CHOP (6 cycles) +/- RT; or R-CHOP (4 cycles) followed by ICE with or without rituximab (3 cycles) with or without RT (category 2B).¹⁴³

Post-treatment PET/CT is considered essential; Residual mediastinal masses are common. For patients initially treated with R-CHOP, consolidation with RT should be considered, particularly if increased FDG-activity persists in the primary tumor. If PET/CT is negative at the end of treatment after more intensive therapies (e.g., DA-EPOCH-R) and initial disease was non-bulky, observation may be appropriate. If PET/CT is positive, biopsy is recommended before additional treatment is contemplated.

Relapsed or refractory disease should be managed as described above for DLBCL. However, outcomes of second-line therapy followed by HDT/ASCR remains undefined in patients with relapsed or refractory PMBL. Retrospective analyses that evaluated the outcomes of patients with relapsed/refractory PMBL have reported OS rates of 67% to 68% and PFS rates of 57% in patients undergoing HDT/ASCR after second-line therapy.^{148,149} In a multivariate analysis, incomplete response to initial therapy, advanced Ann Arbor stage at disease progression, and failure to achieve a \geq PR after second-line therapy were independently associated with inferior EFS and OS.¹⁴⁹

Pembrolizumab (a humanized anti-PD-1 monoclonal antibody) has demonstrated promising activity resulting in an ORR of 41% after a median follow-up of 11 months in a cohort of 18 patients with relapsed/refractory PMBL enrolled in the KEYNOTE-013 study (a multicenter, international trial evaluating the safety and efficacy of

pembrolizumab in patients with hematologic malignancies).¹⁵⁰ Pembrolizumab is FDA-approved for the treatment of relapsed or refractory PMBL after \geq 2 prior lines of therapy. As discussed above, CAR T-cell therapy with axicabtagene ciloleucel is also approved for relapsed or refractory PMBL after \geq 2 prior systemic therapy regimens.^{123,124}

Grey Zone Lymphomas

Grey zone lymphoma, officially known in the 2017 WHO classification as B-cell lymphoma unclassifiable with features intermediate between DLBCL and CHL,⁴ refers to lymphomas with overlapping pathologic features between CHL and PMBL, and with a poorer clinical outcome than either CHL or PMBL.¹⁵¹⁻¹⁵⁵ In a study that evaluated epigenetic changes based on DNA methylation analysis of microdissected tumor cells from patients with mediastinal grey zone lymphomas, PMBL, CHL, and DLBCL, the methylation profiles of patients with grey zone lymphoma were intermediate to those of PMBL and CHL, but distinct from patients with DLBCL.¹⁵⁶ Among 235 CpG targets that were identified as being differentially methylated between the lymphomas, 22 targets could be used to readily distinguish between PMBL and CHL, with grey zone lymphomas showing an overlap of both signatures. This unique epigenetic signature of mediastinal grey zone lymphomas provide validation of its classification as a separate disease entity in the WHO classification.^{4,156}

The morphology of grey zone lymphomas is characterized by sheet-like growth of pleomorphic cells in a diffusely fibrous stroma; cells are typically larger and more pleomorphic than those in PMBL, and may sometimes resemble lacunar or Hodgkin-like cells; necrosis without neutrophilic infiltration is frequently present.^{4,153} Mediastinal grey zone lymphomas with biologic and morphologic features of both CHL and DLBCL, are more commonly seen in young adult males between the ages of 20 to 40 years and are characterized by the presence of a large anterior mediastinal mass with or without supraclavicular lymph node involvement.^{151,152,154}



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Non-mediastinal grey zone lymphomas occur in older patients, have a higher incidence of bone marrow involvement, include more than one extranodal disease, and have advanced stage disease and higher risk IPI score than mediastinal grey zone lymphomas.¹⁵⁷ In a retrospective multicenter analysis of 112 patients with grey zone lymphomas, mediastinal presentations were found in 43% of patients, while 57% presented with non-mediastinal grey zone lymphomas.¹⁵⁷

The immunophenotype is atypical, often showing transitional features between PMBL and CHL. In general, CD45 is often positive, and CD15, CD20, CD30, and CD79a are also frequently positive. CD10 and ALK are usually negative. B-cell transcription factors such as PAX5, BOB.1, and OCT-2 are often positive; BCL6 is variably expressed and EBV is more often negative.^{151-153,155} If the morphology more closely resembles PMBL, dim or no expression of CD20 and strong expression of CD30 and CD15 would be suggestive of grey zone lymphoma. If the morphology more closely resembles CHL, strong CD20 expression (and/or other B-cell markers) and absence of CD15 would be suggestive of grey zone lymphoma.⁴

The treatment of patients with grey zone lymphomas poses a challenge, as these lymphomas appear to be associated with a worse prognosis compared with PMBL or CHL.^{151,153,158} No standard of care or consensus exists for the management of patients with grey zone lymphomas, although patients are typically treated with multiagent chemotherapy regimens used for patients with DLBCL. In a multicenter retrospective analysis of gray zone lymphoma (that did not have central pathology review), patients treated with CHOP-like regimens with or without rituximab had superior outcomes compared to subjects treated with ABVD, with 2 year PFS rates of 52% and 22%, respectively.¹⁵⁷ DA-EPOCH-R has also been associated with improved clinical outcomes.^{159,160} In a prospective study that evaluated 6 to 8 cycles of DA-EPOCH-R in a small

group of patients with mediastinal grey zone lymphoma (n=24), the EFS and OS were 62% and 74%, respectively, at the median follow-up of 59 months.¹⁵⁹ With a median follow-up of 5 years, the EFS (62% vs 93%; $P = .0005$) and OS (74% vs 97%; $P = .001$), were significantly lower for patients with mediastinal grey zone lymphoma compared to patients with PMBL (n = 51) enrolled in the same study.

Patients with grey zone lymphomas are best managed in cancer centers with experience in treating this type of lymphoma. In the absence of suitable clinical trials, R-CHOP-21 or DA-EPOCH-R should be considered for initial therapy. Given the apparent inferior outcomes among gray zone lymphomas treated with traditional chemotherapy regimens, consolidative RT should be strongly considered for patients with limited stage disease amenable to RT.

High-Grade B-Cell Lymphomas with translocations of *MYC* and *BCL2* and/or *BCL6*

HGBL with translocations of *MYC* and *BCL2* and/or *BCL6* have been observed in 2% to 11% of newly diagnosed patients with DLBCL. The vast majority are GCB-DLBCL and are characterized by highly aggressive clinical behavior and overlapping pathologic features with DLBCL, B-cell lymphoblastic lymphoma/leukemia (B-LBL) and Burkitt lymphoma (BL).¹⁶¹ FISH for *MYC*, *BCL2*, and *BCL6* gene rearrangements is recommended for those with expression of *MYC* and either *BCL2* or *BCL6* by IHC.

IHC will also identify DLBCL with dual expression of both *MYC* and *BCL2* proteins without associated rearrangements (double expressing lymphomas; DEL), which also have an inferior prognosis.^{162,163} In a series of 193 patients with DLBCL uniformly treated with standard R-CHOP, the 3-year PFS rate (46% vs. 65%; $P = .012$) and 3-year OS rate (46% vs. 75%; $P = .002$) were significantly lower in patients with DEL compared with those without protein expression of *MYC* and *BCL2*.¹⁶² In another study with a longer follow-up, 5-year PFS and OS were 18% and 27%,



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

respectively, in patients with DHL/THL and DEL treated with R-CHOP.¹⁶³ MYC or BCL2 expression alone has not been associated with a worse outcome.^{162,163} In a multivariate analysis that included IPI score and cell of origin, concurrent MYC/BCL2 expression remained a significant independent predictor of poorer PFS and OS after R-CHOP.^{162,163}

Data from retrospective studies suggest that more intensive chemoimmunotherapy regimens may result in better outcomes in patients with DHL/THL.¹⁶⁴⁻¹⁶⁶ In a multicenter retrospective analysis of 106 patients with DHL/THL, treatment with intensive regimens such as DA-EPOCH-R, R-HyperCVAD (rituximab, cyclophosphamide, vincristine, doxorubicin, and dexamethasone) or R-CODOX-M/IVAC (rituximab, cyclophosphamide, doxorubicin, vincristine, methotrexate/ifosfamide, etoposide, and high dose cytarabine) resulted in superior CR and PFS compared to R-CHOP.¹⁶⁴ A meta-analysis compared survival outcomes in patients with double-hit cytogenetics treated with more aggressive regimens including R-HyperCVAD, R-CODOX-M/IVAC or R-EPOCH versus R-CHOP in the first-line setting.¹⁶⁷ The median PFS for the R-CHOP, DA-EPOCH-R and other dose intensive regimens was 12, 22, and 19 months, respectively. DA-EPOCH-R significantly reduced the risk of progression compared with R-CHOP; however, OS was not significantly different across treatment approaches.

DA-EPOCH-R was evaluated in a prospective phase II study of 53 patients with untreated aggressive B-cell lymphoma with MYC gene rearrangement (19 patients had confirmed MYC rearrangement alone and 24 patients also had rearrangement of BCL2, BCL6, or both [double-hit]). After a median follow-up of 56 months, 48-month EFS and OS rates were 71% and 77%, respectively for all patients.¹⁶⁸ Additional prospective studies are needed to evaluate the efficacy of DA-EPOCH-R as well as other regimens and stem cell transplantation strategies in patients with HGBL with translocations of MYC and BCL2 and/or BCL6. Alternative

treatment strategies are needed to improve outcomes in this poor-risk patient population.

The standard of care for the treatment of patients with HGBL with translocations of MYC and BCL2 and/or BCL6 nor for DEL has not been established. R-CHOP is associated with inferior outcomes. DA-EPOCH-R, R-hyperCVAD (alternating with high-dose methotrexate and cytarabine) or R-CODOX-M/R-IVAC are used in NCCN Member Institutions for the treatment of HGBL with translocations of MYC and BCL2 and/or BCL6. CNS prophylaxis should be considered according to institutional standards since these patients are at higher risk for CNS involvement.^{164,165} Consolidation with HDT/ASCR is done at some NCCN Member Institutions; however its role is not established. Consolidative RT has also been reported to result in improved PFS among patients with a CR to chemotherapy.¹⁶⁹ Currently, no data supports the use of any regimen other than R-CHOP for DEL, and additional clinical trials are needed.

Relapsed/refractory disease should be managed as described for DLBCL. However, limited data are available regarding the outcome of relapsed/refractory disease following HDT/ASCR or allogeneic HCT in patients with HGBL with translocations of MYC and BCL2 and/or BCL6 or DEL.^{170,171} Polatuzumab vedotin + BR is an appropriate treatment option for patients with relapsed or refractory HGBL with translocations of MYC and BCL2 and/or BCL6 (after ≥2 prior lines of therapies) ineligible for HDT/ASCR.¹²² CAR T-cell therapy (axicabtagene ciloleucel or tisagenlecleucel) is FDA approved for the treatment of relapsed/refractory HGBL after ≥2 prior systemic therapy regimens.¹²³⁻¹²⁵



References

1. Al-Hamadani M, Habermann TM, Cerhan JR, et al. Non-Hodgkin lymphoma subtype distribution, geodemographic patterns, and survival in the US: A longitudinal analysis of the National Cancer Data Base from 1998 to 2011. *Am J Hematol* 2015;90:790-795. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26096944>.
2. U.S. National Library of Medicine Key MEDLINE® Indicators Available at: http://www.nlm.nih.gov/bsd/bsd_key.html.
3. Alizadeh AA, Eisen MB, Davis RE, et al. Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling. *Nature* 2000;403:503-511. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10676951>.
4. Swerdlow SH, Harris NL, Jaffe ES, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Revised 4th ed. Lyon, France: IARC; 2017.
5. Hans CP, Weisenburger DD, Greiner TC, et al. Confirmation of the molecular classification of diffuse large B-cell lymphoma by immunohistochemistry using a tissue microarray. *Blood* 2004;103:275-282. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14504078>.
6. Choi WWL, Weisenburger DD, Greiner TC, et al. A new immunostain algorithm classifies diffuse large B-cell lymphoma into molecular subtypes with high accuracy. *Clin Cancer Res* 2009;15:5494-5502. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19706817>.
7. Meyer PN, Fu K, Greiner TC, et al. Immunohistochemical methods for predicting cell of origin and survival in patients with diffuse large B-cell lymphoma treated with rituximab. *J Clin Oncol* 2011;29:200-207. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21135273>.
8. Visco C, Li Y, Xu-Monette ZY, et al. Comprehensive gene expression profiling and immunohistochemical studies support application of immunophenotypic algorithm for molecular subtype classification in diffuse large B-cell lymphoma: a report from the International DLBCL
- Rituximab-CHOP Consortium Program Study. *Leukemia* 2012;26:2103-2113. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22437443>.
9. Nowakowski GS, LaPlant B, Macon WR, et al. Lenalidomide combined with R-CHOP overcomes negative prognostic impact of non-germinal center B-cell phenotype in newly diagnosed diffuse large B-Cell lymphoma: a phase II study. *J Clin Oncol* 2015;33:251-257. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25135992>.
10. Czuczman MS, Trneny M, Davies A, et al. A Phase 2/3 Multicenter, Randomized, Open-Label Study to Compare the Efficacy and Safety of Lenalidomide Versus Investigator's Choice in Patients with Relapsed or Refractory Diffuse Large B-Cell Lymphoma. *Clin Cancer Res* 2017;23:4127-4137. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28381416>.
11. Wilson WH, Young RM, Schmitz R, et al. Targeting B cell receptor signaling with ibrutinib in diffuse large B cell lymphoma. *Nat Med* 2015;21:922-926. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26193343>.
12. Savage KJ, Johnson NA, Ben-Neriah S, et al. MYC gene rearrangements are associated with a poor prognosis in diffuse large B-cell lymphoma patients treated with R-CHOP chemotherapy. *Blood* 2009;114:3533-3537. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19704118>.
13. Barrans S, Crouch S, Smith A, et al. Rearrangement of MYC is associated with poor prognosis in patients with diffuse large B-cell lymphoma treated in the era of rituximab. *J Clin Oncol* 2010;28:3360-3365. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20498406>.
14. Tzankov A, Xu-Monette ZY, Gerhard M, et al. Rearrangements of MYC gene facilitate risk stratification in diffuse large B-cell lymphoma patients treated with rituximab-CHOP. *Mod Pathol* 2014;27:958-971. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24336156>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

15. Batlle-Lopez A, Gonzalez de Villambrosia S, Francisco M, et al. Stratifying diffuse large B-cell lymphoma patients treated with chemoimmunotherapy: GCB/non-GCB by immunohistochemistry is still a robust and feasible marker. *Oncotarget* 2016;7:18036-18049. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26910115>.

16. Mozos A, Royo C, Hartmann E, et al. SOX11 expression is highly specific for mantle cell lymphoma and identifies the cyclin D1-negative subtype. *Haematologica* 2009;94:1555-1562. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19880778>.

17. Zeng W, Fu K, Quintanilla-Fend L, et al. Cyclin D1-negative blastoid mantle cell lymphoma identified by SOX11 expression. *Am J Surg Pathol* 2012;36:214-219. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22251940>.

18. Juweid ME. FDG-PET/CT in lymphoma. *Methods Mol Biol* 2011;727:1-19. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21331925>.

19. Cheson BD, Fisher RI, Barrington SF, et al. Recommendations for initial evaluation, staging, and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. *J Clin Oncol* 2014;32:3059-3068. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25113753>.

20. Adams HJ, Kwee TC, de Keizer B, et al. FDG PET/CT for the detection of bone marrow involvement in diffuse large B-cell lymphoma: systematic review and meta-analysis. *Eur J Nucl Med Mol Imaging* 2014;41:565-574. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24281821>.

21. A predictive model for aggressive non-Hodgkin's lymphoma. The International Non-Hodgkin's Lymphoma Prognostic Factors Project. *N Engl J Med* 1993;329:987-994. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8141877>.

22. Sehn LH, Berry B, Chhanabhai M, et al. The revised international prognostic index (R-IPI) is a better predictor of outcome than the standard

IPI for patients with diffuse large B-cell lymphoma treated with R-CHOP. *Blood* 2006;109:1857-1861. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17105812>.

23. Zhou Z, Sehn LH, Rademaker AW, et al. An enhanced International Prognostic Index (NCCN-IPI) for patients with diffuse large B-cell lymphoma treated in the rituximab era. *Blood* 2014;123:837-842. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24264230>.

24. El-Galaly TC, Villa D, Alzahrani M, et al. Outcome prediction by extranodal involvement, IPI, R-IPI, and NCCN-IPI in the PET/CT and rituximab era: A Danish-Canadian study of 443 patients with diffuse-large B-cell lymphoma. *Am J Hematol* 2015;90:1041-1046. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26260224>.

25. Persky DO, Unger JM, Spier CM, et al. Phase II study of rituximab plus three cycles of CHOP and involved-field radiotherapy for patients with limited-stage aggressive B-cell lymphoma: Southwest Oncology Group study 0014. *J Clin Oncol* 2008;26:2258-2263. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18413640>.

26. Pfreundschuh M, Kuhnt E, Trumper L, et al. CHOP-like chemotherapy with or without rituximab in young patients with good-prognosis diffuse large-B-cell lymphoma: 6-year results of an open-label randomised study of the MabThera International Trial (MINT) Group. *Lancet Oncol* 2011;12:1013-1022. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21940214>.

27. Held G, Murawski N, Ziepert M, et al. Role of radiotherapy to bulky disease in elderly patients with aggressive B-cell lymphoma. *J Clin Oncol* 2014;32:1112-1118. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24493716>.

28. Odejide OO, Cronin AM, Davidoff AJ, et al. Limited stage diffuse large B-cell lymphoma: comparative effectiveness of treatment strategies in a large cohort of elderly patients. *Leuk Lymphoma* 2015;56:716-724. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24913508>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

29. Lamy T, Damaj G, Soubeyran P, et al. R-CHOP 14 with or without radiotherapy in nonbulky limited-stage diffuse large B-cell lymphoma. *Blood* 2018;131:174-181. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29061568>.

30. Reyes F, Lepage E, Ganem G, et al. ACVBP versus CHOP plus radiotherapy for localized aggressive lymphoma. *N Engl J Med* 2005;352:1197-1205. Available at:

31. Recher C, Coiffier B, Haioun C, et al. Intensified chemotherapy with ACVBP plus rituximab versus standard CHOP plus rituximab for the treatment of diffuse large B-cell lymphoma (LNH03-2B): an open-label randomised phase 3 trial. *Lancet* 2011;378:1858-1867. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22118442>.

32. Peyrade F, Jardin F, Thieblemont C, et al. Attenuated immunochemotherapy regimen (R-miniCHOP) in elderly patients older than 80 years with diffuse large B-cell lymphoma: a multicentre, single-arm, phase 2 trial. *Lancet Oncol* 2011;12:460-468. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21482186>.

33. Peyrade F, Fain O, Fabiani B, et al. Long-term follow-up of the GELA LNH 03-7B study: A prospective phase II study of 150 patients over 80 years with diffuse large B-cell lymphoma (DLBCL) treated with RminiCHOP [abstract]. *J Clin Oncol* 2013;31(15_suppl):Abstract 8536. Available at: http://meeting.ascopubs.org/cgi/content/abstract/31/15_suppl/8536.

34. Vitolo U, Chiappella A, Ferreri AJ, et al. First-line treatment for primary testicular diffuse large B-cell lymphoma with rituximab-CHOP, CNS prophylaxis, and contralateral testis irradiation: final results of an international phase II trial. *J Clin Oncol* 2011;29:2766-2772. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21646602>.

35. Coiffier B, Lepage E, Briere J, et al. CHOP chemotherapy plus rituximab compared with CHOP alone in elderly patients with diffuse large-B-cell lymphoma. *N Engl J Med* 2002;346:235-242. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11807147>.

36. Coiffier B, Thieblemont C, Van Den Neste E, et al. Long-term outcome of patients in the LNH-98.5 trial, the first randomized study comparing rituximab-CHOP to standard CHOP chemotherapy in DLBCL patients: a study by the Groupe d'Etudes des Lymphomes de l'Adulte. *Blood* 2010;116:2040-2045. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20548096>.

37. Sonneveld P, van Putten W, Holte H, et al. Intensified CHOP with rituximab for intermediate or high-risk Non-hodgkin's lymphoma: interim analysis of a randomized phase III trial in elderly patients by the Dutch HOVON and Nordic Lymphoma Groups [abstract]. *Blood* 2005;106:Abstract 16. Available at: <http://www.bloodjournal.org/content/106/11/16>.

38. Habermann TM, Weller EA, Morrison VA, et al. Rituximab-CHOP versus CHOP alone or with maintenance rituximab in older patients with diffuse large B-cell lymphoma. *J Clin Oncol* 2006;24:3121-3127. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16754935>.

39. Pfreundschuh M, Schubert J, Ziepert M, et al. Six versus eight cycles of bi-weekly CHOP-14 with or without rituximab in elderly patients with aggressive CD20+ B-cell lymphomas: a randomised controlled trial (RICOVER-60). *Lancet Oncol* 2008;9:105-116. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18226581>.

40. Kuhn A, Cunningham D, Counsell N, et al. Outcome of elderly patients with diffuse large B-cell lymphoma treated with R-CHOP: results from the UK NCRI R-CHOP14v21 trial with combined analysis of molecular characteristics with the DSHNHL RICOVER-60 trial. *Ann Oncol* 2017;28:1540-1546. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28398499>.

41. Cunningham D, Hawkes EA, Jack A, et al. Rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisolone in patients with newly diagnosed diffuse large B-cell non-Hodgkin lymphoma: a phase 3 comparison of dose intensification with 14-day versus 21-day cycles. *Lancet* 2013;381:1817-1826. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23615461>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

42. Delarue R, Tilly H, Mounier N, et al. Dose-dense rituximab-CHOP compared with standard rituximab-CHOP in elderly patients with diffuse large B-cell lymphoma (the LNH03-6B study): a randomised phase 3 trial. *Lancet Oncol* 2013;14:525-533. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23578722>.

43. Schmitz N, Nickelsen M, Ziepert M, et al. Conventional chemotherapy (CHOEP-14) with rituximab or high-dose chemotherapy (MegaCHOEP) with rituximab for young, high-risk patients with aggressive B-cell lymphoma: an open-label, randomised, phase 3 trial (DSHNHL 2002-1). *Lancet Oncol* 2012;13:1250-1259. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23168367>.

44. Friedrichs B, Nickelsen M, Ziepert M, et al. Doubling rituximab in high-risk patients with aggressive B-cell lymphoma -results of the DENSE-R-MegaCHOEP trial. *Br J Haematol* 2019;184:760-768. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30520013>.

45. Huttman A, Rekowski J, Muller SP, et al. Six versus eight doses of rituximab in patients with aggressive B cell lymphoma receiving six cycles of CHOP: results from the "Positron Emission Tomography-Guided Therapy of Aggressive Non-Hodgkin Lymphomas" (PETAL) trial. *Ann Hematol* 2019. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30610279>.

46. Wilson WH, Jung SH, Porcu P, et al. A Cancer and Leukemia Group B multi-center study of DA-EPOCH-rituximab in untreated diffuse large B-cell lymphoma with analysis of outcome by molecular subtype. *Haematologica* 2012;97:758-765. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22133772>.

47. Purroy N, Bergua J, Gallur L, et al. Long-term follow-up of dose-adjusted EPOCH plus rituximab (DA-EPOCH-R) in untreated patients with poor prognosis large B-cell lymphoma. A phase II study conducted by the Spanish PETHEMA Group. *Br J Haematol* 2015;169:188-198. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25521006>.

48. Bartlett NL, Wilson WH, Jung SH, et al. Dose-Adjusted EPOCH-R Compared With R-CHOP as Frontline Therapy for Diffuse Large B-Cell Lymphoma: Clinical Outcomes of the Phase III Intergroup Trial Alliance/CALGB 50303. *J Clin Oncol* 2019:[Epub ahead of print]. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30939090>.

49. Pfreundschuh M, Muller C, Zeynalova S, et al. Suboptimal dosing of rituximab in male and female patients with DLBCL. *Blood* 2014;123:640-646. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24297867>.

50. Pfreundschuh M, Murawski N, Zeynalova S, et al. Optimization of rituximab for the treatment of DLBCL: increasing the dose for elderly male patients. *Br J Haematol* 2017;179:410-420. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28990173>.

51. Fletcher CD, Kahl BS. Central nervous system involvement in diffuse large B-cell lymphoma: an analysis of risks and prevention strategies in the post-rituximab era. *Leuk Lymphoma* 2014;55:2228-2240. Available at:

52. Hosein PJ, Maragulia JC, Salzberg MP, et al. A multicentre study of primary breast diffuse large B-cell lymphoma in the rituximab era. *Br J Haematol* 2014;165:358-363. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24467658>.

53. Ludmir EB, Milgrom SA, Pinnix CC, et al. Primary breast diffuse large B-cell lymphoma: treatment strategies and patterns of failure. *Leuk Lymphoma* 2018;59:2896-2903. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29697005>.

54. Schmitz N, Zeynalova S, Nickelsen M, et al. CNS International Prognostic Index: A Risk Model for CNS Relapse in Patients With Diffuse Large B-Cell Lymphoma Treated With R-CHOP. *J Clin Oncol* 2016;34:3150-3156. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27382100>.

55. El-Galaly TC, Villa D, Michaelsen TY, et al. The number of extranodal sites assessed by PET/CT scan is a powerful predictor of CNS relapse for patients with diffuse large B-cell lymphoma: An international multicenter



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

study of 1532 patients treated with chemoimmunotherapy. Eur J Cancer 2017;75:195-203. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28237865>.

56. Abramson JS, Hellmann M, Barnes JA, et al. Intravenous methotrexate as central nervous system (CNS) prophylaxis is associated with a low risk of CNS recurrence in high-risk patients with diffuse large B-cell lymphoma. Cancer 2010;116:4283-4290. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20564149>.

57. Holte H, Leppa S, Bjorkholm M, et al. Dose-densified chemoimmunotherapy followed by systemic central nervous system prophylaxis for younger high-risk diffuse large B-cell/follicular grade 3 lymphoma patients: results of a phase II Nordic Lymphoma Group study. Ann Oncol 2013;24:1385-1392. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23247661>.

58. Cheah CY, Herbert KE, O'Rourke K, et al. A multicentre retrospective comparison of central nervous system prophylaxis strategies among patients with high-risk diffuse large B-cell lymphoma. Br J Cancer 2014;111:1072-1079. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25072255>.

59. Ferreri AJ, Bruno-Ventre M, Donadoni G, et al. Risk-tailored CNS prophylaxis in a mono-institutional series of 200 patients with diffuse large B-cell lymphoma treated in the rituximab era. Br J Haematol 2015;168:654-662. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25312994>.

60. Kumar A, Vanderplas A, LaCasce AS, et al. Lack of benefit of central nervous system prophylaxis for diffuse large B-cell lymphoma in the rituximab era: findings from a large national database. Cancer 2012;118:2944-2951. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22006274>.

61. Tomita N, Takasaki H, Ishiyama Y, et al. Intrathecal methotrexate prophylaxis and central nervous system relapse in patients with diffuse large B-cell lymphoma following rituximab plus cyclophosphamide, doxorubicin, vincristine and prednisone. Leuk Lymphoma

2015;56:725-729. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24913502>.

62. Haioun C, Lepage E, Gisselbrecht C, et al. Survival benefit of high-dose therapy in poor-risk aggressive non-Hodgkin's lymphoma: final analysis of the prospective LNH87-2 protocol--a groupe d'Etude des lymphomes de l'Adulte study. J Clin Oncol 2000;18:3025-3030. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10944137>.

63. Le Gouill S, Milpied NJ, Lamy T, et al. First-line rituximab (R) high-dose therapy (R-HDT) versus R-CHOP14 for young adults with diffuse large B-cell lymphoma: Preliminary results of the GOELAMS 075 prospective multicenter randomized trial [abstract]. J Clin Oncol 2011;29:Abstract 8003. Available at:

http://meeting.ascopubs.org/cgi/content/abstract/29/15_suppl/8003.

64. Stiff PJ, Unger JM, Cook JR, et al. Autologous transplantation as consolidation for aggressive non-Hodgkin's lymphoma. N Engl J Med 2013;369:1681-1690. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24171516>.

65. Chiappella A, Martelli M, Angelucci E, et al. Rituximab-dose-dense chemotherapy with or without high-dose chemotherapy plus autologous stem-cell transplantation in high-risk diffuse large B-cell lymphoma (DLCL04): final results of a multicentre, open-label, randomised, controlled, phase 3 study. Lancet Oncol 2017;18:1076-1088. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28668386>.

66. Mikhaeel NG, Timothy AR, O'Doherty MJ, et al. 18-FDG-PET as a prognostic indicator in the treatment of aggressive Non-Hodgkin's Lymphoma-comparison with CT. Leuk Lymphoma 2000;39:543-553. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11342337>.

67. Spaepen K, Stroobants S, Dupont P, et al. Early restaging positron emission tomography with (18)F-fluorodeoxyglucose predicts outcome in patients with aggressive non-Hodgkin's lymphoma. Ann Oncol 2002;13:1356-1363. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12196360>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

68. Haioun C, Itti E, Rahmouni A, et al. [18F]fluoro-2-deoxy-D-glucose positron emission tomography (FDG-PET) in aggressive lymphoma: an early prognostic tool for predicting patient outcome. *Blood* 2005;106:1376-1381. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15860666>.

69. Dupuis J, Itti E, Rahmouni A, et al. Response assessment after an inductive CHOP or CHOP-like regimen with or without rituximab in 103 patients with diffuse large B-cell lymphoma: integrating 18fluorodeoxyglucose positron emission tomography to the International Workshop Criteria. *Ann Oncol* 2009;20:503-507. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19074215>.

70. Moskowitz CH, Schoder H, Teruya-Feldstein J, et al. Risk-adapted dose-dense immunochemotherapy determined by interim FDG-PET in advanced-stage diffuse large B-cell lymphoma. *J Clin Oncol* 2010;28:1896-1903. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20212248>.

71. Pregno P, Chiappella A, Bello M, et al. Interim 18-FDG-PET/CT failed to predict the outcome in diffuse large B-cell lymphoma patients treated at the diagnosis with rituximab-CHOP. *Blood* 2012;119:2066-2073. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22234681>.

72. Carr R, Fanti S, Paez D, et al. Prospective international cohort study demonstrates inability of interim PET to predict treatment failure in diffuse large B-cell lymphoma. *J Nucl Med* 2014;55:1936-1944. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25429159>.

73. Dabaja BS, Vanderplas AM, Crosby-Thompson AL, et al. Radiation for diffuse large B-cell lymphoma in the rituximab era: analysis of the National Comprehensive Cancer Network lymphoma outcomes project. *Cancer* 2015;121:1032-1039. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25492236>.

74. Mamot C, Klingbiel D, Hitz F, et al. Final Results of a Prospective Evaluation of the Predictive Value of Interim Positron Emission Tomography in Patients With Diffuse Large B-Cell Lymphoma Treated

With R-CHOP-14 (SAKK 38/07). *J Clin Oncol* 2015;33:2523-2529. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26150440>.

75. Swinnen LJ, Li H, Quon A, et al. Response-adapted therapy for aggressive non-Hodgkin's lymphomas based on early [18F] FDG-PET scanning: ECOG-ACRIN Cancer Research Group study (E3404). *Br J Haematol* 2015;170:56-65. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25823885>.

76. Duhrsen U, Muller S, Hertenstein B, et al. Positron Emission Tomography-Guided Therapy of Aggressive Non-Hodgkin Lymphomas (PETAL): A Multicenter, Randomized Phase III Trial. *J Clin Oncol* 2018;36:2024-2034. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29750632>.

77. Barrington SF, Mikhaeel NG, Kostakoglu L, et al. Role of imaging in the staging and response assessment of lymphoma: consensus of the International Conference on Malignant Lymphomas Imaging Working Group. *J Clin Oncol* 2014;32:3048-3058. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25113771>.

78. Barrington SF, Qian W, Somer EJ, et al. Concordance between four European centres of PET reporting criteria designed for use in multicentre trials in Hodgkin lymphoma. *European Journal of Nuclear Medicine and Molecular Imaging* 2010;37:1824-1833. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20505930>.

79. Meignan M, Gallamini A, Haioun C, Polliack A. Report on the Second International Workshop on interim positron emission tomography in lymphoma held in Menton, France, 8-9 April 2010. *Leuk Lymphoma* 2010;51:2171-2180. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21077737>.

80. Meignan M, Gallamini A, Itti E, et al. Report on the Third International Workshop on Interim Positron Emission Tomography in Lymphoma held in Menton, France, 26-27 September 2011 and Menton 2011 consensus. *Leuk Lymphoma* 2012;53:1876-1881. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22432519>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

81. Held G, Zeinalova S, Murawski N, et al. Impact of rituximab and radiotherapy on outcome of patients with aggressive B-cell lymphoma and skeletal involvement. *J Clin Oncol* 2013;31:4115-4122. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24062391>.
82. Thieblemont C, Tilly H, Gomes da Silva M, et al. Lenalidomide Maintenance Compared With Placebo in Responding Elderly Patients With Diffuse Large B-Cell Lymphoma Treated With First-Line Rituximab Plus Cyclophosphamide, Doxorubicin, Vincristine, and Prednisone. *J Clin Oncol* 2017;35:2473-2481. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28426350>.
83. Guppy AE, Tebbutt NC, Norman A, Cunningham D. The role of surveillance CT scans in patients with diffuse large B-cell non-Hodgkin's lymphoma. *Leuk Lymphoma* 2003;44:123-125. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12691151>.
84. Liedtke M, Hamlin PA, Moskowitz CH, Zelenetz AD. Surveillance imaging during remission identifies a group of patients with more favorable aggressive NHL at time of relapse: a retrospective analysis of a uniformly-treated patient population. *Ann Oncol* 2006;17:909-913. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16672295>.
85. Zinzani PL, Stefoni V, Tani M, et al. Role of [18F]fluorodeoxyglucose positron emission tomography scan in the follow-up of lymphoma. *J Clin Oncol* 2009;27:1781-1787. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19273712>.
86. Petrausch U, Samaras P, Haile SR, et al. Risk-adapted FDG-PET/CT-based follow-up in patients with diffuse large B-cell lymphoma after first-line therapy. *Ann Oncol* 2010;21:1694-1698. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20139151>.
87. Avivi I, Zilberlicht A, Dann EJ, et al. Strikingly high false positivity of surveillance FDG-PET/CT scanning among patients with diffuse large cell lymphoma in the rituximab era. *Am J Hematol* 2013;88:400-405. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23423884>.
88. Cheah CY, Hofman MS, Dickinson M, et al. Limited role for surveillance PET-CT scanning in patients with diffuse large B-cell lymphoma in complete metabolic remission following primary therapy. *Br J Cancer* 2013;109:312-317. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23807169>.
89. Thompson CA, Ghesquieres H, Maurer MJ, et al. Utility of routine post-therapy surveillance imaging in diffuse large B-cell lymphoma. *J Clin Oncol* 2014;32:3506-3512. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25267745>.
90. El-Galaly TC, Jakobsen LH, Hutchings M, et al. Routine Imaging for Diffuse Large B-Cell Lymphoma in First Complete Remission Does Not Improve Post-Treatment Survival: A Danish-Swedish Population-Based Study. *J Clin Oncol* 2015;33:3993-3998. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26438115>.
91. Maurer MJ, Ghesquieres H, Jais JP, et al. Event-free survival at 24 months is a robust end point for disease-related outcome in diffuse large B-cell lymphoma treated with immunochemotherapy. *J Clin Oncol* 2014;32:1066-1073. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24550425>.
92. Philip T, Guglielmi C, Hagenbeek A, et al. Autologous bone marrow transplantation as compared with salvage chemotherapy in relapses of chemotherapy-sensitive non-Hodgkin's lymphoma. *N Engl J Med* 1995;333:1540-1545. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7477169>.
93. Mounier N, Canals C, Gisselbrecht C, et al. High-dose therapy and autologous stem cell transplantation in first relapse for diffuse large B cell lymphoma in the rituximab era: an analysis based on data from the European Blood and Marrow Transplantation Registry. *Biol Blood Marrow Transplant* 2012;18:788-793. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22005647>.
94. Farooq U, Maurer MJ, Thompson CA, et al. Clinical heterogeneity of diffuse large B cell lymphoma following failure of front-line



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

immunochemotherapy. Br J Haematol 2017;179:50-60. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28653407>.

95. Spaepen K, Stroobants S, Dupont P, et al. Prognostic value of pretransplantation positron emission tomography using fluorine 18-fluorodeoxyglucose in patients with aggressive lymphoma treated with high-dose chemotherapy and stem cell transplantation. Blood 2003;102:53-59. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12609836>.

96. Derenzini E, Musuraca G, Fanti S, et al. Pretransplantation positron emission tomography scan is the main predictor of autologous stem cell transplantation outcome in aggressive B-cell non-Hodgkin lymphoma. Cancer 2008;113:2496-2503. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18833583>.

97. Hoppe BS, Moskowitz CH, Zhang Z, et al. The role of FDG-PET imaging and involved field radiotherapy in relapsed or refractory diffuse large B-cell lymphoma. Bone Marrow Transplant 2009;43:941-948. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19139730>.

98. Vose JM, Zhang MJ, Rowlings PA, et al. Autologous transplantation for diffuse aggressive Non-hodgkin's lymphoma in patients never achieving remission: a report from the autologous Blood and Marrow Transplant Registry. J Clin Oncol 2001;19:406-413. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11208832>.

99. Caballero MD, Pérez-Simón JA, Iriondo A, et al. High-dose therapy in diffuse large cell lymphoma: results and prognostic factors in 452 patients from the GEL-TAMO Spanish Cooperative Group. Annals of Oncology 2003;14:140-151. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12488306>.

100. Rodriguez J, Caballero MD, Gutierrez A, et al. Autologous stem-cell transplantation in diffuse large B-cell non-Hodgkin's lymphoma not achieving complete response after induction chemotherapy: the GEL/TAMO experience. Annals of Oncology 2004;15:1504-1509. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15367411>.

101. Vose J, Sneller V. Outpatient regimen rituximab plus ifosfamide, carboplatin and etoposide (R-ICE) for relapsed non-Hodgkin's lymphoma. Ann Oncol 2003;14 Suppl 1:17-20. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12736226>.

102. Kewalramani T, Zelenetz AD, Nimer SD, et al. Rituximab and ICE as second-line therapy before autologous stem cell transplantation for relapsed or primary refractory diffuse large B-cell lymphoma. Blood 2004;103:3684-3688. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14739217>.

103. Gisselbrecht C, Schmitz N, Mounier N, et al. Rituximab Maintenance Therapy After Autologous Stem-Cell Transplantation in Patients With Relapsed CD20+ Diffuse Large B-Cell Lymphoma: Final Analysis of the Collaborative Trial in Relapsed Aggressive Lymphoma. J Clin Oncol 2012;30:4462-4469. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23091101>.

104. Mey UJ, Orlopp KS, Flieger D, et al. Dexamethasone, high-dose cytarabine, and cisplatin in combination with rituximab as salvage treatment for patients with relapsed or refractory aggressive non-Hodgkin's lymphoma. Cancer Invest 2006;24:593-600. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16982464>.

105. Lignon J, Sibon D, Madelaine I, et al. Rituximab, dexamethasone, cytarabine, and oxaliplatin (R-DHAX) is an effective and safe salvage regimen in relapsed/refractory B-cell non-Hodgkin lymphoma. Clin Lymphoma Myeloma Leuk 2010;10:262-269. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20709662>.

106. Rigacci L, Fabbri A, Puccini B, et al. Oxaliplatin-based chemotherapy (dexamethasone, high-dose cytarabine, and oxaliplatin)+/-rituximab is an effective salvage regimen in patients with relapsed or refractory lymphoma. Cancer 2010;116:4573-4579. Available at:

107. Crump M, Kuruvilla J, Couban S, et al. Randomized comparison of gemcitabine, dexamethasone, and cisplatin versus dexamethasone, cytarabine, and cisplatin chemotherapy before autologous stem-cell transplantation for relapsed and refractory aggressive lymphomas:



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

NCIC-CTG LY.12. J Clin Oncol 2014;32:3490-3496. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25267740>.

108. Gopal AK, Press OW, Shustov AR, et al. Efficacy and safety of gemcitabine, carboplatin, dexamethasone, and rituximab in patients with relapsed/refractory lymphoma: a prospective multi-center phase II study by the Puget Sound Oncology Consortium. Leuk Lymphoma 2010;51:1523-1529. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/20578815>.

109. Hamlin PA, Zelenetz AD, Kewalramani T, et al. Age-adjusted International Prognostic Index predicts autologous stem cell transplantation outcome for patients with relapsed or primary refractory diffuse large B-cell lymphoma. Blood 2003;102:1989-1996. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12676776>.

110. Lerner RE, Thomas W, Defor TE, et al. The International Prognostic Index assessed at relapse predicts outcomes of autologous transplantation for diffuse large-cell non-Hodgkin's lymphoma in second complete or partial remission. Biol Blood Marrow Transplant 2007;13:486-492. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17382255>.

111. Thieblemont C, Briere J, Mounier N, et al. The germinal center/activated B-cell subclassification has a prognostic impact for response to salvage therapy in relapsed/refractory diffuse large B-cell lymphoma: a bio-CORAL study. J Clin Oncol 2011;29:4079-4087. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21947824>.

112. Cuccuini W, Briere J, Mounier N, et al. MYC+ diffuse large B-cell lymphoma is not salvaged by classical R-ICE or R-DHAP followed by BEAM plus autologous stem cell transplantation. Blood 2012;119:4619-4624. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22408263>.

113. Hoppe BS, Moskowitz CH, Filippa DA, et al. Involved-field radiotherapy before high-dose therapy and autologous stem-cell rescue in diffuse large-cell lymphoma: long-term disease control and toxicity. J Clin

Oncol 2008;26:1858-1864. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18332466>.

114. Lopez A, Gutierrez A, Palacios A, et al. GEMOX-R regimen is a highly effective salvage regimen in patients with refractory/relapsing diffuse large-cell lymphoma: a phase II study. Eur J Haematol 2008;80:127-132. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18005385>.

115. Corazzelli G, Capobianco G, Arcamone M, et al. Long-term results of gemcitabine plus oxaliplatin with and without rituximab as salvage treatment for transplant-ineligible patients with refractory/relapsing B-cell lymphoma. Cancer Chemother Pharmacol 2009;64:907-916. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19219604>.

116. Mounier N, El Gnaoui T, Tilly H, et al. Rituximab plus gemcitabine and oxaliplatin in patients with refractory/relapsed diffuse large B-cell lymphoma who are not candidates for high-dose therapy. A phase II Lymphoma Study Association trial. Haematologica 2013;98:1726-1731. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23753028>.

117. Weidmann E, Kim SZ, Rost A, et al. Bendamustine is effective in relapsed or refractory aggressive non-Hodgkin's lymphoma. Ann Oncol 2002;13:1285-1289. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12181253>.

118. Ohmachi K, Niitsu N, Uchida T, et al. Multicenter Phase II Study of Bendamustine Plus Rituximab in Patients With Relapsed or Refractory Diffuse Large B-Cell Lymphoma. J Clin Oncol 2013;31:2103-2109. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23650408>.

119. Vacirca JL, Acs PI, Tabbara IA, et al. Bendamustine combined with rituximab for patients with relapsed or refractory diffuse large B cell lymphoma. Ann Hematol 2014;93:403-409. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23955074>.

120. Jacobsen ED, Sharman JP, Oki Y, et al. Brentuximab vedotin demonstrates objective responses in a phase 2 study of relapsed/refractory DLBCL with variable CD30 expression. Blood



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

2015;125:1394-1402. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25573987>.

121. Crump M, Neelapu SS, Farooq U, et al. Outcomes in refractory diffuse large B-cell lymphoma: results from the international SCHOLAR-1 study. *Blood* 2017;130:1800-1808. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28774879>.

122. Sehn LH, Herrera AF, Matasar MJ, et al. Polatuzumab Vedotin (Pola) Plus Bendamustine (B) with Rituximab (R) or Obinutuzumab (G) in Relapsed/Refractory (R/R) Diffuse Large B-Cell Lymphoma (DLBCL): Updated Results of a Phase (Ph) Ib/II Study [abstract]. *Blood* 2018;132:Abstract 1683. Available at:

http://www.bloodjournal.org/content/132/Suppl_1/1683.abstract.

123. Neelapu SS, Locke FL, Bartlett NL, et al. Axicabtagene Ciloleucel CAR T-Cell Therapy in Refractory Large B-Cell Lymphoma. *N Engl J Med* 2017;377:2531-2544. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29226797>.

124. Locke FL, Ghobadi A, Jacobson CA, et al. Long-term safety and activity of axicabtagene ciloleucel in refractory large B-cell lymphoma (ZUMA-1): a single-arm, multicentre, phase 1-2 trial. *Lancet Oncol* 2019;20:31-42. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30518502>.

125. Schuster SJ, Bishop MR, Tam CS, et al. Tisagenlecleucel in Adult Relapsed or Refractory Diffuse Large B-Cell Lymphoma. *N Engl J Med* 2019;380:45-56. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30501490>.

126. Neelapu SS, Tummala S, Kebriaei P, et al. Chimeric antigen receptor T-cell therapy - assessment and management of toxicities. *Nat Rev Clin Oncol* 2018;15:47-62. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28925994>.

127. Cazals-Hatem D, Lepage E, Brice P, et al. Primary mediastinal large B-cell lymphoma. A clinicopathologic study of 141 cases compared with 916 nonmediastinal large B-cell lymphomas, a GELA ("Groupe d'Etude

des Lymphomes de l'Adulte") study. *Am J Surg Pathol* 1996;20:877-888.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8669537>.

128. Faris JE, LaCasce AS. Primary mediastinal large B-cell lymphoma. *Clin Adv Hematol Oncol* 2009;7:125-133. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19367254>.

129. Rosenwald A, Wright G, Leroy K, et al. Molecular diagnosis of primary mediastinal B cell lymphoma identifies a clinically favorable subgroup of diffuse large B cell lymphoma related to Hodgkin lymphoma. *J Exp Med* 2003;198:851-862. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12975453>.

130. Savage KJ, Monti S, Kutok JL, et al. The molecular signature of mediastinal large B-cell lymphoma differs from that of other diffuse large B-cell lymphomas and shares features with classical Hodgkin lymphoma. *Blood* 2003;102:3871-3879. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12933571>.

131. Hamlin PA, Portlock CS, Straus DJ, et al. Primary mediastinal large B-cell lymphoma: optimal therapy and prognostic factor analysis in 141 consecutive patients treated at Memorial Sloan Kettering from 1980 to 1999. *Br J Haematol* 2005;130:691-699. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16115124>.

132. Savage KJ, Al-Rajhi N, Voss N, et al. Favorable outcome of primary mediastinal large B-cell lymphoma in a single institution: the British Columbia experience. *Ann Oncol* 2006;17:123-130. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16236753>.

133. Zinzani PL, Martelli M, Bertini M, et al. Induction chemotherapy strategies for primary mediastinal large B-cell lymphoma with sclerosis: a retrospective multinational study on 426 previously untreated patients. *Haematologica* 2002;87:1258-1264. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12495899>.

134. Todeschini G, Secchi S, Morra E, et al. Primary mediastinal large B-cell lymphoma (PMLBCL): long-term results from a retrospective multicentre Italian experience in 138 patients treated with CHOP or



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

MACOP-B/VACOP-B. Br J Cancer 2004;90:372-376. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/14735179>.

135. Mazzarotto R, Boso C, Vianello F, et al. Primary mediastinal large B-cell lymphoma: results of intensive chemotherapy regimens (MACOP-B/VACOP-B) plus involved field radiotherapy on 53 patients. A single institution experience. Int J Radiat Oncol Biol Phys 2007;68:823-829. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17379431>.

136. De Sanctis V, Finolezzi E, Osti MF, et al. MACOP-B and involved-field radiotherapy is an effective and safe therapy for primary mediastinal large B cell lymphoma. Int J Radiat Oncol Biol Phys 2008;72:1154-1160. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18472357>.

137. Zinzani PL, Stefoni V, Finolezzi E, et al. Rituximab combined with MACOP-B or VACOP-B and radiation therapy in primary mediastinal large B-cell lymphoma: a retrospective study. Clin Lymphoma Myeloma 2009;9:381-385. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19858058>.

138. Rieger M, Osterborg A, Pettengell R, et al. Primary mediastinal B-cell lymphoma treated with CHOP-like chemotherapy with or without rituximab: results of the Mabthera International Trial Group study. Ann Oncol 2011;22:664-670. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/20724576>.

139. Lisenko K, Dingeldein G, Cremer M, et al. Addition of rituximab to CHOP-like chemotherapy in first line treatment of primary mediastinal B-cell lymphoma. BMC Cancer 2017;17:359. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28532396>.

140. Vassilakopoulos TP, Pangalis GA, Katsigiannis A, et al. Rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisone with or without radiotherapy in primary mediastinal large B-cell lymphoma: the emerging standard of care. Oncologist 2012;17:239-249. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22282906>.

141. Avigdor A, Sirotkin T, Kedmi M, et al. The impact of R-VACOP-B and interim FDG-PET/CT on outcome in primary mediastinal large B cell lymphoma. Ann Hematol 2014;93:1297-1304. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24595734>.

142. Soumerai JD, Hellmann MD, Feng Y, et al. Treatment of primary mediastinal B-cell lymphoma with rituximab, cyclophosphamide, doxorubicin, vincristine and prednisone is associated with a high rate of primary refractory disease. Leuk Lymphoma 2014;55:538-543. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23734654>.

143. Moskowitz C, Hamlin PA, Jr., Maragulia J, et al. Sequential dose-dense RCHOP followed by ICE consolidation (MSKCC protocol 01-142) without radiotherapy for patients with primary mediastinal large B-cell lymphoma[abstract]. Blood 2010;116:Abstract 420. Available at:
<http://www.bloodjournal.org/content/116/21/420>.

144. Dunleavy K, Pittaluga S, Maeda LS, et al. Dose-adjusted EPOCH-rituximab therapy in primary mediastinal B-cell lymphoma. N Engl J Med 2013;368:1408-1416. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23574119>.

145. Binkley MS, Hiniker SM, Wu S, et al. A single-institution retrospective analysis of outcomes for stage I-II primary mediastinal large B-cell lymphoma treated with immunochemotherapy with or without radiotherapy. Leuk Lymphoma 2015:1-5. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/26159046>.

146. Giulino-Roth L, O'Donohue T, Chen Z, et al. Outcomes of adults and children with primary mediastinal B-cell lymphoma treated with dose-adjusted EPOCH-R. Br J Haematol 2017;179:739-747. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/29082519>.

147. Shah NN, Szabo A, Huntington SF, et al. R-CHOP versus dose-adjusted R-EPOCH in frontline management of primary mediastinal B-cell lymphoma: a multi-centre analysis. Br J Haematol 2018;180:534-544. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/29265182>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

148. Kuruvilla J, Pintilie M, Tsang R, et al. Salvage chemotherapy and autologous stem cell transplantation are inferior for relapsed or refractory primary mediastinal large B-cell lymphoma compared with diffuse large B-cell lymphoma. *Leuk Lymphoma* 2008;49:1329-1336. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18604722>.

149. Vardhana S, Hamlin PA, Yang J, et al. Outcomes of Relapsed and Refractory Primary Mediastinal (Thymic) Large B Cell Lymphoma Treated with Second-Line Therapy and Intent to Transplant. *Biol Blood Marrow Transplant* 2018;24:2133-2138. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29909154>.

150. Zinzani PL, Ribrag V, Moskowitz CH, et al. Safety and tolerability of pembrolizumab in patients with relapsed/refractory primary mediastinal large B-cell lymphoma. *Blood* 2017;130:267-270. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28490569>.

151. Traverse-Glehen A, Pittaluga S, Gaulard P, et al. Mediastinal gray zone lymphoma: the missing link between classic Hodgkin's lymphoma and mediastinal large B-cell lymphoma. *Am J Surg Pathol* 2005;29:1411-1421. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16224207>.

152. Quintanilla-Martinez L, de Jong D, de Mascarel A, et al. Gray zones around diffuse large B cell lymphoma. Conclusions based on the workshop of the XIV meeting of the European Association for Hematopathology and the Society of Hematopathology in Bordeaux, France. *J Hematop* 2009;2:211-236. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20309430>.

153. Hoeller S, Copie-Bergman C. Grey zone lymphomas: lymphomas with intermediate features. *Adv Hematol* 2012;2012:460801. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22548066>.

154. Gualco G, Natkunam Y, Bacchi CE. The spectrum of B-cell lymphoma, unclassifiable, with features intermediate between diffuse large B-cell lymphoma and classical Hodgkin lymphoma: a description of 10 cases. *Mod Pathol* 2012;25:661-674. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22222636>.

155. Pilichowska M, Pittaluga S, Ferry JA, et al. Clinicopathologic consensus study of gray zone lymphoma with features intermediate between DLBCL and classical HL. *Blood Adv* 2017;1:2600-2609. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29296913>.

156. Eberle FC, Rodriguez-Canales J, Wei L, et al. Methylation profiling of mediastinal gray zone lymphoma reveals a distinctive signature with elements shared by classical Hodgkin's lymphoma and primary mediastinal large B-cell lymphoma. *Haematologica* 2011;96:558-566. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21454882>.

157. Evens AM, Kanakry JA, Sehn LH, et al. Gray zone lymphoma with features intermediate between classical Hodgkin lymphoma and diffuse large B-cell lymphoma: Characteristics, outcomes, and prognostication among a large multicenter cohort. *Am J Hematol* 2015;90:778-783. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26044261>.

158. Garcia JF, Mollejo M, Fraga M, et al. Large B-cell lymphoma with Hodgkin's features. *Histopathology* 2005;47:101-110. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15982329>.

159. Wilson WH, Pittaluga S, Nicolae A, et al. A prospective study of mediastinal gray-zone lymphoma. *Blood* 2014;124:1563-1569. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25024303>.

160. Chihara D, Westin JR, Miranda RN, et al. Dose adjusted-EPOCH-R and mediastinal disease may improve outcomes for patients with gray-zone lymphoma. *Br J Haematol* 2017;179:503-506. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27378601>.

161. Snuderl M, Kolman OK, Chen YB, et al. B-cell lymphomas with concurrent IGH-BCL2 and MYC rearrangements are aggressive neoplasms with clinical and pathologic features distinct from Burkitt lymphoma and diffuse large B-cell lymphoma. *Am J Surg Pathol* 2010;34:327-340. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20118770>.

162. Green TM, Young KH, Visco C, et al. Immunohistochemical double-hit score is a strong predictor of outcome in patients with diffuse



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

large B-cell lymphoma treated with rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisone. J Clin Oncol 2012;30:3460-3467. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22665537>.

163. Johnson NA, Slack GW, Savage KJ, et al. Concurrent expression of MYC and BCL2 in diffuse large B-cell lymphoma treated with rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisone. J Clin Oncol 2012;30:3452-3459. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22851565>.

164. Petrich AM, Gandhi M, Jovanovic B, et al. Impact of induction regimen and stem cell transplantation on outcomes in double-hit lymphoma: a multicenter retrospective analysis. Blood 2014;124:2354-2361. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25161267>.

165. Oki Y, Noorani M, Lin P, et al. Double hit lymphoma: the MD Anderson Cancer Center clinical experience. Br J Haematol 2014;166:891-901. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24943107>.

166. Sun H, Savage KJ, Karsan A, et al. Outcome of Patients With Non-Hodgkin Lymphomas With Concurrent MYC and BCL2 Rearrangements Treated With CODOX-M/IVAC With Rituximab Followed by Hematopoietic Stem Cell Transplantation. Clin Lymphoma Myeloma Leuk 2015;15:341-348. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25656914>.

167. Howlett C, Snedecor SJ, Landsburg DJ, et al. Front-line, dose-escalated immunochemotherapy is associated with a significant progression-free survival advantage in patients with double-hit lymphomas: a systematic review and meta-analysis. Br J Haematol 2015;170:504-514. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25907897>.

168. Dunleavy K, Fanale MA, Abramson JS, et al. Dose-adjusted EPOCH-R (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin, and rituximab) in untreated aggressive diffuse large B-cell lymphoma with MYC rearrangement: a prospective, multicentre,

single-arm phase 2 study. Lancet Haematol 2018;5:e609-e617. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30501868>.

169. Tumati V, Trivedi L, Li HC, et al. Patterns of Failure in Patients With Double Hit or Double Expressor Lymphomas: Implications for Radiation Therapy. Int J Radiat Oncol Biol Phys 2018;100:1126-1132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29722657>.

170. Herrera AF, Mei M, Low L, et al. Relapsed or Refractory Double-Expressor and Double-Hit Lymphomas Have Inferior Progression-Free Survival After Autologous Stem-Cell Transplantation. J Clin Oncol 2017;35:24-31. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28034071>.

171. Herrera AF, Rodig SJ, Song JY, et al. Outcomes after Allogeneic Stem Cell Transplantation in Patients with Double-Hit and Double-Expressor Lymphoma. Biol Blood Marrow Transplant 2018;24:514-520. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29196080>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Burkitt Lymphoma

Burkitt Lymphoma (BL) is a rare and aggressive B-cell lymphoma typically involving extranodal disease sites. Endemic, sporadic, and immunodeficiency-associated BL are the three clinical variants of BL that are described in the WHO classification.¹ The endemic variant is the most common form of childhood malignancy occurring in equatorial Africa, and the majority of cases are associated with Epstein-Barr virus (EBV) infection. Sporadic BL accounts for 1% to 2% of all adult lymphomas in the United States and Western Europe, and can be associated with EBV infection in about 30% of cases.² Immunodeficiency-associated BL occurs mainly in patients infected with human immunodeficiency virus (HIV), in individuals with congenital immunodeficiency, and in some patients following hematopoietic cell transplant (HCT). An analysis from the NCI SEER database reported improved survival outcomes in patients with BL diagnosed during the last decade (1922 patients diagnosed between 2002 and 2008).³ The estimated 5-year survival rate was 56% compared with 43% in patients diagnosed prior to 2002. Thus, durable remission may be possible in approximately 60% of patients with BL.

Literature Search Criteria and Guidelines Update Methodology

Prior to the update of this version of the NCCN Guidelines® for B-Cell Lymphomas an electronic search of the PubMed database was performed to obtain key literature in BL published since the previous Guidelines update. The PubMed database was chosen as it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.⁴

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.

The data from key PubMed articles deemed as relevant to these Guidelines and discussed by the panel have been included in this version of the Discussion section (eg, e-publications ahead of print, meeting abstracts). Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.

The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

Diagnosis

Adequate immunophenotyping using immunohistochemistry (IHC) with or without flow cytometry analysis is essential to establish the diagnosis of BL. The typical immunophenotype of BL is slg+, CD10+, CD19+, CD20+, CD22+, TdT-, Ki67+ (>95%), BCL2-, BCL6+, and simple karyotype with *MYC* rearrangement as sole abnormality. The IHC panel should include the following: CD3, CD10, CD20, CD45, TdT, Ki-67, BCL2, and BCL6. Flow cytometry analysis should include the following markers: CD5, CD10, CD19, CD20, CD45, TDT, and kappa/lambda. If immunophenotyping is performed using flow cytometry first, then IHC using selected markers (Ki-67 and BCL2) can supplement the findings from flow cytometry. EBV-encoded RNA in situ hybridization (EBER-ISH) may be useful to evaluate for EBV infection status in some cases.

By definition, classical BL is characterized by *MYC* gene rearrangement as the sole cytogenetic abnormality[(due to t(8;14) in 80% of cases, or its variants, t(2;8) and t(8;22) in the remaining 20% of cases], which results in the juxtaposition of *MYC* gene on chromosome 8 with the immunoglobulin heavy chain variable (*IGHV*) region on chromosome 14 or the immunoglobulin light chain genes.⁵ Cytogenetic analysis (with or without fluorescence in situ hybridization [FISH]) for detection of t(8;14) or variants should be performed in all cases with evaluation of *BCL2* or *BCL6* gene rearrangements under certain circumstances. FISH using a



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

break-apart probe, or long-segment PCR, are more reliable for the detection of t(8;14) and its variants, and are often utilized to detect these rearrangements.⁶

There are other mature aggressive B-cell lymphomas with morphologic, immunophenotypic, and/or molecular features similar to BL that do not meet this strict definition.^{7,8} Recent studies have identified a molecularly distinct subset of B-cell lymphomas with morphologic and clinical features of BL, characterized by deregulation of *11q* gene but lacking *MYC* rearrangements.⁹⁻¹² In the 2017 WHO classification, these are included as a provisional entity designated as Burkitt-like lymphoma with *11q* aberration.¹ These subset of lymphomas have a clinical course similar to BL, but the optimum management of this rare subtype is undefined, though it is most often treated like typical BL.

As discussed above, *MYC* translocations are detected in all cases of BL, but *MYC* translocations also occur in diffuse large B-cell lymphoma (DLBCL) and in a subset of B-cell lymphomas with features intermediate between BL and DLBCL.¹³ Double-hit or triple-hit lymphomas that harbor both *MYC* and *BCL2* and/or *BCL6* translocations have a highly aggressive disease course with poor prognosis (median survival of 4–6 months) and the optimal management has not been identified.¹⁴⁻¹⁸ In the 2017 classification, these lymphomas have been included in a unique category called high-grade B-cell lymphomas (HGBL) with translocations of *MYC* and *BCL2* and/or *BCL6*.¹

B-cell lymphoma, unclassifiable, with features intermediate between DLBCL and BL was added as a provisional category in the 2008 WHO I classification to include the following subsets of B-cell lymphomas with features intermediate between BL and DLBCL: lymphomas with features of both DLBCL and BL, that cannot be diagnosed as DLBCL or BL for biological and clinical reasons; lymphomas that are morphologically intermediate between BL and DLBCL include lymphomas with

immunophenotype suggestive of BL (CD10 positive, BCL6 positive, BCL2 negative, and IRF4/MUM1 negative or weakly positive); and lymphomas that are morphologically similar to BL but are strongly BCL2 positive.¹⁹ These lymphomas often present with poor prognostic factors (eg, elevated lactate dehydrogenase [LDH], bone marrow and CNS involvement, and a high International Prognostic Index [IPI]) and are associated with poor survival outcomes (median survival of 9 months and a 5-year survival rate of only 30%).¹³ In the 2017 WHO classification, the provisional category of BCLU has been eliminated and lymphomas that appear blastoid or lymphomas with features intermediate between DLBCL and BL but lack a *MYC* and *BCL2* and/or *BCL6* rearrangement are placed in the new category, HGBL, NOS.¹

Workup

The initial diagnostic workup includes a detailed physical examination (with special attention to the node-bearing areas, liver, and spleen) and chest/abdomen/pelvic CT with contrast of diagnostic quality. CT scan of the neck may be useful in certain cases. Adult patients with BL commonly present with bulky abdominal masses, B symptoms, and laboratory evidence of tumor lysis; in addition, bone marrow involvement (up to 70% of cases) and leptomeningeal central nervous system (CNS) involvement (up to 40% of cases) may also be common findings at the time of diagnosis. Brain MRI may be useful under certain circumstances (eg, if CNS involvement is suspected at time of diagnosis due to neurologic signs or symptoms). PET or integrated PET/CT scans are not recommended for routine use, since it is unlikely that findings of PET or PET/CT would alter therapy for patients with newly diagnosed BL. Multigated acquisition (MUGA) scan or echocardiogram is recommended, particularly for older patients, if the treatment includes an anthracycline-based regimen.

Evaluations of bone marrow aspirate and biopsy, lumbar puncture, and flow cytometry of cerebrospinal fluid are essential. In these highly



aggressive lymphomas, as in DLBCLs, the serum LDH level has prognostic significance. Because BL is frequently associated with HIV infection, HIV serology should be part of the diagnostic workup for these diseases (for cases with positive HIV serology, see recommendations for AIDS-related B-cell lymphomas). In addition, hepatitis B virus (HBV) testing prior to initiation of treatment is recommended for all patients who will receive anti-CD20 monoclonal antibody (MAB)-based regimens due to increased risk of HBV reactivation.

Induction Therapy

BL is curable in a significant subset of patients when treated with dose-intensive, multiagent chemotherapy regimens that also incorporate CNS prophylaxis. About 60% to 90% of pediatric and young adult patients with BL achieve durable remission if treated appropriately.²⁰ However, the survival of older adults with BL appears to be less favorable, compared with younger patients.²¹ Although the SEER database suggests that older adults (patients aged >40 years) represent about 60% of BL cases (with about 30% aged >60 years), this patient population is underrepresented in published clinical trials.^{20,21}

Most contemporary regimens used in adult patients have been developed from the pediatric protocols, and include intensive multiagent chemotherapy along with CNS prophylaxis with systemic and/or intrathecal chemotherapy. Tumor lysis syndrome is more common in patients with BL and should be managed as outlined under “Tumor Lysis Syndrome” in the *Supportive Care* section of the algorithm.

It is important to note that CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) or similar regimens are not considered adequate therapy for the management of BL. Results from retrospective analyses have shown that more intensive multiagent chemotherapy regimens such as hyper-CVAD (hyperfractionated cyclophosphamide,

vincristine, doxorubicin, and dexamethasone alternating with methotrexate and cytarabine, including intrathecal methotrexate) or CODOX-M/IVAC (cyclophosphamide, doxorubicin, and high-dose methotrexate, alternating with ifosfamide, etoposide, and high-dose cytarabine) are associated with more favorable outcomes than CHOP or CHOP plus etoposide.^{22,23} In a population-based analysis of data from 258 patients with HIV-negative BL from a Swedish/Danish registry, the 2-year overall survival (OS) rate was only 39% for CHOP or CHOP with etoposide compared with 83%, and 69%, respectively, for hyper-CVAD and CODOX-M/IVAC.²²

Thus, for patients with BL who can tolerate aggressive therapies, intensive multiagent chemotherapy may offer the best chance for durable disease control. Data from clinical trials that have evaluated induction therapy with more intensive multiagent chemotherapy regimens are discussed below.

CODOX-M/IVAC (originally published in 1998) is a highly effective regimen with a 1-year event-free survival (EFS) rate of 85% in pediatric and adult patients with previously untreated BL or Burkitt-like lymphoma.^{24,25} Patients with high-risk disease received 4 alternating treatments (ABAB) of CODOX-M (A) and IVAC (B), and those with low-risk disease received 3 cycles of CODOX-M. Both cycles included intrathecal chemotherapy (cytarabine or methotrexate) for CNS prophylaxis in addition to high-dose systemic cytarabine and methotrexate.

Subsequent phase II and retrospective studies have confirmed the efficacy of this regimen, and a “modified” regimen (inclusion of rituximab and dose-modified to decrease toxicity).^{15,26-29}

The efficacy of modified CODOX-M/IVAC regimen (modified slightly with vincristine dose capped at 2 mg) in adult patients with BL was confirmed in an international phase II study (n =52; 12 patients had low-risk disease; 40 patients had high-risk disease).²⁶ The overall 2-year EFS and OS rates were 65% and 73%, respectively. The 2-year EFS and OS rates were 83%



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

and 81%, respectively, for patients with low-risk disease. The corresponding survival rates were 60% and 70%, respectively, for those with high-risk disease.²⁶ Subsequent reports showed that lowering the dose of systemic methotrexate to 3 g/m² from the originally reported 7 g/m² maintained the efficacy and was also associated with decreased toxicity, with a 2-year progression-free survival (PFS) rate of 62% to 64% and 2-year OS rate of 71%.^{15,27} Another small study that evaluated modified CODOX-M/IVAC (reducing the dose of methotrexate to 3 g/m²) with or without rituximab in 15 patients with BL or B-cell lymphoma unclassifiable, also reported 5-year PFS and OS rates of 87%.²⁸

The addition of rituximab to modified CODOX-M/IVAC was evaluated in a retrospective study of 80 patients with BL.²⁹ There was a trend for improvement in outcomes with the addition of rituximab but the differences were not statistically significant. The 3-year PFS and OS rates with rituximab were 74% and 77%, respectively; the 3-year PFS and OS rates without the addition of rituximab were 61% and 66%, respectively.

An additional modification of CODOX-M/IVAC utilizing liposomal doxorubicin instead of doxorubicin, plus rituximab, has also been shown to be effective in patients with newly diagnosed classic BL.^{30,31} In a phase II trial of 25 patients, this modified CODOX-M/IVAC regimen resulted in an overall response rate of (ORR) of 100% (92% CR).³¹ At 34-month median follow-up, the 2-year PFS and OS rates for all patients were 80% and 84%, respectively (PFS and OS rates were both 100% for patients with low-risk BL and the corresponding survival rates were 76% and 81%, respectively, for those with high-risk BL). The 2-year PFS and OS rates for high-risk, HIV-negative patients were 84% and 89%, respectively.

The hyper-CVAD regimen is also effective, resulting in CR rates of 81% and a 3-year OS rate of 49% in patients with BL or leukemia.³² The OS rate was higher among patients ≤60 years (77% vs. 17% for patients >60 years). The results of a phase II study (57 patients; 30 patients with

non-HIV BL and 27 patients with B-ALL) showed that the addition of rituximab to hyper-CVAD (R-hyper-CVAD) improved long-term outcomes in patients with newly diagnosed BL or B-ALL, particularly in the older patient subgroup.^{33,34} R-hyper-CVAD induced complete response (CR) in 86% of patients and the 5-year OS rate was 74%, with a median follow-up of 62 months.³⁴ The 5-year OS rates were 72% and 70%, respectively, in patients <60 years and those >60 years. These survival outcomes were superior in historical comparison with patients treated with hyper-CVAD alone.

Dose-adjusted EPOCH with rituximab (DA-EPOCH-R) also resulted in highly favorable outcomes in patients with previously untreated BL. In a prospective study that evaluated DA-EPOCH-R in HIV-negative patients (n = 19) and a short course of EPOCH with dose-dense rituximab (SC-EPOCH-RR) in HIV-positive patients (n = 11), at a median follow-up of 86 months, the freedom-from-progression (FFP) and OS rates with DA-EPOCH-R were 95% and 100%, respectively.³⁵ The highly favorable outcomes seen in this study may reflect the inclusion of more low-risk patients compared to other studies, with approximately 53% of all patients (37% in the DA-EPOCH-R group) presenting with normal LDH levels. A risk-adapted treatment approach with DA-EPOCH-R was validated in a subsequent multicenter phase II trial (n = 113).³⁶ Low-risk patients received 2 cycles of DA-EPOCH-R without intrathecal therapy followed by PET and one more cycle, if interim PET was negative. High-risk patients with no CNS involvement received 2 cycles of DA-EPOCH-R without intrathecal therapy followed by PET and 4 additional cycles of DA-EPOCH-R, unless interim PET showed progression. After a median follow-up of 36 months, the PFS, FFP, and OS rates were 86%, 92% and 86%, respectively.

A large, prospective, multicenter study from the German study group evaluated the efficacy and safety of a short-intensive multiagent



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

chemotherapy regimen (including high-dose methotrexate, high-dose cytarabine, cyclophosphamide, etoposide, ifosfamide, and corticosteroids) combined with rituximab in patients with BL (n = 225) and Burkitt leukemia (n = 138).³⁷ Patients also received triple intrathecal therapy with methotrexate, cytarabine, and dexamethasone. The CR rate was 88% for the total cohort. At a median follow-up of >7 years, the PFS and OS rates were 71% and 80%, respectively. The diagnosis of BL versus Burkitt-like lymphoma had no influence on outcome. These outcomes appear highly promising, with a manageable toxicity profile. The results of a more recent randomized phase III trial (n = 260) from the French lymphoma study group (GELA) also showed that the addition of rituximab to short intensive chemotherapy is associated with improved EFS in adults with BL.³⁸ With a median follow-up of 38 months, the 3-year EFS rates were significantly higher for patients treated with rituximab and dose-dense chemotherapy than for patients treated with dose-dense chemotherapy alone (75% vs. 62%; $P = .024$).

Several studies have evaluated the role of HCT in patients with BL. The Dutch-Belgian Hemato-Oncology Cooperative Group (HOVON) demonstrated the feasibility of intensive high-dose induction chemotherapy (prednisone, cyclophosphamide, doxorubicin, etoposide, and mitoxantrone, without high-dose methotrexate or high-dose cytarabine) followed by consolidation with BEAM and autologous HCT in untreated BL, Burkitt-like lymphoma, or B-ALL.³⁹ Among the patients with BL/Burkitt-like lymphoma (n = 27), CR was achieved in 81% of patients with a partial response (PR) in 11%; the 5-year EFS and OS rates were 73% and 81%, respectively.³⁹ In a recent analysis of outcomes with HCT (autologous or allogeneic transplant) in patients with BL from the CIBMTR database (N = 241), the 5-year PFS and OS rates with autologous HCT at first remission were 78% and 83%, respectively.⁴⁰ These outcomes with autologous HCT were similar to findings from the above HOVON study, and appeared to compare favorably to the 5-year PFS and OS rates with

allogeneic HCT in first remission, which were 50% and 53%, respectively. Not surprisingly, patients who underwent HCT with less than a first remission had poorer outcomes regardless of transplant type. The 5-year PFS and OS rates with autologous HCT in those without a first remission were 27% and 31%, respectively; the corresponding rates with allogeneic HCT without first remission were only 19% and 20%, respectively. For patients in a second remission, autologous HCT resulted in a 5-year PFS of 44%.⁴⁰ An earlier retrospective analysis from the CIBMTR database in patients with relapsed or refractory BL (children and adolescents age ≤18 years; n = 41) showed similar 5-year EFS outcomes between autologous and allogeneic HCT (27% vs. 31%).⁴¹ As would be expected, EFS rates were lower among patients who were not in CR at the time of transplant.

NCCN Recommendations

It is preferred that patients with BL receive treatment at centers with expertise in the management of this highly aggressive disease. Participation in clinical trials is recommended for all patients. As mentioned earlier, CHOP or CHOP-like therapy is not adequate for the treatment of BL.^{22,23} The management of patients with B-cell lymphomas with features intermediate between BL and DLBCL (now included in the new category, HGBL, NOS) as well as those patients with double-hit or triple-hit lymphomas (HGBL with translocations of *MYC* and *BCL2* and/or *BCL6*) has not been well studied. Therefore, these patients are best managed in the context of clinical trials evaluating novel targeted agents.

Patients with either of the following clinical characteristics are generally considered to have low-risk disease: normal serum LDH or stage I disease and a completely resected abdominal lesion or a single extra-abdominal mass <10 cm. All other patients should be considered as having high-risk disease.

CODOX-M (original or modified) ± rituximab (3 cycles), DA-EPOCH-R (3 cycles with one more additional cycle for patients achieving CR), and



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

hyper-CVAD + rituximab are included as options for patients with low-risk BL.

CODOX-M/IVAC (original or modified) with or without rituximab, DA-EPOCH-R (for patients not able to tolerate aggressive treatments), and R-hyper-CVAD are included as options for patients with high-risk BL.

Adequate CNS prophylaxis with systemic and/or intrathecal chemotherapy with methotrexate and/or cytarabine is recommended with all regimens.

Patients with CR to induction therapy should be followed up every 2 to 3 months for 1 year then every 3 months for the next 1 year, and then every 6 months thereafter. Disease relapse after 2 years is rare following CR to induction therapy, and follow-up should be individualized according to patient characteristics. Consolidation therapy in the context of a clinical trial may be considered for high-risk patients with CR to induction therapy. Patients with less than CR to induction therapy should be treated in the context of a clinical trial. In the absence of suitable clinical trials palliative involved-site radiation therapy (ISRT) may be considered appropriate.

Relapsed or Refractory Disease

Patients with relapsed or refractory disease should be treated in the context of a clinical trial. In the absence of suitable clinical trials, selected patients with a reasonable remission duration following induction therapy may be managed with second-line therapy using multiagent chemotherapy regimens.

DA-EPOCH-R, IVAC combined with rituximab (R-IVAC), R-GDP (gemcitabine, dexamethasone, cisplatin, combined with rituximab), R-ICE (ifosfamide, carboplatin, etoposide, combined with rituximab), and high-dose cytarabine are included as options for second-line therapy. However, it should be noted that treatment options remain undefined for patients who relapse after first-line therapy and these suggestions are

based on very limited, retrospective studies with only a few patients. For instance, the R-ICE regimen was evaluated in a small group of pediatric patients with relapsed BL and B-ALL (n = 14), which resulted in CR in 29% and PR in 36% of patients.⁴²

Consolidation with high-dose therapy and autologous stem cell rescue (HDT/ASCR) or allogeneic HCT (if donor available) may be considered for selected patients achieving a CR or PR to second-line therapy. Clinical trial or best supportive care including palliative ISRT are recommended for patients with disease not responding to second-line therapy or those with progressive disease.



References

1. Swerdlow SH, Harris NL, Jaffe ES, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Revised 4th ed. Lyon, France: IARC; 2017.
2. Blum KA, Lozanski G, Byrd JC. Adult Burkitt leukemia and lymphoma. *Blood* 2004;104:3009-3020. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15265787>.
3. Costa LJ, Xavier AC, Wahlquist AE, Hill EG. Trends in survival of patients with Burkitt lymphoma/leukemia in the USA: an analysis of 3691 cases. *Blood* 2013;121:4861-4866. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23641015>.
4. U.S. National Library of Medicine Key MEDLINE® Indicators Available at: http://www.nlm.nih.gov/bsd/bsd_key.html.
5. Ferry JA. Burkitt's lymphoma: clinicopathologic features and differential diagnosis. *Oncologist* 2006;11:375-383. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16614233>.
6. Burmeister T, Schwartz S, Horst HA, et al. Molecular heterogeneity of sporadic adult Burkitt-type leukemia/lymphoma as revealed by PCR and cytogenetics: correlation with morphology, immunology and clinical features. *Leukemia* 2005;19:1391-1398. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15973450>.
7. Macpherson N, Lesack D, Klasa R, et al. Small noncleaved, non-Burkitt's (Burkitt-Like) lymphoma: cytogenetics predict outcome and reflect clinical presentation. *J Clin Oncol* 1999;17:1558-1567. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10334544>.
8. Hummel MI, Bentink S, Berger H, et al. A biologic definition of Burkitt's lymphoma from transcriptional and genomic profiling. *N Engl J Med* 2006;354:2419-2430. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16760442>.
9. Ferreira JF, Morscio J, Dierickx D, et al. Post-transplant molecularly defined Burkitt lymphomas are frequently MYC-negative and characterized by the 11q-gain/loss pattern. *Haematologica* 2015;100:e275-279. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25795716>.
10. Salaverria I, Martin-Guerrero I, Wagener R, et al. A recurrent 11q aberration pattern characterizes a subset of MYC-negative high-grade B-cell lymphomas resembling Burkitt lymphoma. *Blood* 2014;123:1187-1198. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24398325>.
11. Rymkiewicz G, Grygalewicz B, Chechlinska M, et al. A comprehensive flow-cytometry-based immunophenotypic characterization of Burkitt-like lymphoma with 11q aberration. *Mod Pathol* 2018;31:732-743. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29327714>.
12. Wagener R, Seufert J, Raimondi F, et al. The mutational landscape of Burkitt-like lymphoma with 11q aberration is distinct from that of Burkitt lymphoma. *Blood* 2019;133:962-966. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30567752>.
13. Perry AM, Crockett D, Dave BJ, et al. B-cell lymphoma, unclassifiable, with features intermediate between diffuse large B-cell lymphoma and burkitt lymphoma: study of 39 cases. *Br J Haematol* 2013;162:40-49. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23600716>.
14. Le Gouill S, Talmant P, Touzeau C, et al. The clinical presentation and prognosis of diffuse large B-cell lymphoma with t(14;18) and 8q24/c-MYC rearrangement. *Haematologica* 2007;92:1335-1342. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18024371>.
15. Mead GM, Barrans SL, Qian W, et al. A prospective clinicopathologic study of dose-modified CODOX-M/IVAC in patients with sporadic Burkitt lymphoma defined using cytogenetic and immunophenotypic criteria (MRC/NCRI LY10 trial). *Blood* 2008;112:2248-2260. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18612102>.
16. Tomita N, Tokunaka M, Nakamura N, et al. Clinicopathological features of lymphoma/leukemia patients carrying both BCL2 and MYC translocations. *Haematologica* 2009;94:935-943. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19535347>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

17. Johnson NA, Savage KJ, Ludkovski O, et al. Lymphomas with concurrent BCL2 and MYC translocations: the critical factors associated with survival. *Blood* 2009;114:2273-2279. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19597184>.

18. Snuderl M, Kolman OK, Chen YB, et al. B-cell lymphomas with concurrent IGH-BCL2 and MYC rearrangements are aggressive neoplasms with clinical and pathologic features distinct from Burkitt lymphoma and diffuse large B-cell lymphoma. *Am J Surg Pathol* 2010;34:327-340. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20118770>.

19. Swerdlow SH, Campo E, Harris NL, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues, Fourth Edition, Lyon, France: IARC; ; 2008.

20. Perkins AS, Friedberg JW. Burkitt lymphoma in adults. *Hematology Am Soc Hematol Educ Program* 2008:341-348. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19074108>.

21. Kelly JL, Tothaker SR, Ciminello L, et al. Outcomes of patients with Burkitt lymphoma older than age 40 treated with intensive chemotherapeutic regimens. *Clin Lymphoma Myeloma* 2009;9:307-310. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19717381>.

22. Wasterlid T, Brown PN, Hagberg O, et al. Impact of chemotherapy regimen and rituximab in adult Burkitt lymphoma: a retrospective population-based study from the Nordic Lymphoma Group. *Ann Oncol* 2013;24:1879-1886. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23446093>.

23. Oosten LEM, Chamuleau MED, Thielen FW, et al. Treatment of sporadic Burkitt lymphoma in adults, a retrospective comparison of four treatment regimens. *Ann Hematol* 2018;97:255-266. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29209924>.

24. Magrath I, Adde M, Shad A, et al. Adults and children with small non-cleaved-cell lymphoma have a similar excellent outcome when treated

with the same chemotherapy regimen. *J Clin Oncol* 1996;14:925-934. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8622041>.

25. Adde M, Shad A, Venzon D, et al. Additional chemotherapy agents improve treatment outcome for children and adults with advanced B-cell lymphomas. *Semin Oncol* 1998;25:33-39; discussion 45-38. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9578060>.

26. Mead GM, Sydes MR, Walewski J, et al. An international evaluation of CODOX-M and CODOX-M alternating with IVAC in adult Burkitt's lymphoma: results of United Kingdom Lymphoma Group LY06 study. *Ann Oncol* 2002;13:1264-1274. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12181251>.

27. Lacasce A, Howard O, Lib S, et al. Modified Magrath regimens for adults with Burkitt and Burkitt-like lymphomas: preserved efficacy with decreased toxicity. *Leuk Lymphoma* 2004;45:761-767. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15160953>.

28. Maruyama D, Watanabe T, Maeshima AM, et al. Modified cyclophosphamide, vincristine, doxorubicin, and methotrexate (CODOX-M)/ifosfamide, etoposide, and cytarabine (IVAC) therapy with or without rituximab in Japanese adult patients with Burkitt lymphoma (BL) and B cell lymphoma, unclassifiable, with features intermediate between diffuse large B cell lymphoma and BL. *Int J Hematol* 2010;92:732-743. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21120644>.

29. Barnes JA, Lacasce AS, Feng Y, et al. Evaluation of the addition of rituximab to CODOX-M/IVAC for Burkitt's lymphoma: a retrospective analysis. *Ann Oncol* 2011;22:1859-1864. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21339382>.

30. Corazzelli G, Frigeri F, Russo F, et al. RD-CODOX-M/IVAC with rituximab and intrathecal liposomal cytarabine in adult Burkitt lymphoma and 'unclassifiable' highly aggressive B-cell lymphoma. *Br J Haematol* 2012;156:234-244. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22098541>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

31. Evens AM, Carson KR, Kolesar J, et al. A multicenter phase II study incorporating high-dose rituximab and liposomal doxorubicin into the CODOX-M/IVAC regimen for untreated Burkitt's lymphoma. *Ann Oncol* 2013;24:3076-3081. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24146219>.

32. Thomas DA, Cortes J, O'Brien S, et al. Hyper-CVAD program in Burkitt's-type adult acute lymphoblastic leukemia. *J Clin Oncol* 1999;17:2461-2470. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10561310>.

33. Thomas DA, Faderl S, O'Brien S, et al. Chemoimmunotherapy with hyper-CVAD plus rituximab for the treatment of adult Burkitt and Burkitt-type lymphoma or acute lymphoblastic leukemia. *Cancer* 2006;106:1569-1580. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16502413>.

34. Thomas DA, Kantarjian HM, Faderl S, et al. Hyper-CVAD and Rituximab for De Novo Burkitt Lymphoma/Leukemia [abstract]. *Blood* 2011;118:Abstract 2698. Available at:

<http://www.bloodjournal.org/content/118/21/2698>.

35. Dunleavy K, Pittaluga S, Shovlin M, et al. Low-intensity therapy in adults with Burkitt's lymphoma. *N Engl J Med* 2013;369:1915-1925. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24224624>.

36. Roschewski M, Dunleavy K, Abramson JS, et al. Risk-Adapted Therapy in Adults with Burkitt Lymphoma: Results of NCI 9177, a Multicenter Prospective Phase II Study of DA-EPOCH-R [abstract]. *Blood* 2017;130:Abstract 188. Available at:

http://www.bloodjournal.org/content/130/Suppl_1/188.abstract.

37. Hoelzer D, Walewski J, Dohner H, et al. Improved outcome of adult Burkitt lymphoma/leukemia with rituximab and chemotherapy: report of a large prospective multicenter trial. *Blood* 2014;124:3870-3879. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25359988>.

38. Ribrag V, Koscielny S, Bosq J, et al. Rituximab and dose-dense chemotherapy for adults with Burkitt's lymphoma: a randomised,

controlled, open-label, phase 3 trial. *Lancet* 2016;387:2402-2411.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27080498>.

39. van Imhoff GW, van der Holt B, MacKenzie MA, et al. Short intensive sequential therapy followed by autologous stem cell transplantation in adult Burkitt, Burkitt-like and lymphoblastic lymphoma. *Leukemia* 2005;19:945-952. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15800666>.

40. Maramattom LV, Hari PN, Burns LJ, et al. Autologous and allogeneic transplantation for burkitt lymphoma outcomes and changes in utilization: a report from the center for international blood and marrow transplant research. *Biol Blood Marrow Transplant* 2013;19:173-179. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23200705>.

41. Gross TG, Hale GA, He W, et al. Hematopoietic stem cell transplantation for refractory or recurrent non-Hodgkin lymphoma in children and adolescents. *Biol Blood Marrow Transplant* 2010;16:223-230. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19800015>.

42. Griffin TC, Weitzman S, Weinstein H, et al. A study of rituximab and ifosfamide, carboplatin, and etoposide chemotherapy in children with recurrent/refractory B-cell (CD20+) non-Hodgkin lymphoma and mature B-cell acute lymphoblastic leukemia: a report from the Children's Oncology Group. *Pediatr Blood Cancer* 2009;52:177-181. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18816698>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

AIDS-Related B-Cell Lymphomas

Non-Hodgkin's lymphomas (NHL), Kaposi sarcoma (KS) and lung cancer are the most common cancer types diagnosed in people living with human immunodeficiency virus (HIV) in the United States and the incidences of cancer in this patient population is expected to shift substantially through 2030, with the largest declines projected for NHL and KS.^{1,2}

Diffuse large B-cell lymphoma (DLBCL), Burkitt lymphoma (BL), and primary central nervous system lymphoma (PCNSL) are the most common subtypes of NHL in people living with HIV.³ The distribution of systemic lymphomas versus PCNSL may vary between published reports depending upon the different factors such as geographic regions, time period covered and referral patterns of the institutions. The incidences of Hodgkin lymphoma (HL) and indolent lymphoma are also elevated in people living with HIV, but are much less common than BL or DLBCL.³ Plasmablastic lymphoma (PBL) and primary effusion lymphoma (PEL) are the less common forms of systemic lymphomas, accounting for less than 5% of lymphomas in people living with HIV.

Literature Search Criteria and Guidelines Update Methodology

Prior to the update of this version of the NCCN Guidelines® for B-Cell Lymphomas an electronic search of the PubMed database was performed to obtain key literature in AIDS-related lymphomas published since the previous Guidelines update, using the following search terms: AIDS-related lymphomas or HIV-associated lymphomas or HIV-positive lymphomas. The PubMed database was chosen as it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.⁴

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types:

Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.

The PubMed search resulted in 26 citations and their potential relevance was examined. The data from key PubMed articles deemed as relevant to these Guidelines and discussed by the panel have been included in this version of the Discussion section (eg, e-publications ahead of print, meeting abstracts). Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.

The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

Prognosis

The routine use of antiretroviral therapy (ART) has improved the prognosis of patients with HIV-associated lymphomas and the pathologic spectrum of lymphomas in people living with HIV has also changed in the era of ART, with a drastic decrease in the incidence of PBL and a much lower incidence of systemic HIV-associated lymphomas.^{5,6} Nevertheless, HIV continues to be independently associated with increased risk of death among patients with HIV-associated lymphomas even in the ART era in the United States, and the association varies by lymphoma subtype.⁷⁻⁹ Increased incidence of lymphoma due to unmasking immune reconstitution inflammatory syndrome soon after receiving ART has also been described in people living with HIV.¹⁰

In a report from the COHERE (Collaboration of Observational HIV Epidemiological Research Europe) study that evaluated the outcomes of patients with HIV-associated lymphomas treated in the ART era (1998-2006), the 1-year OS rates among patients with systemic lymphoma and PCNSL were 66% and 54%, respectively.⁸ In large cohort



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

study that evaluated the trends in presentation and survival of lymphoma a large HIV-infected cohort, the 5-year survival rates were 50%, 44%, 23% and 43% respectively for DLBCL, BL, PCNSL and other NHL subtypes.³ Older age, lymphoma occurrence during antiretroviral therapy, lower CD4 count at lymphoma diagnosis, higher HIV RNA and histologic category were identified as independent risk factors for mortality. In a recent analysis that evaluated the characteristics and outcomes of DLBCL in HIV-infected patients in the era of ART, the 2-year OS and progression-free survival (PFS) were both 75% after treatment with RCHOP; PFS after treatment with RCHOP did not differ from that of the HIV-negative counterparts.¹¹

PCNSL in people with HIV may have a better prognosis and is actually curable with less intensive chemotherapy regimens when used along with immune reconstitution.¹²⁻¹⁵ Immune reconstitution with ART administered along with high-dose methotrexate or radiation therapy (RT) is associated with improved survival outcomes in patients with HIV-associated PCNSL, even among those with a history of opportunistic infections, limited access to healthcare and medical non-adherence.¹⁴

PBL is an aggressive CD20-negative large B-cell lymphoma that mainly involves the jaw and oral cavity in people living with HIV and is also associated with EBV infection.^{16,17} The prognosis of PBL has improved in the ART era even in patients with higher stage disease and more extranodal involvement, with some case series reporting favorable outcomes for patients with HIV-associated PBL treated with anthracycline-based multiagent chemotherapy in conjunction with ART.¹⁸⁻²⁰ In a cohort study of 61 patients with PBL, age (<50 years of age) and early stage disease (stage I/II disease) were identified as prognostic factors for favorable survival outcomes.¹⁹ EBV-positive status was associated with a better event-free survival (EFS) compared to EBV-negative status. In a case series of 12 patients with newly diagnosed PBL treated exclusively in the ART era at the AIDS

Malignancy Consortium (AMC) sites, at a median follow up of 73 weeks, the 1-year survival rate was 67% with no reported deaths in the follow up period after one year.²⁰ However, other studies suggest that the prognosis of patients with HIV-associated PBL remains poor even in the ART era.²¹⁻²⁴ In the German AIDS-related-Lymphoma-Cohort-Study, the 2-year OS rate for patients with PBL was 43% compared to the 2-year OS rates of 69% and 63% respectively for patients with BL and DLBCL.²³ PBL histology, international prognostic index (IPI) and bone marrow involvement were identified as independent predictors of mortality.

PEL is characterized by neoplastic effusions in body cavities without detectable tumor masses (occurring most often in the pleural, pericardial, and abdominal cavities) and is associated with Kaposi sarcoma-associated herpesvirus (KSHV), otherwise known as human herpes virus 8 (HHV8) and it may also be co-infected with Epstein Barr virus (EBV).^{25,26} PEL has a poor survival compared to HIV-associated DLBCL or BL, even in the ART era. In a study that analyzed the outcomes of 10,769 patients with HIV-associated lymphomas identified in the National Cancer Data Base, the estimated 5-year OS rates were 42%, 45%, 22% and 28% respectively for DLBCL, BL, PCNSL, and PEL.²⁷ Poor performance status, advanced stage disease and absence of ART prior to the diagnosis of PEL have been identified as prognostic factors for shorter survival.^{28,29} The use of ART with chemotherapy is essential to improve the outcomes of PEL.

Diagnosis

The major factor in the diagnostic evaluation of HIV-associated lymphoma is to distinguish between different subtypes. Adequate immunophenotyping using immunohistochemistry (IHC) with or without flow cytometry analysis is essential to establish the diagnosis of the subtype of HIV-associated lymphomas.³⁰ Epstein-Barr virus (EBV)-encoded RNA in situ hybridization (EBER-ISH) is recommended



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

for all patients since EBV is the most commonly found oncogenic virus in patients HIV-associated lymphomas.³¹ HHV8/KHSV testing would be useful to confirm the diagnosis of PEL since this oncogenic virus is implicated in the pathogenesis of PEL.^{25,26}

Workup

The diagnostic evaluation and workup are as outlined in the NCCN Guidelines section for HIV-associated lymphomas. However, all patients (without regard to histology) should have a lumbar puncture to rule out CNS involvement. In addition, baseline values for CD4 counts and HIV viral load should be obtained. Among the systemic HIV-associated lymphomas, BL is generally associated with a higher CD4+ cell count at diagnosis compared with DLBCL. PCNSL, PBL and PEL are particularly associated with low CD4+ count levels.

Initial Treatment

Several key factors have emerged as being important to improve the outcome in patients with HIV-associated lymphomas. The introduction of ART has allowed for the administration of more dose-intensive chemotherapy regimens and a reduction in treatment-associated toxicity. In addition, the use of concurrent ART is also associated with superior outcomes (improved CR rates, a trend towards improved OS and faster immune recovery).³²⁻³⁷

In prospective phase II studies, several combination chemotherapy regimens (with or without rituximab) given with concomitant ART, have been proven to be active and tolerable in patients with HIV-associated lymphomas. In a pooled analysis of 1546 patients with HIV-associated lymphomas included in prospective clinical trials, initial therapy with more dose-intensive regimens and the use of rituximab resulted in higher CR rates; the use of rituximab was also associated improved PFS and OS.³⁵

EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin) + rituximab

In the AMC 034 trial, patients with HIV-associated aggressive B-cell NHL were randomized to receive infusional EPOCH regimen either concurrently with rituximab or followed sequentially by rituximab (106 patients with HIV-associated lymphomas; 75% DLBCL; 25% BL, BL-like).³⁸ The complete response (CR) rate was 73% and 55%, respectively for concurrent and sequential rituximab. The 2-year PFS rate (66% vs. 63%) and OS rate (70% vs. 67%) were similar between treatment arms. Toxicity was comparable in the 2 treatment arms, although the use of concurrent rituximab was associated with a higher incidence of treatment-related deaths among the patients with a baseline CD4+ count <50/mcL. Overall, treatment-related deaths occurred in 5 patients (10%) in the concurrent rituximab arm and 4 patients (7%) in the sequential rituximab arm. In this trial, ART was given concurrently with EPOCH or delayed until chemotherapy completion per investigator choice. Although the use of concurrent ART was not associated with improved survival outcomes compared to delayed use of ART, concurrent use of ART was associated with faster immune recovery (CD4 counts higher than baseline 6 months after EPOCH and decrease in HIV viral load during chemotherapy).³⁷

In a pooled analysis that included 150 patients with HIV-associated NHL treated in the AMC trials (AMC 010 and AMC 034), low age-adjusted IPI score and baseline CD4 count ≥100/mcL were significantly associated with improved CR rate, EFS and OS outcomes.³⁹ Among the patients who were treated with concurrent EPOCH-R, both EFS and OS were significantly improved compared with RCHOP (after adjusting for aalPI and CD4 counts). The incidence of treatment-related deaths were higher in patients with baseline CD4 counts <50/mcL compared with those with higher CD4 counts (37% vs. 6%; $P<0.01$).³⁹



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Dose-adjusted EPOCH (DA-EPOCH) is also effective resulting in an ORR of 87% with a CR rate of 74% in patients with untreated HIV-associated NHL (n = 39; 79% DLBCL; 18% BL).⁴⁰ At a median follow up of 53 months, PFS and OS rates were 73% and 60%, respectively. The disease-free survival (DFS) rate was 92% with only 2 of the patients with a CR experiencing disease recurrence at last follow up). The OS outcomes were decreased among the patients with low baseline CD4 counts ($\leq 100/\text{mCL}$) compared with those with higher CD4 counts (16% vs. 87%). In a multivariate analysis, low CD4 counts and CNS involvement were the only significant factors associated with decreased OS.⁴⁰

In a more recent phase II study, a short course of EPOCH with dose-dense rituximab (SC-EPOCH-RR) was shown to be effective resulting in a CR rate of 91% in patients with HIV-associated DLBCL (n = 33). At a median follow up of 5 years, the PFS and OS rates were 84% and 68%, respectively.⁴¹ In this study, the addition of rituximab did not appear to cause serious infection-related complications or deaths. The safety and efficacy of SC-EPOCH-RR in patients with low-risk HIV-associated BL was demonstrated in a subsequent prospective study that included 11 patients with HIV-associated BL.⁴² At a median follow-up of 86 months, the rates of freedom from progression and OS were 100% and 90%, respectively for patients treated with SC-EPOCH-RR. This regimen was also associated with lower incidences of fever and neutropenia than DA-EPOCH-R.

The safety and preliminary efficacy of vorinostat in combination with EPOCH-R regimen in patients with high-risk HIV-associated NHL (with at least one of the following high-risk characteristics: age-adjusted IPI 2–3, Ki-67 80%, non-germinal center B-cell (GCB) DLBCL, non-Burkitt B-cell NHL) was established in a phase I study and the utility of adding vorinostat to EPOCH-R regimen is being evaluated in a phase II randomized trial.⁴³ The result of a retrospective study showed that the addition of bortezomib

to EPOCH is also safe and effective as a front-line treatment for HIV-associated PBL but this has to be confirmed in a prospective study.⁴⁴

CHOP (cyclophosphamide, doxorubicin, vincristine and prednisone with or without rituximab)

CHOP regimen has been shown to induce CR rates of 30% to 48%, with a median OS of approximately 25 months and the addition of rituximab has been associated with improved CR rates in patients with HIV-associated lymphomas.^{45–49}

In the randomized phase III trial (AMC 010 study) of 150 patients with HIV-associated NHL (80% DLBCL; 9% BL), the addition of rituximab to CHOP (RCHOP) was associated with improved CR rates (58% vs. 47%) as well as longer median time to progression (29 months vs. 20 months) and longer median OS (32 months vs. 25 months) compared with CHOP alone.⁴⁶ The median PFS was similar between treatment groups (10 months vs. 9 months). It should also be noted that in this study, 35 patients randomized to the RCHOP arm had received maintenance rituximab following initial RCHOP.⁴⁶ However, clinical trials evaluating maintenance rituximab in non-HIV patients did not show survival benefit. In addition, in the aforementioned clinical trial, all deaths due to infection occurred during the maintenance phase.⁴⁶ Therefore, the use of maintenance rituximab is not recommended for patients with HIV-associated lymphomas. In subsequent phase II trials, 6 cycles of the RCHOP resulted in CR/CRu rates of 69% to 77%, with 2-year OS and 3-year OS rates of 75% and 56%, respectively, in patients with HIV-associated NHL (majority with DLBCL), with manageable toxicities.^{48,49}

Liposomal doxorubicin or pegylated liposomal doxorubicin in combination with cyclophosphamide, vincristine and prednisone (CDOP) has also been shown to be effective in patients with HIV-associated lymphoma.^{50,51} In a multicenter phase II trial (AMC 047 study) of 40



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

patients with HIV-associated NHL (DLBCL in 98% of cases), infusional CDOP (with pegylated liposomal doxorubicin) in combination with rituximab (RCDOP), resulted in an ORR of 68% (48% CR). The 1-year PFS and OS rates were 61% and 70%, respectively; the 2-year PFS and OS were 52% and 62%, respectively. Infectious complications were reported in 40% of patients (grade 4 in 5%) but no infection-related deaths occurred. This may in part be explained by the fact that patients received concomitant ART and those with low CD4 counts ($\leq 100/\text{mCL}$ at baseline or during anti-tumor therapy) received antimicrobial prophylaxis. Factors such as decreased CD4 counts or increased HIV viral load did not appear to influence treatment response. However, these results, however, appeared less favorable compared with the EPOCH regimen.

CODOX-M/IVAC regimen (cyclophosphamide, doxorubicin and high-dose methotrexate, alternating with ifosfamide, etoposide and high-dose cytarabine) with or without rituximab

The CODOX-M/IVAC regimen (cyclophosphamide, doxorubicin and high-dose methotrexate, alternating with ifosfamide, etoposide and high-dose cytarabine) with or without rituximab, commonly used in the management of patients with BL is also effective in patients with HIV-associated BL.⁵²⁻⁵⁶

In one retrospective study that evaluated CODOX-M/IVAC with or without rituximab in 80 patients with BL, the CR rates (93% and 88%, respectively) 3-year PFS (68% for both subgroups) and 3-year OS rates (68% and 72%, respectively) were similar among patients with and without HIV infection. There was a trend toward improved 3-year PFS rate (74% vs. 61%) and OS rate (77% vs. 66%) with the addition of rituximab.⁵³ The AMC 048 trial prospectively evaluated modified CODOX-M/IVAC with rituximab in 34 patients with HIV-associated BL (2 patients had low-risk disease; 32 patients had high-risk disease).⁵⁶ Patients with low-risk disease were treated with 3 cycles of rituximab and CODOX-M (R-CODOX-M) whereas all other patients with high-risk

disease were treated with R-CODOX-M/IVAC. The median follow-up was 26 months. The 1-year PFS and OS rates were 69% and 72%, respectively; the 2-year OS rate was 69%.

HyperCVAD regimen (hyperfractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone, alternating with high-dose methotrexate and cytarabine)

The hyperCVAD regimen (hyperfractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone, alternating with high-dose methotrexate and cytarabine) with or without rituximab has demonstrated efficacy in the treatment of *de novo* BL/leukemia and incorporation of rituximab was associated with improved disease-related outcomes particularly for the older subset of patients.^{57,58} HyperCVAD regimen given in combination with ART has also been shown to be effective in patients with HIV-associated BL/leukemia and Burkitt-like lymphoma resulting in CR rates of 64–92% and a median OS of 12 months.⁵⁹

NCCN Recommendations

The NCCN Guidelines recommend the use of ART and growth factor support along with full-dose chemotherapy with or without rituximab. Patients on ART with persistently low CD4+ count of less than 50 to 100/mCL tend to have a poorer prognosis, higher risk of infection and cytopenias.^{38,46,60} Therefore, omission of rituximab is strongly suggested for these patients due to the higher risk of serious infectious complications. Maximizing supportive care and close monitoring for cytopenias and infections is recommended while administering lymphoma therapy for this groups of patients.

ART can be administered safely with chemotherapy. However, certain antiviral drugs can interfere with the metabolism of cancer therapies, commonly by CYP3A4 inhibition (protease inhibitors) or CYP3A4 induction (non-nucleoside reverse transcriptase inhibitors).^{61,62} Therefore, effective alternatives for the existing ART should be



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

considered to minimize toxicities and drug-drug interactions when anticipated. In general, avoidance of zidovudine, cobicistat, and ritonavir is strongly recommended. Any change in antiviral therapy should be made in consultation with an infectious disease specialist.

CODOX-M/IVAC (with or without rituximab), EPOCH-R and hyperCVAD with rituximab are recommended for patients with HIV-associated BL. EPOCH or CHOP in combination with rituximab is recommended for patients with HIV-associated DLBCL. EPOCH-R is included as the preferred regimen for the treatment of HIV-associated BL and DLBCL. CNS prophylaxis with intrathecal methotrexate or intrathecal cytarabine is indicated for patients with HIV-associated DLBCL with selected high-risk features (e.g., double-hit lymphoma, involvement of 2 or more extranodal sites with elevated LDH, bone marrow involvement, or other high-risk site involvement such as epidural, testicular or paranasal sinuses).

Patients with lymphoma associated with HHV8-positive DLBCL, NOS and PEL can also be treated with the same regimens as described for patients with HIV-associated DLBCL. Since most cases of PEL are CD20-negative, the addition of rituximab to the chemotherapy is not indicated.

CODOX-M/IVAC, EPOCH or hyper-CVAD are recommended for patients with PBL, with the realization that only limited data are available on the management of these patients at this time.^{18,21,63} A CD4 count of $>100/\mu\text{L}$ at diagnosis, chemosensitive disease and CR to first-line chemotherapy were associated with better outcomes.^{21,63} Consolidation with high-dose therapy followed by autologous stem cell rescue (HDT/ASCR) can be considered following CR after initial therapy for patients with high-risk features (age-adjusted IPI >2 , presence of *MYC* gene rearrangement or *TP53* deletion).^{64,65}

Immune reconstitution with ART along with the use of high-dose methotrexate should be considered for patients with PCNSL.¹²⁻¹⁵ RT should be reserved for patients who are not able to tolerate systemic therapy or for those with disease that is refractory to systemic therapy. Selected patients with good performance status receiving ART may also be treated as per the NCCN Guidelines for Primary CNS Lymphoma.

Relapsed/Refractory Disease

In a recent multicenter study that evaluated the risk factors and incidence of relapse in a large cohort of patients with HIV-associated lymphomas (after achieving CR to first-line treatment), unclassifiable histology, advanced stage disease, no concomitant ART during chemotherapy and the use of R-CHOP-based regimens were independently associated with higher risk of relapse in patients with BL.³⁶

Bortezomib in combination with ICE (ifosfamide, carboplatin and etoposide) + rituximab is an effective second-line therapy for patients with relapsed/refractory HIV-associated lymphomas.⁶⁶ Brentuximab vedotin has also been shown to be effective in patients with PEL (albeit in case reports).⁶⁷ Additional prospective studies are warranted to confirm these findings.

HDT/ASCR is associated with favorable survival outcome in patients with chemosensitive relapsed/refractory disease, similar to the HIV-seronegative population.⁶⁸⁻⁷⁴

In a retrospective analysis that evaluated the outcomes of 88 patients with relapsed or refractory HIV-associated lymphomas treated with curative intent at 13 AMC sites, ICE (39%), DA-EPOCH (19%) and ESHAP (etoposide, methylprednisone, cytarabine and cisplatin, 13%) were the most commonly used second-line regimens.⁷¹ The ORR was 31% and the 1-year OS rate was 37% for the entire study population. Baseline CD4 counts did not influence OS outcomes. Subsequent



treatment with autologous hematopoietic cell transplant (HCT) was associated with improved 1-year OS (63% vs. 37%) compared with no HCT. There was no difference in 1-year OS rate based on HCT (88% with HCT vs. 82% with no transplant) for patients who experienced a response (CR or PR) after second-line therapy. The response rate and survival outcomes were better for patients with non-BL histology than that of patients with BL histology. The ORR and 1-year OS rate were 33% and 42% respectively for patients with non-BL histology compared to 17% and 12% respectively for those with BL histology. Patients with primary refractory disease (n=54) had significantly decreased ORR (24% vs. 56%; $P=.003$) and decreased 1-year OS (31% vs. 59%; $P=.022$) compared with those with relapsed disease.

In a phase II study of 43 patients with chemosensitive, relapsed or refractory HIV-associated lymphoma, at a median follow-up of 25 months, autologous HCT was associated with an estimated 1-year and 2-year OS rates of 87% and 82%, respectively.⁷² The estimated 2-year PFS rate and the 1-year transplant related mortality rates were 80% and 5%, respectively. A matched population of control patients from the CIBMTR data registry revealed that these outcomes did not differ statistically from that of non-HIV-infected patients.

In a more recent retrospective study that evaluated the outcome of patients with HIV-associated lymphomas (n = 118; DLBCL, 47%, HL, 24%, BL, 18%, and PEL, 7%) following autologous HCT in the rituximab and ART era, at a median follow-up of 4 years, the 3-year non-relapse mortality, incidence of relapse, PFS and OS rates were 10%, 27%, 63% and 66%, respectively.⁷⁴ In the multivariate analysis, disease status less than PR at the time of transplant was a significant predictor of unfavorable PFS and OS.

These results suggest that HDT/ASCR should be considered for patients with chemosensitive relapsed or refractory HIV-associated lymphomas if they are candidates for transplant.



References

1. Rihana N, Nanjappa S, Sullivan C, et al. Malignancy Trends in HIV-Infected Patients Over the Past 10 Years in a Single-Center Retrospective Observational Study in the United States. *Cancer Control* 2018;25:1073274818797955. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30185062>.
2. Shiels MS, Islam JY, Rosenberg PS, et al. Projected Cancer Incidence Rates and Burden of Incident Cancer Cases in HIV-Infected Adults in the United States Through 2030. *Ann Intern Med* 2018;168:866-873. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29801099>.
3. Gopal S, Patel MR, Yanik EL, et al. Temporal trends in presentation and survival for HIV-associated lymphoma in the antiretroviral therapy era. *J Natl Cancer Inst* 2013;105:1221-1229. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23892362>.
4. U.S. National Library of Medicine Key MEDLINE® Indicators Available at: http://www.nlm.nih.gov/bsd/bsd_key.html.
5. Levine AM, Seneviratne L, Espina BM, et al. Evolving characteristics of AIDS-related lymphoma. *Blood* 2000;96:4084-4090. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11110677>.
6. Besson C, Goubar A, Gabarre J, et al. Changes in AIDS-related lymphoma since the era of highly active antiretroviral therapy. *Blood* 2001;98:2339-2344. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11588028>.
7. Han X, Jemal A, Hulland E, et al. HIV Infection and Survival of Lymphoma Patients in the Era of Highly Active Antiretroviral Therapy. *Cancer Epidemiol Biomarkers Prev* 2017;26:303-311. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27756777>.
8. Collaboration of Observational HIV Epidemiological Research Europe (COHERE) study group. Prognosis of HIV-associated non-hodgkin lymphoma in patients starting combination antiretroviral therapy. *AIDS* 2009;23:2029-2037. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19531926>.
9. Cingolani A, Cozzi Lepri A, Teofili L, et al. Survival and predictors of death in people with HIV-associated lymphoma compared to those with a diagnosis of lymphoma in general population. *PLoS One* 2017;12:e0186549. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29088223>.
10. Gopal S, Patel MR, Achenbach CJ, et al. Lymphoma immune reconstitution inflammatory syndrome in the center for AIDS research network of integrated clinical systems cohort. *Clin Infect Dis* 2014;59:279-286. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24755860>.
11. Besson C, Lancar R, Prevot S, et al. Outcomes for HIV-associated diffuse large B-cell lymphoma in the modern combined antiretroviral therapy era. *AIDS* 2017;31:2493-2501. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28926410>.
12. Newell ME, Hoy JF, Cooper SG, et al. Human immunodeficiency virus-related primary central nervous system lymphoma. *Cancer* 2004;100:2627-2636. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15197806>.
13. Diamond C, Taylor TH, Im T, et al. Highly active antiretroviral therapy is associated with improved survival among patients with AIDS-related primary central nervous system non-Hodgkin's lymphoma. *Curr HIV Res* 2006;4:375-378. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16842088>.
14. Gupta NK, Nolan A, Omuro A, et al. Long-term survival in AIDS-related primary central nervous system lymphoma. *Neuro Oncol* 2017;19:99-108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27576871>.
15. Moulignier A, Lamirel C, Picard H, et al. Long-term AIDS-related PCNSL outcomes with HD-MTX and combined antiretroviral therapy. *Neurology* 2017;89:796-804. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28747447>.
16. Dong HY, Scadden DT, de Leval L, et al. Plasmablastic lymphoma in HIV-positive patients: an aggressive Epstein-Barr virus-associated



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

extramedullary plasmacytic neoplasm. Am J Surg Pathol 2005;29:1633-1641. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16327436>.

17. Rodrigues-Fernandes CI, de Souza LL, Santos-Costa SFD, et al. Clinicopathological analysis of oral plasmablastic lymphoma: A systematic review. J Oral Pathol Med 2018;47:915-922. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29917262>.

18. Teruya-Feldstein J, Chiao E, Filippa DA, et al. CD20-negative large-cell lymphoma with plasmablastic features: a clinically heterogeneous spectrum in both HIV-positive and -negative patients. Ann Oncol 2004;15:1673-1679. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15520070>.

19. Loghavi S, Alayed K, Aladily TN, et al. Stage, age, and EBV status impact outcomes of plasmablastic lymphoma patients: a clinicopathologic analysis of 61 patients. J Hematol Oncol 2015;8:65. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26055271>.

20. Noy A, Lensing SY, Moore PC, et al. Plasmablastic lymphoma is treatable in the HAART era. A 10 year retrospective by the AIDS Malignancy Consortium. Leuk Lymphoma 2016;57:1731-1734. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26674561>.

21. Castillo JJ, Winer ES, Stachurski D, et al. Prognostic factors in chemotherapy-treated patients with HIV-associated Plasmablastic lymphoma. Oncologist 2010;15:293-299. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/20167839>.

22. Castillo JJ, Furman M, Beltran BE, et al. Human immunodeficiency virus-associated plasmablastic lymphoma: poor prognosis in the era of highly active antiretroviral therapy. Cancer 2012;118:5270-5277. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22510767>.

23. Schommers P, Hentrich M, Hoffmann C, et al. Survival of AIDS-related diffuse large B-cell lymphoma, Burkitt lymphoma, and plasmablastic lymphoma in the German HIV Lymphoma Cohort. Br J Haematol

2015;168:806-810. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25403997>.

24. Schommers P, Wyen C, Hentrich M, et al. Poor outcome of HIV-infected patients with plasmablastic lymphoma: results from the German AIDS-related lymphoma cohort study. AIDS 2013;27:842-845. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23574794>.

25. Nador RG, Cesarman E, Chadburn A, et al. Primary effusion lymphoma: a distinct clinicopathologic entity associated with the Kaposi's sarcoma-associated herpes virus. Blood 1996;88:645-656. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/8695812>.

26. Shimada K, Hayakawa F, Kiyoi H. Biology and management of primary effusion lymphoma. Blood 2018;132:1879-1888. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30154110>.

27. Olszewski AJ, Fallah J, Castillo JJ. Human immunodeficiency virus-associated lymphomas in the antiretroviral therapy era: Analysis of the National Cancer Data Base. Cancer 2016;122:2689-2697. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27337679>.

28. Boulanger E, Gerard L, Gabarre J, et al. Prognostic factors and outcome of human herpesvirus 8-associated primary effusion lymphoma in patients with AIDS. J Clin Oncol 2005;23:4372-4380. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15994147>.

29. El-Fattah MA. Clinical characteristics and survival outcome of primary effusion lymphoma: A review of 105 patients. Hematol Oncol 2017;35:878-883. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27859456>.

30. Raphael MM, Audouin J, Lamine M, et al. Immunophenotypic and genotypic analysis of acquired immunodeficiency syndrome-related non-Hodgkin's lymphomas. Correlation with histologic features in 36 cases. French Study Group of Pathology for HIV-Associated Tumors. Am J Clin Pathol 1994;101:773-782. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/8209868>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

31. Arvey A, Ojesina AI, Pedomallu CS, et al. The tumor virus landscape of AIDS-related lymphomas. *Blood* 2015;125:e14-22. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25827832>.

32. Cheung MC, Pantanowitz L, Dezube BJ. AIDS-related malignancies: emerging challenges in the era of highly active antiretroviral therapy. *Oncologist* 2005;10:412-426. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15967835>.

33. Mounier N, Spina M, Gabarre J, et al. AIDS-related non-Hodgkin lymphoma: final analysis of 485 patients treated with risk-adapted intensive chemotherapy. *Blood* 2006;107:3832-3840. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16410446>.

34. Mounier N, Spina M, Gisselbrecht C. Modern management of non-Hodgkin lymphoma in HIV-infected patients. *Br J Haematol* 2007;136:685-698. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17229246>.

35. Barta SK, Xue X, Wang D, et al. Treatment factors affecting outcomes in HIV-associated non-Hodgkin lymphomas: a pooled analysis of 1546 patients. *Blood* 2013;122:3251-3262. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24014242>.

36. Schommers P, Gillor D, Hentrich M, et al. Incidence and risk factors for relapses in HIV-associated non-Hodgkin lymphoma as observed in the German HIV-related lymphoma cohort study. *Haematologica* 2018;103:857-864. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29439188>.

37. Tan CRC, Barta SK, Lee J, et al. Combination antiretroviral therapy accelerates immune recovery in patients with HIV-related lymphoma treated with EPOCH: a comparison within one prospective trial AMC034. *Leuk Lymphoma* 2018;59:1851-1860. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29160731>.

38. Sparano JA, Lee JY, Kaplan LD, et al. Rituximab plus concurrent infusional EPOCH chemotherapy is highly effective in HIV-associated

B-cell non-Hodgkin lymphoma. *Blood* 2010;115:3008-3016. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20023215>.

39. Barta SK, Lee JY, Kaplan LD, et al. Pooled analysis of AIDS malignancy consortium trials evaluating rituximab plus CHOP or infusional EPOCH chemotherapy in HIV-associated non-Hodgkin lymphoma. *Cancer* 2012;118:3977-3983. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22180164>.

40. Little RF, Pittaluga S, Grant N, et al. Highly effective treatment of acquired immunodeficiency syndrome-related lymphoma with dose-adjusted EPOCH: impact of antiretroviral therapy suspension and tumor biology. *Blood* 2003;101:4653-4659. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12609827>.

41. Dunleavy K, Little RF, Pittaluga S, et al. The role of tumor histogenesis, FDG-PET, and short-course EPOCH with dose-dense rituximab (SC-EPOCH-RR) in HIV-associated diffuse large B-cell lymphoma. *Blood* 2010;115:3017-3024. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20130244>.

42. Dunleavy K, Pittaluga S, Shovlin M, et al. Low-intensity therapy in adults with Burkitt's lymphoma. *N Engl J Med* 2013;369:1915-1925. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24224624>.

43. Ramos JC, Sparano JA, Rudek MA, et al. Safety and Preliminary Efficacy of Vorinostat With R-EPOCH in High-risk HIV-associated Non-Hodgkin's Lymphoma (AMC-075). *Clin Lymphoma Myeloma Leuk* 2018;18:180-190 e182. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29426719>.

44. Castillo JJ, Guerrero-Garcia T, Baldini F, et al. Bortezomib plus EPOCH is effective as frontline treatment in patients with plasmablastic lymphoma. *Br J Haematol* 2019;184:679-682. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29527667>.

45. Ratner L, Lee J, Tang S, et al. Chemotherapy for human immunodeficiency virus-associated non-Hodgkin's lymphoma in combination with highly active antiretroviral therapy. *J Clin Oncol*



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

2001;19:2171-2178. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11304769>.

46. Kaplan LD, Lee JY, Ambinder RF, et al. Rituximab does not improve clinical outcome in a randomized phase 3 trial of CHOP with or without rituximab in patients with HIV-associated non-Hodgkin lymphoma: AIDS-Malignancies Consortium Trial 010. *Blood* 2005;106:1538-1543.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15914552>.

47. Weiss R, Mitrou P, Arasteh K, et al. Acquired immunodeficiency syndrome-related lymphoma: simultaneous treatment with combined cyclophosphamide, doxorubicin, vincristine, and prednisone chemotherapy and highly active antiretroviral therapy is safe and improves survival--results of the German Multicenter Trial. *Cancer* 2006;106:1560-1568. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16502436>.

48. Boue F, Gabarre J, Gisselbrecht C, et al. Phase II trial of CHOP plus rituximab in patients with HIV-associated non-Hodgkin's lymphoma. *J Clin Oncol* 2006;24:4123-4128. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16896005>.

49. Ribera JM, Oriol A, Morgades M, et al. Safety and efficacy of cyclophosphamide, adriamycin, vincristine, prednisone and rituximab in patients with human immunodeficiency virus-associated diffuse large B-cell lymphoma: results of a phase II trial. *Br J Haematol* 2008;140:411-419. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18162120>.

50. Levine AM, Tulpule A, Espina B, et al. Liposome-encapsulated doxorubicin in combination with standard agents (cyclophosphamide, vincristine, prednisone) in patients with newly diagnosed AIDS-related non-Hodgkin's lymphoma: results of therapy and correlates of response. *J Clin Oncol* 2004;22:2662-2670. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15226333>.

51. Levine AM, Noy A, Lee JY, et al. Pegylated liposomal doxorubicin, rituximab, cyclophosphamide, vincristine, and prednisone in AIDS-related lymphoma: AIDS Malignancy Consortium Study 047. *J Clin Oncol*

2013;31:58-64. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23169503>.

52. Wang ES, Straus DJ, Teruya-Feldstein J, et al. Intensive chemotherapy with cyclophosphamide, doxorubicin, high-dose methotrexate/ifosfamide, etoposide, and high-dose cytarabine (CODOX-M/IVAC) for human immunodeficiency virus-associated Burkitt lymphoma. *Cancer* 2003;98:1196-1205. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12973843>.

53. Barnes JA, Lacasce AS, Feng Y, et al. Evaluation of the addition of rituximab to CODOX-M/IVAC for Burkitt's lymphoma: a retrospective analysis. *Ann Oncol* 2011;22:1859-1864. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21339382>.

54. Rodrigo JA, Hicks LK, Cheung MC, et al. HIV-Associated Burkitt Lymphoma: Good Efficacy and Tolerance of Intensive Chemotherapy Including CODOX-M/IVAC with or without Rituximab in the HAART Era. *Adv Hematol* 2012;2012:735392. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22190945>.

55. Alwan F, He A, Montoto S, et al. Adding rituximab to CODOX-M/IVAC chemotherapy in the treatment of HIV-associated Burkitt lymphoma is safe when used with concurrent combination antiretroviral therapy. *AIDS* 2015;29:903-910. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25730506>.

56. Noy A, Lee JY, Cesarman E, et al. AMC 048: modified CODOX-M/IVAC-rituximab is safe and effective for HIV-associated Burkitt lymphoma. *Blood* 2015;126:160-166. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25957391>.

57. Thomas DA, Faderl S, O'Brien S, et al. Chemoimmunotherapy with hyper-CVAD plus rituximab for the treatment of adult Burkitt and Burkitt-type lymphoma or acute lymphoblastic leukemia. *Cancer* 2006;106:1569-1580. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16502413>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

58. Thomas DA, Kantarjian HM, Faderl S, et al. Hyper-CVAD and Rituximab for De Novo Burkitt Lymphoma/Leukemia [abstract]. Blood 2011;118:Abstract 2698. Available at: <http://www.bloodjournal.org/content/118/21/2698>.

59. Cortes J, Thomas D, Rios A, et al. Hyperfractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone and highly active antiretroviral therapy for patients with acquired immunodeficiency syndrome-related Burkitt lymphoma/leukemia. Cancer 2002;94:1492-1499. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11920506>.

60. Spina M, Jaeger U, Sparano JA, et al. Rituximab plus infusional cyclophosphamide, doxorubicin, and etoposide in HIV-associated non-Hodgkin lymphoma: pooled results from 3 phase 2 trials. Blood 2005;105:1891-1897. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15550484>.

61. Foca E, Cavaglia G, Rusconi S, et al. Survival in HIV-infected patients with lymphoma according to the choice of antiretroviral treatment: an observational multicentre study. HIV Med 2018. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29862615>.

62. Sombogaard F, Franssen EJF, Terpstra WE, et al. Outcome effects of antiretroviral drug combinations in HIV-positive patients with chemotherapy for lymphoma: a retrospective analysis. Int J Clin Pharm 2018;40:1402-1408. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29948741>.

63. Koizumi Y, Uehira T, Ota Y, et al. Clinical and pathological aspects of human immunodeficiency virus-associated plasmablastic lymphoma: analysis of 24 cases. Int J Hematol 2016;104:669-681. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27604616>.

64. Al-Malki MM, Castillo JJ, Sloan JM, Re A. Hematopoietic cell transplantation for plasmablastic lymphoma: a review. Biol Blood Marrow Transplant 2014;20:1877-1884. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24946718>.

65. Cattaneo C, Finel H, McQuaker G, et al. Autologous hematopoietic stem cell transplantation for plasmablastic lymphoma: the European Society for Blood and Marrow Transplantation experience. Biol Blood Marrow Transplant 2015;21:1146-1147. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25783635>.

66. Reid EG, Looney D, Maldarelli F, et al. Safety and efficacy of an oncolytic viral strategy using bortezomib with ICE/R in relapsed/refractory HIV-positive lymphomas. Blood Adv 2018;2:3618-3626. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30573564>.

67. Chang VA, Wang HY, Reid EG. Activity of brentuximab vedotin in AIDS-related primary effusion lymphoma. Blood Adv 2019;3:766-768. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30837213>.

68. Serrano D, Carrion R, Balsalobre P, et al. HIV-associated lymphoma successfully treated with peripheral blood stem cell transplantation. Exp Hematol 2005;33:487-494. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15781340>.

69. Spitzer TR, Ambinder RF, Lee JY, et al. Dose-reduced busulfan, cyclophosphamide, and autologous stem cell transplantation for human immunodeficiency virus-associated lymphoma: AIDS Malignancy Consortium study 020. Biol Blood Marrow Transplant 2008;14:59-66. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18158962>.

70. Re A, Michieli M, Casari S, et al. High-dose therapy and autologous peripheral blood stem cell transplantation as salvage treatment for AIDS-related lymphoma: long-term results of the Italian Cooperative Group on AIDS and Tumors (GICAT) study with analysis of prognostic factors. Blood 2009;114:1306-1313. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19451551>.

71. Bayraktar UD, Ramos JC, Petrich A, et al. Outcome of patients with relapsed/refractory acquired immune deficiency syndrome-related lymphoma diagnosed 1999-2008 and treated with curative intent in the AIDS Malignancy Consortium. Leuk Lymphoma 2012;53:2383-2389. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22642936>.



72. Alvarnas JC, Le Rademacher J, Wang Y, et al. Autologous hematopoietic cell transplantation for HIV-related lymphoma: results of the BMT CTN 0803/AMC 071 trial. Blood 2016;128:1050-1058. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27297790>.

73. Ayala E, Chavez JC, Gomez A, et al. Feasibility and Efficacy of High-Dose Chemotherapy and Autologous Hematopoietic Cell Transplantation for HIV-Associated Lymphoma: A Single-Institution Experience. Clin Lymphoma Myeloma Leuk 2018;18:548-551. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29937399>.

74. Hubel K, Re A, Boumendil A, et al. Autologous stem cell transplantation for HIV-associated lymphoma in the antiretroviral and rituximab era: a retrospective study by the EBMT Lymphoma Working Party. Bone Marrow Transplant 2019. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30804486>.

Discussion
update in
progress



Post-Transplant Lymphoproliferative Disorders

Post-transplant lymphoproliferative disorders (PTLD) are a heterogeneous group of lymphomas that occur after solid organ transplant (SOT) or allogeneic hematopoietic cell transplant (HCT) that are related to immunosuppression and the Epstein-Barr virus (EBV).¹ PTLD following SOT are of recipient origin in the majority of cases, often involving the grafted organ, whereas PTLD following allogeneic HCT are usually of donor origin.²⁻⁹

The incidence of PTLD following SOT varies significantly depending on the transplanted organ (kidney transplants, 0.8% to 3%; pancreatic transplants, 0.5% to 5%; liver transplants, 1% to 6%; heart transplants, 2.0% to 8%; lung transplants, 3% to 10%; and multiorgan and intestinal transplants, ≤20%).¹ The incidence of PTLD following allogeneic HCT varies depending on the degree of human leucocyte antigen (HLA) matching and the need for T-cell depletion protocol prior to transplantation.^{4,5,9} Thus, the incidence of PTLD is the highest following haploidentical allogeneic HCT especially in cases of selective T-cell depletion (>20%) followed by cases of unrelated donors (4% to 10%), umbilical-cord transplants (4% to 5%), and matched, related donors (1% to 3%).

About 50% of PTLD following SOT are considered late-onset PTLD (diagnosed >1 year after transplant) and are more likely EBV-negative.¹⁰⁻¹⁵ Gene expression profiling studies have also shown that EBV-negative PTLD are clinically and biologically distinct from EBV-positive PTLD.¹⁶⁻²⁰ EBV-negative PTLD are more likely to be of germinal center B-cell (GCB) type and EBV-positive PTLD are usually of non-GCB type.^{17,21}

Literature Search Criteria and Guidelines Update Methodology

Prior to the update of this version of the NCCN Guidelines® for B-Cell Lymphomas an electronic search of the PubMed database was performed

to obtain key literature in PTLD published since the previous Guidelines update. The PubMed database was chosen as it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.²²

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase II; Clinical Trial, Phase III; Clinical Trial, Phase IV; Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.

The data from key PubMed articles deemed as relevant to these Guidelines and discussed by the panel have been included in this version of the Discussion section (eg, e-publications ahead of print, meeting abstracts). Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.

The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

Risk factors for developing PTLD

EBV serology mismatch (recipient EBV-negative and donor EBV-positive), type of transplanted organ (highest risks for multiorgan, bowel, lung, heart/lung transplants), intensity of induction immunosuppression, and the type of immunosuppression are considered as established risk factors for developing PTLD.²³⁻³⁰ The risk is higher among children compared with adults, because primary EBV infection in EBV-negative organ recipients is the most common driver in children.³⁰⁻³⁷

Unrelated or HLA-mismatched allografts, the use of anti-thymocyte globulin (ATG) or anti-CD3 monoclonal antibody (MAB) for the prevention or treatment of graft-versus-host disease (GVHD), and T-cell depletion of the allograft are associated with increased risks for PTLD in patients



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

undergoing allogeneic HCT.^{4,5,34} The use of ATG or anti-CD3 MAB (OKT3), calcineurin inhibition with tacrolimus as primary immunosuppressive therapy along with the use of azathioprine and new agents (eg. belatacept in EBV-negative transplant recipients) are associated with increased risks for developing PTLD following SOT.^{25,29} In an analysis of 523 patients who underwent heart transplant, switching from calcineurin inhibitor-based immunosuppression to sirolimus-based immunosuppression was associated with decreased rates of malignancies following heart transplant.³⁸

Risk factors for surviving PTLD

Older age, poor performance status (Eastern Cooperative Oncology performance score of ≥ 2), elevated lactate dehydrogenase (LDH), organ dysfunction, multiple involved lymph nodes, multi organ involvement, graft organ involvement, central nervous system (CNS) involvement, number of extranodal sites (one versus more than one), the type of organ transplanted, hypoalbuminemia, the International Prognostic Factor Index (IPI), and comorbidities have been identified as prognostic factors for poor survival in patients with PTLD following SOT.³⁹⁻⁴³ In the PTLD-1 trial, the IPI, lung transplants, and inadequate response to rituximab induction therapy were associated with a worse prognosis.⁴⁴

Classification

In the 2017 WHO classification, PTLD are classified into 6 subclasses: 3 early lesions (nondestructive PTLD), monomorphic PTLD (B-cell, T-cell and NK cell type), polymorphic PTLD, and classical Hodgkin lymphoma (CHL) PTLD.⁴⁵

Early lesions consist of 3 histologic subtypes: plasmacytic hyperplasia PTLD, infectious mononucleosis PTLD, and florid follicular hyperplasia PTLD. Early lesions typically develop within a year of transplantation and are EBV-positive in almost all cases.⁴⁶

Monomorphic PTLD appear to be the most common subtype of PTLD and the majority are of B-cell origin with diffuse large B cell lymphoma (DLBCL) being the most frequent subtype.^{21,37} Although uncommon, Burkitt lymphoma (BL),⁴⁷⁻⁵⁰ plasma cell myeloma or plasmacytoma⁵¹⁻⁵⁵ have also been reported. Indolent B-cell lymphomas arising in transplant recipients are not included among PTLD, with the exception of EBV-positive marginal zone lymphomas. Monomorphic PTLD of T-cell or natural killer (NK)-cell origin (although very rare) tend to occur later (after a median of 6 years following transplant in one series).^{56,57} Peripheral T-cell lymphoma, not-otherwise specified (PTCL, NOS) is the most prevalent subtype followed by anaplastic large cell lymphoma (ALCL) and hepatosplenic T-cell lymphoma (HSTCL).⁵⁷ From a pathology perspective, monomorphic PTLD cannot be distinguished from lymphomas with a similar lineage and cell of origin in immunocompetent patients suggesting that the subclassification of these types should be the same.

Polymorphic PTLD are mostly EBV positive, and can be either polyclonal or monoclonal. These are the most common type of PTLD among children and are characterized by a mixed lymphoproliferation consisting of immunoblasts, plasma cells, and intermediate-sized lymphoid cells. Immunohistochemistry (IHC) will show a variable mixtures of B-cells and T-cells. However, subdividing polymorphic PTLD is not indicated in the WHO classification since this does not reliably predict clinical behavior.⁴⁶

CHL PTLD is almost always EBV-positive, and is the least common of the four PTLD categories.⁵⁸

Diagnosis

The clinical presentation of PTLD is heterogeneous and is characterized by a high incidence of extranodal disease which may involve the gastrointestinal tract in (20% to 30%), solid allografts (10% to 15%), and CNS (5% to 20%). The diagnosis of PTLD can be challenging given the



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

nonspecific clinical presentation, and heterogeneity in histopathologic and immunophenotypic presentations. Histopathology and adequate immunophenotyping are essential to confirm the diagnosis of PTLD.^{59,60} Among B-cell PTLD, expression of BCL6, MUM1 and CD138 can be useful in distinguishing between the histological subtypes of PTLD.^{61,62} BCL6 expression was detected in most cases of monomorphic PTLD (71% of centroblastic DLBCL), whereas it was consistently absent in polymorphic PTLD. MUM1 was preferentially expressed in 92% of polymorphic PTLD.⁶¹ Overall, BCL6–, MUM1+ and CD138– phenotype is associated most frequently with polymorphic PTLD; BCL6+, MUM1+/- and CD138– is mostly associated with monomorphic PTLD.^{61,62}

The recommended IHC panel includes the following markers: CD3, CD5, CD10, BCL6, BCL2, IRF4/MUM1, CD20, CD79a, PAX5, Ki67, and kappa, lambda light chains. Cell surface markers CD3, CD5, CD7, CD4, CD8, CD19, CD20, CD10, and kappa/lambda are recommended for flow cytometric analysis. Under certain circumstances, the following additional markers may be useful for an IHC panel: CD15, CD30, CD45, CD7, CD4, CD8, ALK, TIA-1, granzyme B, CD57, CD56, and CD138. In addition, the following markers for flow cytometry may also be useful under certain situations: CD138, CD30, CD57, CD56, CD16, CD25, CD52, and cytoplasmic kappa or lambda.

Up to 50% of PTLD cases that develop after SOT are not associated with EBV.¹³ Although an association with EBV is not required for the diagnosis of PTLD, evaluation of EBV infection status is an essential component of the diagnostic workup. EBV can be detected by either IHC for latent membrane protein 1 (LMP 1) or EBV encoded RNA in situ hybridization (EBER-ISH). EBER-ISH is more sensitive than IHC, and is recommended if EBV-LMP-1 is negative. If immunostaining for EBV-LMP 1 is positive, EBER-ISH is not required. Under certain circumstances, EBV evaluation by Southern blot may also be useful.

Immunoglobulin heavy chain variable (*IGHV*) gene mutations are seen in the majority of B-cell PTLD, with the exception of early lesions.^{46,62,63}

Genetic alterations in *MYC*, *NRAS* and *TP53* are seen only in monomorphic PTLD and *BCL6* mutations (present in 43% of the polymorphic PTLD) have been associated with shorter survival and poor response to therapy.^{46,64,65} In certain situations, molecular analysis to detect *IGHV* rearrangements and *BCL6* mutations could be useful.

Workup

The initial workup for PTLD should include a physical examination and evaluation of performance status. Laboratory assessments should include standard blood work including complete blood count (CBC) with differential and a metabolic panel (to include albumin, electrolytes, BUN, and creatinine), in addition to measurements of serum LDH levels. Bone marrow evaluations may be useful in selected cases. The history of immunosuppression treatment with the transplant course must be assessed. Chest/abdomen/pelvis CT with contrast of diagnostic quality and/or whole-body PET/CT scan are recommended as part of initial diagnostic workup. Brain MRI may be useful in selected cases, especially when CNS involvement is suspected. In addition, multigated acquisition (MUGA) scan or echocardiogram is recommended for patients being considered for treatment regimens containing anthracyclines or anthracenediones. Hepatitis B virus (HBV) testing should be performed prior to initiation of treatment with immunotherapy (with or without chemotherapy) given the potential risks for viral reactivation with such regimens.

SOT and HCT recipients who develop PTLD have a higher EBV viral load than transplant recipients without PTLD.⁶⁶ Evaluation of EBV viral load by quantitative polymerase-chain-reaction (PCR) amplification of EBV DNA can aid in the diagnosis as well as monitoring of treatment responses in patients with PTLD.⁶⁶⁻⁶⁹ The reported positive and negative predictive



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

values for this approach have varied significantly for SOT (28% to 100% and 75% to 100%, respectively) and HCT (25% to 40% and 67% to 86%, respectively) recipients.⁷⁰⁻⁷² Plasma or peripheral blood mononuclear cells (PBMC) are useful for measuring EBV viral load. Although some studies have shown that viral load in plasma is more sensitive than PBMC, especially in patients with EBV-positive disease.^{68,73,74} In recent studies, cell-free plasma EBV DNA was a better marker than EBV DNA from peripheral blood mononuclear cells.^{70,71} EBV serology to assess primary infection versus reactivation may be useful. CMV infection has been associated with risks for EBV-positive PTLD.^{24,75} Thus, EBV PCR for the measurement of cell-free plasma EBV DNA marker and CMV PCR can be useful for selected patients.

Treatment

Treatment approaches are largely dependent on the PTLD subtype.⁷⁶ Published reports have included reduction in immunosuppression (RI), antiviral therapy, rituximab monotherapy, chemotherapy, and/or chemoimmunotherapy regimens. The optimal treatment for PTLD is not well defined due to the lack of randomized controlled trials and the heterogeneity of the disease.

The role of antiviral therapy is controversial since the majority of PTLD are associated with latent EBV. Replicating EBV DNA has been reported in about 40% of EBV associated lymphoproliferative disorders in immunocompromised patients.⁷⁷⁻⁷⁹ Antiviral drugs targeting EBV replication may be beneficial in the subset of patients with early or polymorphic PTLD.^{80,81}

Reduction in immunosuppression

RI remains the initial step in the management of nearly all cases of PTLD. RI leads to regression of PTLD in 20% to 80% of patients with polyclonal and monoclonal PTLD.⁸²⁻⁸⁴ EBV-negative disease is less responsive to RI, but responses have been reported.^{21,37} In a prospective phase II study that

evaluated a sequential approach (RI first, then interferon-alfa for less than complete response (CR), then multiagent chemotherapy if less than CR to interferon) in 16 eligible patients with PTLD following SOT, RI resulted in one partial response (PR).⁸⁵

Rituximab monotherapy

The efficacy of rituximab monotherapy in the treatment of patients with B-cell PTLD has been confirmed in phase II studies and retrospective analyses.^{43,86-91} In a prospective multicenter phase II study of 43 eligible patients with PTLD after SOT, rituximab monotherapy resulted in an overall response rate (ORR) of 44% (28% CR) with a 1-year overall survival (OS) rate of 67%.⁸⁷ A prospective multicenter phase II study of 38 patients with PTLD after SOT demonstrated that risk-adapted extended treatment with 4 additional doses of rituximab increased the CR rate from 34% to 61% without increasing toxicity.⁹² Among the patients who could not achieve a CR with rituximab monotherapy and subsequently received rituximab in combination with chemotherapy (R-CHOP or R-EPOCH; n=8), 6 patients achieved a CR (75%). At a median follow up of 28 months, the event-free survival (EFS) and OS rates were 42% and 47%, respectively.⁹²

The results of a multicenter retrospective analysis (80 patients with PTLD following SOT) suggests that the inclusion of rituximab as part of initial therapy significantly improved both progression-free survival (PFS) and OS.⁴³ All patients initially underwent RI, and 74% were treated with rituximab with or without chemotherapy. The 3-year PFS and OS rates for all patients were 57% and 62%, respectively. The 3-year PFS and OS rates were 70% and 73% respectively for patients who received rituximab-based therapy as part of initial treatment compared to 21% and 33%, respectively for those who did not receive rituximab-based therapy.

Chemotherapy with or without rituximab

Anthracycline-based chemotherapy with or without rituximab has also been effective in the treatment of patients with PTLD.^{88,93-97} In a



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

retrospective analysis of 26 patients with PTLD after SOT with disease unresponsive to RI alone, CHOP (cyclophosphamide, doxorubicin, vincristine and prednisone) induced an ORR of 65% (50% CR).⁹⁵ With a median follow up of nearly 9 years, the median OS was 14 months. Treatment-related mortality rate was high, at 31%.⁹⁵ Chemotherapy and RI, with or without rituximab has also been reported to induce durable CR with reduced risk of graft impairment when used as first line treatment.^{98,99}

Sequential Chemoimmunotherapy

The prospective multicenter phase II study (PTLD-1) demonstrated the safety and efficacy of sequential chemoimmunotherapy (4 weekly doses of rituximab followed by 4 cycles of CHOP-21) with G-CSF in patients with PTLD after failure with initial RI (n =74; 70 evaluable patients).¹⁰⁰ The large majority of patients presented with monomorphic PTLD (primarily DLBCL), and 44% were EBV positive. The ORR with rituximab was 60% (20% CR), which improved to 90% (68% CR) in the patients who received CHOP chemotherapy following rituximab. The 5-year PFS and OS rates were 50% and 55%, respectively.¹⁰⁰ The most common grade 3 or 4 toxicities included leukopenia (68%) and infectious events (41%). Treatment-related mortality associated with CHOP was reported in 11% of patients.

A risk-stratified treatment strategy based upon initial response to rituximab was evaluated in a subsequent prospective, international, multicenter phase II trial. In this trial, 152 patients with PTLD after SOT unresponsive to RI received induction therapy with 4 weekly doses of rituximab.¹⁰¹ Low-risk patients (defined as those achieving CR after initial rituximab) received consolidation with rituximab monotherapy on days 50, 72, 94, and 116. High-risk patients (defined as non-CR after initial rituximab) received chemoimmunotherapy with R-CHOP-21 (4 cycles) combined with G-CSF. *Pneumocystis jirovecii* prophylaxis was recommended. Among the 126 patients enrolled in the risk-stratified protocol, the ORR was 88% (70% CR). The estimated 3-year OS rate was 70%, which compared

favorably to the OS rate of 61%. The treatment related mortality was 8% and the median OS was 6.6 years. The estimated 3-year time-to-progression (TTP) was 89% for patients with low-risk group treated with rituximab consolidation and response to rituximab remained a prognostic factor for OS despite the risk stratification. This risk-stratified sequential treatment strategy obviates the need for chemoimmunotherapy for low-risk patients achieving a CR to rituximab, while incorporating a more effective chemoimmunotherapy regimen (R-CHOP) for high-risk patients.^{44,101}

Other indications for chemotherapy include specific histologic subtypes, such as PTCL, NOS, BL, CHL, and other uncommon lymphomas. These must be managed with treatment approaches associated with improved outcomes in the specific histologic subtypes. In a retrospective multiinstitutional analysis of 84 patients with primary CNS PTLD, 93% of patients received RI.¹⁰² Additional first-line therapy included high-dose methotrexate (48%), high-dose cytarabine (33%), brain radiotherapy (24%), and/or rituximab (44%). The ORR was 60%. At a median follow-up of 42 months, the 3-year PFS and OS rates were 32% and 43%, respectively.

Retransplantation after diagnosis and treatment of PTLD

Retransplantation after diagnosis and treatment of PTLD is feasible in selected patients with PTLD following SOT.^{83,103} Waiting for at least one year from the control of PTLD to retransplantation is recommended to minimize risk of PTLD recurrence.⁸³ In a French cohort study of 52 patients with kidney transplants who underwent 55 retransplantation, the median time elapsed from PTLD to retransplantation was 90 months and only one patient developed PTLD recurrence after retransplantation.¹⁰³

Adoptive immunotherapy with EBV-specific cytotoxic T-lymphocytes

Adoptive immunotherapy with autologous or allogeneic EBV-specific cytotoxic T-lymphocytes (EBV-CTL) may be a promising strategy in patients with PTLD after failure of conventional treatments.¹⁰⁴⁻¹⁰⁸ A



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

prospective multicenter phase II study evaluated allogeneic EBV-CTL in the treatment of 33 patients with PTLD that failed conventional therapy.¹⁰⁷ The majority of patients (94%) had received SOT. All patients had RI as part of initial therapy for PTLD, and some patients had also received rituximab monotherapy, anti-viral therapy, or chemotherapy. The ORR and OS rate at 6 months were 52% (42% CR) and 79%, respectively. In a study that evaluated the effectiveness of allogeneic EBV-CTL in 114 patients who underwent allogeneic HSCT, EBV-CTL prevented PTLD in 101 patients and induced a durable CR in 85% of patients in the subgroup with existing PTLD (n=13).¹⁰⁸ This study also showed that during long-term follow up, functional EBV-CTL persisted up to 9 years.

Further prospective studies are needed to better define the role of adoptive immunotherapy in the prevention and management of PTLD.

NCCN Recommendations

Treatment options for PTLD depend on the histological subtype and should be individualized. RI, if possible, should be a part of the initial treatment approach for all patients with PTLD.⁸²⁻⁸⁴ Initial management strategies include reduction of calcineurin inhibition (cyclosporin or tacrolimus) by 50% and discontinuation of antimetabolic agents (azathioprine or mycophenolate mofetil). Discontinuation of all non-steroidal immunosuppression should be considered in patients who are critically ill with extensive and life-threatening disease. The response to RI is 20% to 80% in polyclonal or monoclonal cases, and patients should be closely monitored during RI. Graft monitoring is essential to allow for early detection of allograft rejection. Importantly, RI should be initiated and managed in coordination with the transplant team in order to minimize risks for graft rejection. In contrast to the staging of lymphoma in immunocompetent patients, restaging should be performed at 2 to 4 weeks in patients receiving RI as the only treatment option, since responses occur very early. Additional treatment options are necessary

(as described below based on the subtype) for patients who have not achieved a CR or those with persistent or progressive disease after initial RI.

CHL PTLD should be managed as described in the NCCN Guidelines for Hodgkin Lymphoma. In a cohort study of 192 patients with CHL-like PTLD identified in the Scientific Registry of Transplant Recipients (SRTR), among the 145 patients treated with chemotherapy, most of the patients received regimens specific for CHL, and the use of CHL-specific chemotherapy was associated with improved OS and disease-specific survival.⁵⁸

Early lesions

RI alone is an appropriate first-line therapy for patients with early lesions. For patients who achieve a CR, re-escalation of immunosuppressive therapy should be individualized, taking into account the extent of initial RI and the nature of the organ allograft. These decisions should be made in conjunction with the transplant team. Graft organ function and EBV viral load should be monitored.

Rituximab is recommended as second-line therapy for patients with persistent or progressive disease after RI. EBV viral load should be monitored by PCR.

Monomorphic PTLD

Treatment options include RI and/or rituximab monotherapy^{43,86-91} or chemoimmunotherapy for patients with B-cell type.^{88,93-97} A risk stratified approach (as described above) could be used for patients achieving CR to rituximab monotherapy.^{44,92,100,101} Rituximab monotherapy should only be considered as part of a step-wise approach to treatment in patients who are not highly symptomatic or in those who cannot tolerate chemoimmunotherapy due to the presence of comorbid conditions. In addition, RI is effective in plasmacytic PTLD with responses ranging from



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

33% to 75% in small series.^{51,53,54} Patients who achieve a CR with initial therapy should undergo surveillance/follow up according to the NCCN Guidelines for B-Cell Lymphoma specific for the B-cell lymphoma subtype.

Second-line therapy options for patients with persistent or progressive disease are dependent on initial therapy. Rituximab or chemoimmunotherapy are recommended for patients who received RI alone as initial treatment, whereas patients who received rituximab monotherapy should be treated with chemoimmunotherapy. Patients who received chemoimmunotherapy as initial treatment should be managed as described for relapsed or refractory DLBCL. Other options include participation in a suitable clinical trial, if available, or adoptive immunotherapy with EBV-specific cytotoxic T-lymphocytes (if EBV positive).¹⁰⁴⁻¹⁰⁸ Future strategies and other therapies have been reviewed.¹

There are no established treatment options (other than RI) for patients with T-cell subtype. Treatment with anthracycline-based multiagent chemotherapy regimens recommended for T-cell lymphomas could be considered.

Polymorphic PTLD

RI along with rituximab monotherapy^{43,86-91} or chemoimmunotherapy^{88,93-97} are recommended for patients with systemic disease. RI along with involved-site radiation therapy (ISRT) with or without rituximab or surgery with or without rituximab, or rituximab monotherapy is recommended for patients with localized disease. A risk stratified approach (as described above) could be used for patients achieving CR to rituximab monotherapy.^{44,92,100,101}

Observation or continuation of RI with or without rituximab maintenance is recommended for patients who achieve a CR. Chemoimmunotherapy or adoptive immunotherapy with EBV-specific cytotoxic T-lymphocytes (if EBV positive) are included as options for patients with persistent or

progressive disease.¹⁰⁴⁻¹⁰⁸ Participation in a suitable clinical trial, where available, should also be considered.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

References

1. Dierickx D, Habermann TM. Post-Transplantation Lymphoproliferative Disorders in Adults. *N Engl J Med* 2018;378:549-562. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29414277>.
2. Chadburn A, Suci-Foca N, Cesarman E, et al. Post-transplantation lymphoproliferative disorders arising in solid organ transplant recipients are usually of recipient origin. *Am J Pathol* 1995;147:1862-1870. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7495309>.
3. Weissmann DJ, Ferry JA, Harris NL, et al. Posttransplantation lymphoproliferative disorders in solid organ recipients are predominantly aggressive tumors of host origin. *Am J Clin Pathol* 1995;103:748-755. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7785662>.
4. Micallef IN, Chhanabhai M, Gascoyne RD, et al. Lymphoproliferative disorders following allogeneic bone marrow transplantation: the Vancouver experience. *Bone Marrow Transplant* 1998;22:981-987. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9849695>.
5. Curtis RE, Travis LB, Rowlings PA, et al. Risk of lymphoproliferative disorders after bone marrow transplantation: a multi-institutional study. *Blood* 1999;94:2208-2216. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10498590>.
6. Petit B, Le Meur Y, Jaccard A, et al. Influence of host-recipient origin on clinical aspects of posttransplantation lymphoproliferative disorders in kidney transplantation. *Transplantation* 2002;73:265-271. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11821742>.
7. Gulley ML, Swinnen LJ, Plaisance KT, Jr., et al. Tumor origin and CD20 expression in posttransplant lymphoproliferative disorder occurring in solid organ transplant recipients: implications for immune-based therapy. *Transplantation* 2003;76:959-964. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14508361>.
8. Peterson MR, Emery SC, Yung GL, et al. Epstein-Barr virus-associated posttransplantation lymphoproliferative disorder following lung transplantation is more commonly of host origin. *Arch Pathol Lab Med* 2006;130:176-180. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16454557>.
9. Kinch A, Cavelier L, Bengtsson M, et al. Donor or recipient origin of posttransplant lymphoproliferative disorders following solid organ transplantation. *Am J Transplant* 2014;14:2838-2845. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25307322>.
10. Leblond V, Davi F, Charlotte F, et al. Posttransplant lymphoproliferative disorders not associated with Epstein-Barr virus: a distinct entity? *J Clin Oncol* 1998;16:2052-2059. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9626203>.
11. Nelson BP, Nalesnik MA, Bahler DW, et al. Epstein-Barr virus-negative post-transplant lymphoproliferative disorders: a distinct entity? *Am J Surg Pathol* 2000;24:375-385. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10716151>.
12. Ghobrial IM, Habermann TM, Macon WR, et al. Differences between early and late posttransplant lymphoproliferative disorders in solid organ transplant patients: are they two different diseases? *Transplantation* 2005;79:244-247. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15665775>.
13. Luskin MR, Heil DS, Tan KS, et al. The Impact of EBV Status on Characteristics and Outcomes of Posttransplantation Lymphoproliferative Disorder. *Am J Transplant* 2015;15:2665-2673. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25988622>.
14. Quinlan SC, Pfeiffer RM, Morton LM, Engels EA. Risk factors for early-onset and late-onset post-transplant lymphoproliferative disorder in kidney recipients in the United States. *Am J Hematol* 2011;86:206-209. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21264909>.
15. Schober T, Framke T, Kreipe H, et al. Characteristics of early and late PTLD development in pediatric solid organ transplant recipients. *Transplantation* 2013;95:240-246. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23222898>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

16. Craig FE, Johnson LR, Harvey SA, et al. Gene expression profiling of Epstein-Barr virus-positive and -negative monomorphic B-cell posttransplant lymphoproliferative disorders. *Diagn Mol Pathol* 2007;16:158-168. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17721324>.

17. Morscio J, Dierickx D, Ferreiro JF, et al. Gene Expression Profiling Reveals Clear Differences Between EBV-Positive and EBV-Negative Posttransplant Lymphoproliferative Disorders. *Am J Transplant* 2013. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23489474>.

18. Ferreiro JF, Morscio J, Dierickx D, et al. EBV-Positive and EBV-Negative Posttransplant Diffuse Large B Cell Lymphomas Have Distinct Genomic and Transcriptomic Features. *Am J Transplant* 2016;16:414-425. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26780579>.

19. Margolskee E, Jobanputra V, Jain P, et al. Genetic landscape of T- and NK-cell post-transplant lymphoproliferative disorders. *Oncotarget* 2016;7:37636-37648. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27203213>.

20. Menter T, Juskevicius D, Alikian M, et al. Mutational landscape of B-cell post-transplant lymphoproliferative disorders. *Br J Haematol* 2017;178:48-56. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28419429>.

21. Courville EL, Yohe S, Chou D, et al. EBV-negative monomorphic B-cell post-transplant lymphoproliferative disorders are pathologically distinct from EBV-positive cases and frequently contain TP53 mutations. *Mod Pathol* 2016;29:1200-1211. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27443517>.

22. U.S. National Library of Medicine Key MEDLINE® Indicators Available at: http://www.nlm.nih.gov/bsd/bsd_key.html.

23. Swinnen LJ, Costanzo-Nordin MR, Fisher SG, et al. Increased incidence of lymphoproliferative disorder after immunosuppression with the monoclonal antibody OKT3 in cardiac-transplant recipients. *N Engl J*

Med 1990;323:1723-1728. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2100991>.

24. Walker RC, Marshall WF, Strickler JG, et al. Pretransplantation assessment of the risk of lymphoproliferative disorder. *Clin Infect Dis* 1995;20:1346-1353. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7620022>.

25. Caillard S, Dharnidharka V, Agodoa L, et al. Posttransplant lymphoproliferative disorders after renal transplantation in the United States in era of modern immunosuppression. *Transplantation* 2005;80:1233-1243. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16314791>.

26. Caillard S, Lelong C, Pessione F, Moulin B. Post-transplant lymphoproliferative disorders occurring after renal transplantation in adults: report of 230 cases from the French Registry. *Am J Transplant* 2006;6:2735-2742. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17049061>.

27. Knight JS, Tsodikov A, Cibrik DM, et al. Lymphoma after solid organ transplantation: risk, response to therapy, and survival at a transplantation center. *J Clin Oncol* 2009;27:3354-3362. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19451438>.

28. Caillard S, Lamy FX, Quelen C, et al. Epidemiology of posttransplant lymphoproliferative disorders in adult kidney and kidney pancreas recipients: report of the French registry and analysis of subgroups of lymphomas. *Am J Transplant* 2012;12:682-693. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22226336>.

29. Rausch L, Koenecke C, Koch HF, et al. Matched-pair analysis: identification of factors with independent influence on the development of PTLN after kidney or liver transplantation. *Transplant Res* 2016;5:6. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27486513>.

30. Francis A, Johnson DW, Teixeira-Pinto A, et al. Incidence and predictors of post-transplant lymphoproliferative disease after kidney transplantation during adulthood and childhood: a registry study. *Nephrol*



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

Dial Transplant 2018;33:881-889. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/29342279>.

31. Nalesnik M, Jaffe R, Reyes J, et al. Posttransplant lymphoproliferative disorders in small bowel allograft recipients. Transplant Proc 2000;32:1213. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/10995914>.

32. Opelz G, Dohler B. Lymphomas after solid organ transplantation: a collaborative transplant study report. Am J Transplant 2004;4:222-230. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14974943>.

33. Di Benedetto F, Lauro A, Masetti M, et al. Outcomes after adult isolated small bowel transplantation: experience from a single European centre. Dig Liver Dis 2005;37:240-246. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15788207>.

34. Landgren O, Gilbert ES, Rizzo JD, et al. Risk factors for lymphoproliferative disorders after allogeneic hematopoietic cell transplantation. Blood 2009;113:4992-5001. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19264919>.

35. Wudhikarn K, Holman CJ, Linan M, et al. Post-transplant lymphoproliferative disorders in lung transplant recipients: 20-yr experience at the University of Minnesota. Clin Transplant 2011;25:705-713. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/21077949>.

36. Kremer BE, Reshef R, Misleh JG, et al. Post-transplant lymphoproliferative disorder after lung transplantation: a review of 35 cases. J Heart Lung Transplant 2012;31:296-304. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22112992>.

37. Dierickx D, Tousseyn T, Sagaert X, et al. Single-center analysis of biopsy-confirmed posttransplant lymphoproliferative disorder: incidence, clinicopathological characteristics and prognostic factors. Leuk Lymphoma 2013;54:2433-2440. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23442063>.

38. Asleh R, Li X, Alnsasra H, et al. Long-Term Sirolimus for Primary Immunosuppression and Incidence of De Novo Malignancy Following Heart Transplantation. The Journal of Heart and Lung Transplantation 2019;38:S274. Available at: <https://doi.org/10.1016/j.healun.2019.01.682>.

39. Leblond V, Dhedin N, Mamzer Bruneel MF, et al. Identification of prognostic factors in 61 patients with posttransplantation lymphoproliferative disorders. J Clin Oncol 2001;19:772-778. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11157030>.

40. Ghobrial IM, Habermann TM, Ristow KM, et al. Prognostic factors in patients with post-transplant lymphoproliferative disorders (PTLD) in the rituximab era. Leuk Lymphoma 2005;46:191-196. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15621801>.

41. Oton AB, Wang H, Leleu X, et al. Clinical and pathological prognostic markers for survival in adult patients with post-transplant lymphoproliferative disorders in solid transplant. Leuk Lymphoma 2008;49:1738-1744. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18798108>.

42. Caillard S, Porcher R, Provot F, et al. Post-transplantation lymphoproliferative disorder after kidney transplantation: report of a nationwide French registry and the development of a new prognostic score. J Clin Oncol 2013;31:1302-1309. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23423742>.

43. Evens AM, David KA, Helenowski I, et al. Multicenter analysis of 80 solid organ transplantation recipients with post-transplantation lymphoproliferative disease: outcomes and prognostic factors in the modern era. J Clin Oncol 2010;28:1038-1046. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/20085936>.

44. Trappe RU, Choquet S, Dierickx D, et al. International prognostic index, type of transplant and response to rituximab are key parameters to tailor treatment in adults with CD20-positive B cell PTLD: clues from the PTLD-1 trial. Am J Transplant 2015;15:1091-1100. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25736912>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

45. Swerdlow SH, Harris NL, Jaffe ES, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Revised 4th ed. Lyon, France: IARC; 2017.

46. Tsao L, Hsi ED. The clinicopathologic spectrum of posttransplantation lymphoproliferative disorders. Arch Pathol Lab Med 2007;131:1209-1218. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17683183>.

47. Xicoy B, Ribera JM, Esteve J, et al. Post-transplant Burkitt's leukemia or lymphoma. Study of five cases treated with specific intensive therapy (PETHEMA ALL-3/97 trial). Leuk Lymphoma 2003;44:1541-1543. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14565657>.

48. Picarsic J, Jaffe R, Mazariegos G, et al. Post-transplant Burkitt lymphoma is a more aggressive and distinct form of post-transplant lymphoproliferative disorder. Cancer 2011;117:4540-4550. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21446044>.

49. Zimmermann H, Reinke P, Neuhaus R, et al. Burkitt post-transplantation lymphoma in adult solid organ transplant recipients: sequential immunochemotherapy with rituximab (R) followed by cyclophosphamide, doxorubicin, vincristine, and prednisone (CHOP) or R-CHOP is safe and effective in an analysis of 8 patients. Cancer 2012;118:4715-4724. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22392525>.

50. Ferreiro JF, Morscio J, Dierickx D, et al. Post-transplant molecularly defined Burkitt lymphomas are frequently MYC-negative and characterized by the 11q-gain/loss pattern. Haematologica 2015;100:e275-279. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25795716>.

51. Richendollar BG, Hsi ED, Cook JR. Extramedullary plasmacytoma-like posttransplantation lymphoproliferative disorders: clinical and pathologic features. Am J Clin Pathol 2009;132:581-588. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19762536>.

52. Trappe R, Zimmermann H, Fink S, et al. Plasmacytoma-like post-transplant lymphoproliferative disorder, a rare subtype of monomorphic B-cell post-transplant lymphoproliferation, is associated with

a favorable outcome in localized as well as in advanced disease: a prospective analysis of 8 cases. Haematologica 2011;96:1067-1071. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21719885>.

53. Zimmermann H, Oschlies I, Fink S, et al. Plasmablastic posttransplant lymphoma: cytogenetic aberrations and lack of Epstein-Barr virus association linked with poor outcome in the prospective German Posttransplant Lymphoproliferative Disorder Registry. Transplantation 2012;93:543-550. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22234349>.

54. Karuturi M, Shah N, Frank D, et al. Plasmacytic post-transplant lymphoproliferative disorder: a case series of nine patients. Transpl Int 2013;26:616-622. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23551167>.

55. Kormann R, Francois H, Moles T, et al. Plasma cell neoplasia after kidney transplantation: French cohort series and review of the literature. PLoS One 2017;12:e0179406. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28636627>.

56. Berg LC, Copenhaver CM, Morrison VA, et al. B-cell lymphoproliferative disorders in solid-organ transplant patients: detection of Epstein-Barr virus by in situ hybridization. Hum Pathol 1992;23:159-163. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1310951>.

57. Herreman A, Dierickx D, Morscio J, et al. Clinicopathological characteristics of posttransplant lymphoproliferative disorders of T-cell origin: single-center series of nine cases and meta-analysis of 147 reported cases. Leuk Lymphoma 2013;54:2190-2199. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23402267>.

58. Rosenberg AS, Klein AK, Ruthazer R, Evens AM. Hodgkin lymphoma post-transplant lymphoproliferative disorder: A comparative analysis of clinical characteristics, prognosis, and survival. Am J Hematol 2016;91:560-565. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26928381>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

59. Harris NL, Ferry JA, Swerdlow SH. Posttransplant lymphoproliferative disorders: summary of Society for Hematopathology Workshop. Semin Diagn Pathol 1997;14:8-14. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9044505>.

60. Parker A, Bowles K, Bradley JA, et al. Diagnosis of post-transplant lymphoproliferative disorder in solid organ transplant recipients - BCSH and BTS Guidelines. Br J Haematol 2010;149:675-692. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20408847>.

61. Capello D, Cerri M, Muti G, et al. Molecular histogenesis of posttransplantation lymphoproliferative disorders. Blood 2003;102:3775-3785. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12907442>.

62. Capello D, Rossi D, Gaidano G. Post-transplant lymphoproliferative disorders: molecular basis of disease histogenesis and pathogenesis. Hematol Oncol 2005;23:61-67. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16216037>.

63. Capello D, Cerri M, Muti G, et al. Analysis of immunoglobulin heavy and light chain variable genes in post-transplant lymphoproliferative disorders. Hematol Oncol 2006;24:212-219. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16897790>.

64. Knowles DM, Cesarman E, Chadburn A, et al. Correlative morphologic and molecular genetic analysis demonstrates three distinct categories of posttransplantation lymphoproliferative disorders. Blood 1995;85:552-565. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7812011>.

65. Cesarman E, Chadburn A, Liu YF, et al. BCL-6 gene mutations in posttransplantation lymphoproliferative disorders predict response to therapy and clinical outcome. Blood 1998;92:2294-2302. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9746767>.

66. van Esser JW, van der Holt B, Meijer E, et al. Epstein-Barr virus (EBV) reactivation is a frequent event after allogeneic stem cell transplantation (SCT) and quantitatively predicts EBV-lymphoproliferative disease

following T-cell-depleted SCT. Blood 2001;98:972-978. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11493441>.

67. Stevens SJ, Verschuuren EA, Pronk I, et al. Frequent monitoring of Epstein-Barr virus DNA load in unfractionated whole blood is essential for early detection of posttransplant lymphoproliferative disease in high-risk patients. Blood 2001;97:1165-1171. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11222357>.

68. Tsai DE, Douglas L, Andreadis C, et al. EBV PCR in the diagnosis and monitoring of posttransplant lymphoproliferative disorder: results of a two-arm prospective trial. Am J Transplant 2008;8:1016-1024. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18312608>.

69. Omar H, Hagglund H, Gustafsson-Jernberg A, et al. Targeted monitoring of patients at high risk of post-transplant lymphoproliferative disease by quantitative Epstein-Barr virus polymerase chain reaction. Transpl Infect Dis 2009;11:393-399. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19497070>.

70. Kanakry JA, Hegde AM, Durand CM, et al. The clinical significance of EBV DNA in the plasma and peripheral blood mononuclear cells of patients with or without EBV diseases. Blood 2016;127:2007-2017. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26744460>.

71. Semenova T, Lupo J, Alain S, et al. Multicenter Evaluation of Whole-Blood Epstein-Barr Viral Load Standardization Using the WHO International Standard. J Clin Microbiol 2016;54:1746-1750. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27076661>.

72. Styczynski J, van der Velden W, Fox CP, et al. Management of Epstein-Barr Virus infections and post-transplant lymphoproliferative disorders in patients after allogeneic hematopoietic stem cell transplantation: Sixth European Conference on Infections in Leukemia (ECIL-6) guidelines. Haematologica 2016;101:803-811. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27365460>.

73. Wagner HJ, Fischer L, Jabs WJ, et al. Longitudinal analysis of Epstein-Barr viral load in plasma and peripheral blood mononuclear cells



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

of transplanted patients by real-time polymerase chain reaction.

Transplantation 2002;74:656-664. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/12352882>.

74. Ryan JL, Fan H, Swinnen LJ, et al. Epstein-Barr Virus (EBV) DNA in plasma is not encapsidated in patients with EBV-related malignancies.

Diagn Mol Pathol 2004;13:61-68. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15167006>.

75. Manez R, Breinig MC, Linden P, et al. Posttransplant lymphoproliferative disease in primary Epstein-Barr virus infection after liver transplantation: the role of cytomegalovirus disease. J Infect Dis 1997;176:1462-1467. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9395355>.

76. Muti G, Cantoni S, Oreste P, et al. Post-transplant lymphoproliferative disorders: improved outcome after clinico-pathologically tailored treatment. Haematologica 2002;87:67-77. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11801467>.

77. Katz BZ, Raab-Traub N, Miller G. Latent and replicating forms of Epstein-Barr virus DNA in lymphomas and lymphoproliferative diseases. J Infect Dis 1989;160:589-598. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/2551973>.

78. Kroll J, Li S, Levi M, Weinberg A. Lytic and latent EBV gene expression in transplant recipients with and without post-transplant lymphoproliferative disorder. J Clin Virol 2011;52:231-235. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21900040>.

79. Gonzalez-Farre B, Rovira J, Martinez D, et al. In vivo intratumoral Epstein-Barr virus replication is associated with XBP1 activation and early-onset post-transplant lymphoproliferative disorders with prognostic implications. Mod Pathol 2014;27:1599-1611. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24762547>.

80. Hanto DW, Frizzera G, Gajl-Peczalska KJ, et al. Epstein-Barr virus-induced B-cell lymphoma after renal transplantation: acyclovir therapy and transition from polyclonal to monoclonal B-cell proliferation. N

Engl J Med 1982;306:913-918. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/6278307>.

81. Birkeland SA, Andersen HK, Hamilton-Dutoit SJ. Preventing acute rejection, Epstein-Barr virus infection, and posttransplant lymphoproliferative disorders after kidney transplantation: use of aciclovir and mycophenolate mofetil in a steroid-free immunosuppressive protocol. Transplantation 1999;67:1209-1214. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/10342310>.

82. Tsai DE, Hardy CL, Tomaszewski JE, et al. Reduction in immunosuppression as initial therapy for posttransplant lymphoproliferative disorder: analysis of prognostic variables and long-term follow-up of 42 adult patients. Transplantation 2001;71:1076-1088. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11374406>.

83. Parker A, Bowles K, Bradley JA, et al. Management of post-transplant lymphoproliferative disorder in adult solid organ transplant recipients - BCSH and BTS Guidelines. Br J Haematol 2010;149:693-705. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20408848>.

84. Reshef R, Vardhanabhuti S, Luskin MR, et al. Reduction of immunosuppression as initial therapy for posttransplantation lymphoproliferative disorder. Am J Transplant 2011;11:336-347. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21219573>.

85. Swinnen LJ, LeBlanc M, Grogan TM, et al. Prospective study of sequential reduction in immunosuppression, interferon alpha-2B, and chemotherapy for posttransplantation lymphoproliferative disorder. Transplantation 2008;86:215-222. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18645482>.

86. Blaes AH, Peterson BA, Bartlett N, et al. Rituximab therapy is effective for posttransplant lymphoproliferative disorders after solid organ transplantation: results of a phase II trial. Cancer 2005;104:1661-1667. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16149091>.



NCCN Guidelines Version 2.2020

B-Cell Lymphomas

87. Choquet S, Leblond V, Herbrecht R, et al. Efficacy and safety of rituximab in B-cell post-transplantation lymphoproliferative disorders: results of a prospective multicenter phase 2 study. *Blood* 2006;107:3053-3057. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16254143>.

88. Elstrom RL, Andreadis C, Aqui NA, et al. Treatment of PTLT with rituximab or chemotherapy. *Am J Transplant* 2006;6:569-576. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16468968>.

89. Jain AB, Marcos A, Pokharna R, et al. Rituximab (chimeric anti-CD20 antibody) for posttransplant lymphoproliferative disorder after solid organ transplantation in adults: long-term experience from a single center. *Transplantation* 2005;80:1692-1698. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16378063>.

90. Milpied N, Vasseur B, Parquet N, et al. Humanized anti-CD20 monoclonal antibody (Rituximab) in post transplant B-lymphoproliferative disorder: a retrospective analysis on 32 patients. *Ann Oncol* 2000;11 Suppl 1:113-116. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10707791>.

91. Oertel SHK, Verschuuren E, Reinke P, et al. Effect of anti-CD 20 antibody rituximab in patients with post-transplant lymphoproliferative disorder (PTLD). *Am J Transplant* 2005;5:2901-2906. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16303003>.

92. Gonzalez-Barca E, Domingo-Domenech E, Capote FJ, et al. Prospective phase II trial of extended treatment with rituximab in patients with B-cell post-transplant lymphoproliferative disease. *Haematologica* 2007;92:1489-1494. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18024397>.

93. Buadi FK, Heyman MR, Gocke CD, et al. Treatment and outcomes of post-transplant lymphoproliferative disease: a single institution study. *Am J Hematol* 2007;82:208-214. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17022049>.

94. Buell JF, Gross TG, Hanaway MJ, et al. Chemotherapy for posttransplant lymphoproliferative disorder: the Israel Penn International Transplant Tumor Registry experience. *Transplant Proc* 2005;37:956-957. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15848588>.

95. Choquet S, Trappe R, Leblond V, et al. CHOP-21 for the treatment of post-transplant lymphoproliferative disorders (PTLD) following solid organ transplantation. *Haematologica* 2007;92:273-274. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17296588>.

96. Fohrer C, Caillard S, Koumarianou A, et al. Long-term survival in post-transplant lymphoproliferative disorders with a dose-adjusted ACVBP regimen. *Br J Haematol* 2006;134:602-612. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16889621>.

97. Orjuela M, Gross TG, Cheung Y-K, et al. A pilot study of chemoimmunotherapy (cyclophosphamide, prednisone, and rituximab) in patients with post-transplant lymphoproliferative disorder following solid organ transplantation. *Clin Cancer Res* 2003;9:52. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14506193>.

98. Taylor AL, Bowles KM, Callaghan CJ, et al. Anthracycline-based chemotherapy as first-line treatment in adults with malignant posttransplant lymphoproliferative disorder after solid organ transplantation. *Transplantation* 2006;82:375-381. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16906036>.

99. Trappe R, Hinrichs C, Appel U, et al. Treatment of PTLT with rituximab and CHOP reduces the risk of renal graft impairment after reduction of immunosuppression. *Am J Transplant* 2009;9:2331-2337. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19663889>.

100. Trappe R, Oertel S, Leblond V, et al. Sequential treatment with rituximab followed by CHOP chemotherapy in adult B-cell post-transplant lymphoproliferative disorder (PTLD): the prospective international multicentre phase 2 PTLT-1 trial. *Lancet Oncol* 2012;13:196-206. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22173060>.



NCCN Guidelines Version 2.2020 B-Cell Lymphomas

101. Trappe RU, Dierickx D, Zimmermann H, et al. Response to Rituximab Induction Is a Predictive Marker in B-Cell Post-Transplant Lymphoproliferative Disorder and Allows Successful Stratification Into Rituximab or R-CHOP Consolidation in an International, Prospective, Multicenter Phase II Trial. *J Clin Oncol* 2017;35:536-543. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27992268>.

102. Evens AM, Choquet S, Kroll-Desrosiers AR, et al. Primary CNS posttransplant lymphoproliferative disease (PTLD): an international report of 84 cases in the modern era. *Am J Transplant* 2013;13:1512-1522. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23721553>.

103. Caillard S, Cellot E, Dantal J, et al. A French Cohort Study of Kidney Retransplantation after Post-Transplant Lymphoproliferative Disorders. *Clin J Am Soc Nephrol* 2017;12:1663-1670. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28818847>.

104. Rooney CM, Smith CA, Ng CY, et al. Infusion of cytotoxic T cells for the prevention and treatment of Epstein-Barr virus-induced lymphoma in allogeneic transplant recipients. *Blood* 1998;92:1549-1555. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9716582>.

105. Comoli P, Labirio M, Basso S, et al. Infusion of autologous Epstein-Barr virus (EBV)-specific cytotoxic T cells for prevention of EBV-related lymphoproliferative disorder in solid organ transplant recipients with evidence of active virus replication. *Blood* 2002;99:2592-2598. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11895798>.

106. Savoldo B, Goss JA, Hammer MM, et al. Treatment of solid organ transplant recipients with autologous Epstein Barr virus-specific cytotoxic T lymphocytes (CTLs). *Blood* 2006;108:2942-2949. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16835376>.

107. Haque T, Wilkie GM, Jones MM, et al. Allogeneic cytotoxic T-cell therapy for EBV-positive posttransplantation lymphoproliferative disease: results of a phase 2 multicenter clinical trial. *Blood* 2007;110:1123-1131. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17468341>.

108. Heslop HE, Slobod KS, Pule MA, et al. Long-term outcome of EBV-specific T-cell infusions to prevent or treat EBV-related lymphoproliferative disease in transplant recipients. *Blood* 2010;115:925-935. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19880495>.